

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/04520 A2

(51) International Patent Classification⁷: **C07K 14/705**

(21) International Application Number: PCT/US01/21448

(22) International Filing Date: 5 July 2001 (05.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/216,547	7 July 2000 (07.07.2000)	US
60/218,232	14 July 2000 (14.07.2000)	US
60/220,112	21 July 2000 (21.07.2000)	US
60/221,839	28 July 2000 (28.07.2000)	US

(71) Applicant (for all designated States except US): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **RAUMANN, Brigitte, E.** [US/US]; 5801 South Dorchester Avenue #3B, Chicago, IL 60637 (US). **THORNTON, Michael** [US/US]; 9 Medway Road, Woodside, CA 94062 (US). **DING, Li** [CN/US]; 3353 Alma Street #146, Palo Alto, CA 94306 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **HARLAND, Lee** [GB/GB]; 18 Chaucer Court, New Dover Road, Canterbury, Kent CT1 3AU (GB). **BURFORD, Neil** [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **GREENE, Barrie, D.** [US/US]; 1332 10th Avenue #104, San Francisco, CA 94122 (US). **SANJANWALA, Madhu, S.** [US/US]; 210 Sylvia Court, Los Altos, CA 94024 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **YAO, Monique, G.** [US/US]; 111 Frederick Court, Mountain View, CA 94043 (US). **YANG, Junming** [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). **PATTERSON, Chandra** [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). **GANDHI, Ameena, R.** [US/US]; 837 Roble Avenue #1, Menlo Park, CA 94025 (US). **HAFALIA, April, J., A.** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). **TRIBOULEY, Catherine, M.** [FR/US]; 1121 Tennessee Street #5, San Francisco, CA 94107 (US). **WALIA, Narinder, K.** [US/US]; 890 Davis Street #205, San Leandro, CA 94577 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). **WALSH,**

Roderick, T. [IE/GB]; 8 Boundary Court, St. Lawrence Road, Canterbury, Kent CT1 3EZ (GB). **RAMKUMAR, Jayalaxmi** [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). **LU, Yan** [CN/US]; 3885 Corrina Way, Palo Alto, CA 94303 (US). **LU, Dyung, Aina, M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **AZIMZAI, Yalda** [US/US]; 5518 Boulder Canyon Drive, Castro Valley, CA 94552 (US). **LAL, Preeti** [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). **ELLIOTT, Vicki, S.** [US/US]; 3770 Polton Place Way, San Jose, CA 95121 (US). **NGUYEN, Ganniel, B.** [US/US]; 1403 Ridgewood Drive, San Jose, CA 95118 (US). **XU, Yuming** [US/US]; 1739 Walnut Drive, Mountain View, CA 94040 (US). **SEILHAMER, Jeffrey, J.** [US/US]; 12555 La Cresta, Los Altos Hills, CA 94022 (US). **BOROWSKY, Mark, L.** [US/US]; 122 Orchard Avenue, Redwood City, CA 94061 (US). **KHAN, Farrah, A.** [IN/US]; 3617 Central Road #102, Glenview, IL 60025 (US). **KEARNEY, Liam** [US/US]; 50 Woodside Avenue, San Francisco, CA 94127 (US). **THANGAVELU, Kavitha** [IN/US]; 1950 Montecito Avenue #23, Mountain View, CA 94043 (US). **DAS, Debopriya** [IN/US]; Apartment 13, 1179 Bonita Avenue, Mountain View, CA 94040 (US). **POLICKY, Jennifer, L.** [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRANSPORTERS AND ION CHANNELS

(57) Abstract: The invention provides human transporters and ion channels (TRICH) and polynucleotides which identify and encode TRICH. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of TRICH.

WO 02/04520 A2

TRANSPORTERS AND ION CHANNELS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of transporters and ion channels and to the use of these sequences in the diagnosis, treatment, and prevention of transport, neurological, muscle, immunological, and cell proliferative disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of transporters and ion channels.

BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded and subdivided into functionally distinct organelles by hydrophobic lipid bilayer membranes which are highly impermeable to most polar molecules. Cells and organelles require transport proteins to import and export essential nutrients and metal ions including K^+ , NH_4^+ , P_i , SO_4^{2-} , sugars, and vitamins, as well as various metabolic waste products. Transport proteins also play roles in antibiotic resistance, toxin secretion, ion balance, synaptic neurotransmission, kidney function, intestinal absorption, tumor growth, and other diverse cell functions (Griffith, J. and C. Sansom (1998) The Transporter Facts Book, Academic Press, San Diego CA, pp. 3-29). Transport can occur by a passive concentration-dependent mechanism, or can be linked to an energy source such as ATP hydrolysis or an ion gradient. Proteins that function in transport include carrier proteins, which bind to a specific solute and undergo a conformational change that translocates the bound solute across the membrane, and channel proteins, which form hydrophilic pores that allow specific solutes to diffuse through the membrane down an electrochemical solute gradient.

Carrier proteins which transport a single solute from one side of the membrane to the other are called uniporters. In contrast, coupled transporters link the transfer of one solute with simultaneous or sequential transfer of a second solute, either in the same direction (symport) or in the opposite direction (antiport). For example, intestinal and kidney epithelium contains a variety of symporter systems driven by the sodium gradient that exists across the plasma membrane. Sodium moves into the cell down its electrochemical gradient and brings the solute into the cell with it. The sodium gradient that provides the driving force for solute uptake is maintained by the ubiquitous Na^+/K^+ ATPase system. Sodium-coupled transporters include the mammalian glucose transporter (SGLT1), iodide transporter (NIS), and multivitamin transporter (SMVT). All three transporters have twelve putative transmembrane segments, extracellular glycosylation sites, and cytoplasmically-oriented N- and C-termini. NIS plays a crucial role in the evaluation, diagnosis, and treatment of various thyroid pathologies because it is the molecular basis for radioiodide thyroid-imaging

techniques and for specific targeting of radioisotopes to the thyroid gland (Levy, O. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:5568-5573). SMVT is expressed in the intestinal mucosa, kidney, and placenta, and is implicated in the transport of the water-soluble vitamins, e.g., biotin and pantothenate (Prasad, P.D. et al. (1998) *J. Biol. Chem.* 273:7501-7506).

5 One of the largest families of transporters is the major facilitator superfamily (MFS), also called the uniporter-symporter-antiporter family. MFS transporters are single polypeptide carriers that transport small solutes in response to ion gradients. Members of the MFS are found in all classes of living organisms, and include transporters for sugars, oligosaccharides, phosphates, nitrates, nucleosides, monocarboxylates, and drugs. MFS transporters found in eukaryotes all have a structure
10 comprising 12 transmembrane segments (Pao, S.S. et al. (1998) *Microbiol. Molec. Biol. Rev.* 62:1-34). The largest family of MFS transporters is the sugar transporter family, which includes the seven glucose transporters (GLUT1-GLUT7) found in humans that are required for the transport of glucose and other hexose sugars. These glucose transport proteins have unique tissue distributions and physiological functions. GLUT1 provides many cell types with their basal glucose requirements and
15 transports glucose across epithelial and endothelial barrier tissues; GLUT2 facilitates glucose uptake or efflux from the liver; GLUT3 regulates glucose supply to neurons; GLUT4 is responsible for insulin-regulated glucose disposal; and GLUT5 regulates fructose uptake into skeletal muscle. Defects in glucose transporters are involved in a recently identified neurological syndrome causing infantile seizures and developmental delay, as well as glycogen storage disease, Fanconi-Bickel
20 syndrome, and non-insulin-dependent diabetes mellitus (Mueckler, M. (1994) *Eur. J. Biochem.* 219:713-725; Longo, N. and L.J. Elsas (1998) *Adv. Pediatr.* 45:293-313).

Monocarboxylate anion transporters are proton-coupled symporters with a broad substrate specificity that includes L-lactate, pyruvate, and the ketone bodies acetate, acetoacetate, and beta-hydroxybutyrate. At least seven isoforms have been identified to date. The isoforms are predicted
25 to have twelve transmembrane (TM) helical domains with a large intracellular loop between TM6 and TM7, and play a critical role in maintaining intracellular pH by removing the protons that are produced stoichiometrically with lactate during glycolysis. The best characterized H⁺-monocarboxylate transporter is that of the erythrocyte membrane, which transports L-lactate and a wide range of other aliphatic monocarboxylates. Other cells possess H⁺-linked monocarboxylate transporters with differing
30 substrate and inhibitor selectivities. In particular, cardiac muscle and tumor cells have transporters that differ in their K_m values for certain substrates, including stereoselectivity for L- over D-lactate, and in their sensitivity to inhibitors. There are Na⁺-monocarboxylate cotransporters on the luminal surface of intestinal and kidney epithelia, which allow the uptake of lactate, pyruvate, and ketone bodies in these tissues. In addition, there are specific and selective transporters for organic cations and organic anions

in organs including the kidney, intestine and liver. Organic anion transporters are selective for hydrophobic, charged molecules with electron-attracting side groups. Organic cation transporters, such as the ammonium transporter, mediate the secretion of a variety of drugs and endogenous metabolites, and contribute to the maintenance of intercellular pH (Poole, R.C. and A.P. Halestrap (1993) *Am. J. Physiol.* 264:C761-C782; Price, N.T. et al. (1998) *Biochem. J.* 329:321-328; and Martinelle, K. and I. Haggstrom (1993) *J. Biotechnol.* 30:339-350).

ATP-binding cassette (ABC) transporters are members of a superfamily of membrane proteins that transport substances ranging from small molecules such as ions, sugars, amino acids, peptides, and phospholipids, to lipopeptides, large proteins, and complex hydrophobic drugs. ABC transporters consist of four modules: two nucleotide-binding domains (NBD), which hydrolyze ATP to supply the energy required for transport, and two membrane-spanning domains (MSD), each containing six putative transmembrane segments. These four modules may be encoded by a single gene, as is the case for the cystic fibrosis transmembrane regulator (CFTR), or by separate genes. When encoded by separate genes, each gene product contains a single NBD and MSD. These "half-molecules" form homo- and heterodimers, such as Tap1 and Tap2, the endoplasmic reticulum-based major histocompatibility (MHC) peptide transport system. Several genetic diseases are attributed to defects in ABC transporters, such as the following diseases and their corresponding proteins: cystic fibrosis (CFTR, an ion channel), adrenoleukodystrophy (adrenoleukodystrophy protein, ALDP), Zellweger syndrome (peroxisomal membrane protein-70, PMP70), and hyperinsulinemic hypoglycemia (sulfonylurea receptor, SUR). Overexpression of the multidrug resistance (MDR) protein, another ABC transporter, in human cancer cells makes the cells resistant to a variety of cytotoxic drugs used in chemotherapy (Taglicht, D. and S. Michaelis (1998) *Meth. Enzymol.* 292:130-162).

A number of metal ions such as iron, zinc, copper, cobalt, manganese, molybdenum, selenium, nickel, and chromium are important as cofactors for a number of enzymes. For example, copper is involved in hemoglobin synthesis, connective tissue metabolism, and bone development, by acting as a cofactor in oxidoreductases such as superoxide dismutase, ferroxidase (ceruloplasmin), and lysyl oxidase. Copper and other metal ions must be provided in the diet, and are absorbed by transporters in the gastrointestinal tract. Plasma proteins transport the metal ions to the liver and other target organs, where specific transporters move the ions into cells and cellular organelles as needed. Imbalances in metal ion metabolism have been associated with a number of disease states (Danks, D.M. (1986) *J. Med. Genet.* 23:99-106).

Transport of fatty acids across the plasma membrane can occur by diffusion, a high capacity, low affinity process. However, under normal physiological conditions a significant fraction of fatty acid transport appears to occur via a high affinity, low capacity protein-mediated transport process.

Fatty acid transport protein (FATP), an integral membrane protein with four transmembrane segments, is expressed in tissues exhibiting high levels of plasma membrane fatty acid flux, such as muscle, heart, and adipose. Expression of FATP is upregulated in 3T3-L1 cells during adipose conversion, and expression in COS7 fibroblasts elevates uptake of long-chain fatty acids (Hui, T.Y. et al. (1998) J.

5 Biol. Chem. 273:27420-27429).

Mitochondrial carrier proteins are transmembrane-spanning proteins which transport ions and charged metabolites between the cytosol and the mitochondrial matrix. Examples include the ADP, ATP carrier protein; the 2-oxoglutarate/malate carrier; the phosphate carrier protein; the pyruvate carrier; the dicarboxylate carrier which transports malate, succinate, fumarate, and phosphate; the

10 tricarboxylate carrier which transports citrate and malate; and the Grave's disease carrier protein, a protein recognized by IgG in patients with active Grave's disease, an autoimmune disorder resulting in hyperthyroidism. Proteins in this family consist of three tandem repeats of an approximately 100 amino acid domain, each of which contains two transmembrane regions (Stryer, L. (1995) Biochemistry, W.H. Freeman and Company, New York NY, p. 551; PROSITE PDOC00189 Mitochondrial energy transfer

15 proteins signature; Online Mendelian Inheritance in Man (OMIM) *275000 Graves Disease).

This class of transporters also includes the mitochondrial uncoupling proteins, which create proton leaks across the inner mitochondrial membrane, thus uncoupling oxidative phosphorylation from ATP synthesis. The result is energy dissipation in the form of heat. Mitochondrial uncoupling proteins have been implicated as modulators of thermoregulation and metabolic rate, and have been proposed as

20 potential targets for drugs against metabolic diseases such as obesity (Ricquier, D. et al. (1999) J. Int. Med. 245:637-642).

Ion Channels

The electrical potential of a cell is generated and maintained by controlling the movement of ions across the plasma membrane. The movement of ions requires ion channels, which form ion-

25 selective pores within the membrane. There are two basic types of ion channels, ion transporters and gated ion channels. Ion transporters utilize the energy obtained from ATP hydrolysis to actively transport an ion against the ion's concentration gradient. Gated ion channels allow passive flow of an ion down the ion's electrochemical gradient under restricted conditions. Together, these types of ion channels generate, maintain, and utilize an electrochemical gradient that is used in 1) electrical impulse

30 conduction down the axon of a nerve cell, 2) transport of molecules into cells against concentration gradients, 3) initiation of muscle contraction, and 4) endocrine cell secretion.

Ion Transporters

Ion transporters generate and maintain the resting electrical potential of a cell. Utilizing the energy derived from ATP hydrolysis, they transport ions against the ion's concentration gradient.

These transmembrane ATPases are divided into three families. The phosphorylated (P) class ion transporters, including Na⁺-K⁺ ATPase, Ca²⁺-ATPase, and H⁺-ATPase, are activated by a phosphorylation event. P-class ion transporters are responsible for maintaining resting potential distributions such that cytosolic concentrations of Na⁺ and Ca²⁺ are low and cytosolic concentration of K⁺ is high. The vacuolar (V) class of ion transporters includes H⁺ pumps on intracellular organelles, such as lysosomes and Golgi. V-class ion transporters are responsible for generating the low pH within the lumen of these organelles that is required for function. The coupling factor (F) class consists of H⁺ pumps in the mitochondria. F-class ion transporters utilize a proton gradient to generate ATP from ADP and inorganic phosphate (P_i).

The P-ATPases are hexamers of a 100 kD subunit with ten transmembrane domains and several large cytoplasmic regions that may play a role in ion binding (Scarborough, G.A. (1999) *Curr. Opin. Cell Biol.* 11:517-522). The V-ATPases are composed of two functional domains: the V₁ domain, a peripheral complex responsible for ATP hydrolysis; and the V₀ domain, an integral complex responsible for proton translocation across the membrane. The F-ATPases are structurally and evolutionarily related to the V-ATPases. The F-ATPase F₀ domain contains 12 copies of the c subunit, a highly hydrophobic protein composed of two transmembrane domains and containing a single buried carboxyl group in TM2 that is essential for proton transport. The V-ATPase V₀ domain contains three types of homologous c subunits with four or five transmembrane domains and the essential carboxyl group in TM4 or TM3. Both types of complex also contain a single a subunit that may be involved in regulating the pH dependence of activity (Forgacs, M. (1999) *J. Biol. Chem.* 274:12951-12954).

The resting potential of the cell is utilized in many processes involving carrier proteins and gated ion channels. Carrier proteins utilize the resting potential to transport molecules into and out of the cell. Amino acid and glucose transport into many cells is linked to sodium ion co-transport (symport) so that the movement of Na⁺ down an electrochemical gradient drives transport of the other molecule up a concentration gradient. Similarly, cardiac muscle links transfer of Ca²⁺ out of the cell with transport of Na⁺ into the cell (antiport).

Gated Ion Channels

Gated ion channels control ion flow by regulating the opening and closing of pores. The ability to control ion flux through various gating mechanisms allows ion channels to mediate such diverse signaling and homeostatic functions as neuronal and endocrine signaling, muscle contraction, fertilization, and regulation of ion and pH balance. Gated ion channels are categorized according to the manner of regulating the gating function. Mechanically-gated channels open their pores in response to mechanical stress; voltage-gated channels (e.g., Na⁺, K⁺, Ca²⁺, and Cl⁻ channels) open their pores in response to changes in membrane potential; and ligand-gated channels (e.g.,

acetylcholine-, serotonin-, and glutamate-gated cation channels, and GABA- and glycine-gated chloride channels) open their pores in the presence of a specific ion, nucleotide, or neurotransmitter. The gating properties of a particular ion channel (i.e., its threshold for and duration of opening and closing) are sometimes modulated by association with auxiliary channel proteins and/or post

translational modifications, such as phosphorylation.

Mechanically-gated or mechanosensitive ion channels act as transducers for the senses of touch, hearing, and balance, and also play important roles in cell volume regulation, smooth muscle contraction, and cardiac rhythm generation. A stretch-inactivated channel (SIC) was recently cloned from rat kidney. The SIC channel belongs to a group of channels which are activated by pressure or stress on the cell membrane and conduct both Ca^{2+} and Na^{+} (Suzuki, M. et al. (1999) J. Biol. Chem. 274:6330-6335).

The pore-forming subunits of the voltage-gated cation channels form a superfamily of ion channel proteins. The characteristic domain of these channel proteins comprises six transmembrane domains (S1-S6), a pore-forming region (P) located between S5 and S6, and intracellular amino and carboxy termini. In the Na^{+} and Ca^{2+} subfamilies, this domain is repeated four times, while in the K^{+} channel subfamily, each channel is formed from a tetramer of either identical or dissimilar subunits. The P region contains information specifying the ion selectivity for the channel. In the case of K^{+} channels, a GYG tripeptide is involved in this selectivity (Ishii, T.M. et al. (1997) Proc. Natl. Acad. Sci. USA 94:11651-11656).

Voltage-gated Na^{+} and K^{+} channels are necessary for the function of electrically excitable cells, such as nerve and muscle cells. Action potentials, which lead to neurotransmitter release and muscle contraction, arise from large, transient changes in the permeability of the membrane to Na^{+} and K^{+} ions. Depolarization of the membrane beyond the threshold level opens voltage-gated Na^{+} channels. Sodium ions flow into the cell, further depolarizing the membrane and opening more voltage-gated Na^{+} channels, which propagates the depolarization down the length of the cell. Depolarization also opens voltage-gated potassium channels. Consequently, potassium ions flow outward, which leads to repolarization of the membrane. Voltage-gated channels utilize charged residues in the fourth transmembrane segment (S4) to sense voltage change. The open state lasts only about 1 millisecond, at which time the channel spontaneously converts into an inactive state that cannot be opened irrespective of the membrane potential. Inactivation is mediated by the channel's N-terminus, which acts as a plug that closes the pore. The transition from an inactive to a closed state requires a return to resting potential.

Voltage-gated Na^{+} channels are heterotrimeric complexes composed of a 260 kDa pore-forming α subunit that associates with two smaller auxiliary subunits, $\beta 1$ and $\beta 2$. The $\beta 2$ subunit is a integral

membrane glycoprotein that contains an extracellular Ig domain, and its association with α and $\beta 1$ subunits correlates with increased functional expression of the channel, a change in its gating properties, as well as an increase in whole cell capacitance due to an increase in membrane surface area (Isom, L.L. et al. (1995) Cell 83:433-442).

5 Non voltage-gated Na^+ channels include the members of the amiloride-sensitive Na^+ channel/degenerin (NaC/DEG) family. Channel subunits of this family are thought to consist of two transmembrane domains flanking a long extracellular loop, with the amino and carboxyl termini located within the cell. The NaC/DEG family includes the epithelial Na^+ channel (ENaC) involved in Na^+ reabsorption in epithelia including the airway, distal colon, cortical collecting duct of the kidney, and
10 exocrine duct glands. Mutations in ENaC result in pseudohypoaldosteronism type 1 and Liddle's syndrome (pseudohyperaldosteronism). The NaC/DEG family also includes the recently characterized H^+ -gated cation channels or acid-sensing ion channels (ASIC). ASIC subunits are expressed in the brain and form heteromultimeric Na^+ -permeable channels. These channels require acid pH fluctuations for activation. ASIC subunits show homology to the degenerins, a family of mechanically-gated
15 channels originally isolated from *C. elegans*. Mutations in the degenerins cause neurodegeneration. ASIC subunits may also have a role in neuronal function, or in pain perception, since tissue acidosis causes pain (Waldmann, R. and M. Lazdunski (1998) Curr. Opin. Neurobiol. 8:418-424; Eglen, R.M. et al. (1999) Trends Pharmacol. Sci. 20:337-342).

K^+ channels are located in all cell types, and may be regulated by voltage, ATP concentration,
20 or second messengers such as Ca^{2+} and cAMP. In non-excitabile tissue, K^+ channels are involved in protein synthesis, control of endocrine secretions, and the maintenance of osmotic equilibrium across membranes. In neurons and other excitable cells, in addition to regulating action potentials and repolarizing membranes, K^+ channels are responsible for setting resting membrane potential. The cytosol contains non-diffusible anions and, to balance this net negative charge, the cell contains a Na^+ -
25 K^+ pump and ion channels that provide the redistribution of Na^+ , K^+ , and Cl^- . The pump actively transports Na^+ out of the cell and K^+ into the cell in a 3:2 ratio. Ion channels in the plasma membrane allow K^+ and Cl^- to flow by passive diffusion. Because of the high negative charge within the cytosol, Cl^- flows out of the cell. The flow of K^+ is balanced by an electromotive force pulling K^+ into the cell, and a K^+ concentration gradient pushing K^+ out of the cell. Thus, the resting membrane potential is
30 primarily regulated by K^+ flow (Salkoff, L. and T. Jegla (1995) Neuron 15:489-492).

Potassium channel subunits of the Shaker-like superfamily all have the characteristic six transmembrane/1 pore domain structure. Four subunits combine as homo- or heterotetramers to form functional K channels. These pore-forming subunits also associate with various cytoplasmic β subunits that alter channel inactivation kinetics. The Shaker-like channel family includes the voltage-

gated K⁺ channels as well as the delayed rectifier type channels such as the human ether-a-go-go related gene (HERG) associated with long QT, a cardiac dysrhythmia syndrome (Curran, M.E. (1998) *Curr. Opin. Biotechnol.* 9:565-572; Kaczorowski, G.J. and M.L. Garcia (1999) *Curr. Opin. Chem. Biol.* 3:448-458).

5 A second superfamily of K⁺ channels is composed of the inward rectifying channels (Kir). Kir channels have the property of preferentially conducting K⁺ currents in the inward direction. These proteins consist of a single potassium selective pore domain and two transmembrane domains, which correspond to the fifth and sixth transmembrane domains of voltage-gated K⁺ channels. Kir subunits also associate as tetramers. The Kir family includes ROMK1, mutations in which lead to Bartter
10 syndrome, a renal tubular disorder. Kir channels are also involved in regulation of cardiac pacemaker activity, seizures and epilepsy, and insulin regulation (Doupnik, C.A. et al. (1995) *Curr. Opin. Neurobiol.* 5:268-277; Curran, *supra*).

 The recently recognized TWIK K⁺ channel family includes the mammalian TWIK-1, TREK-1 and TASK proteins. Members of this family possess an overall structure with four transmembrane
15 domains and two P domains. These proteins are probably involved in controlling the resting potential in a large set of cell types (Duprat, F. et al. (1997) *EMBO J* 16:5464-5471).

 The voltage-gated Ca²⁺ channels have been classified into several subtypes based upon their electrophysiological and pharmacological characteristics. L-type Ca²⁺ channels are predominantly expressed in heart and skeletal muscle where they play an essential role in excitation-contraction
20 coupling. T-type channels are important for cardiac pacemaker activity, while N-type and P/Q-type channels are involved in the control of neurotransmitter release in the central and peripheral nervous system. The L-type and N-type voltage-gated Ca²⁺ channels have been purified and, though their functions differ dramatically, they have similar subunit compositions. The channels are composed of three subunits. The α_1 subunit forms the membrane pore and voltage sensor, while the $\alpha_2\delta$ and β
25 subunits modulate the voltage-dependence, gating properties, and the current amplitude of the channel. These subunits are encoded by at least six α_1 , one $\alpha_2\delta$, and four β genes. A fourth subunit, γ , has been identified in skeletal muscle (Walker, D. et al. (1998) *J. Biol. Chem.* 273:2361-2367; McCleskey, E.W. (1994) *Curr. Opin. Neurobiol.* 4:304-312).

 The transient receptor family (Trp) of calcium ion channels are thought to mediate capacitative
30 calcium entry (CCE). CCE is the Ca²⁺ influx into cells to resupply Ca²⁺ stores depleted by the action of inositol triphosphate (IP3) and other agents in response to numerous hormones and growth factors. Trp and Trp-like were first cloned from *Drosophila* and have similarity to voltage gated Ca²⁺ channels in the S3 through S6 regions. This suggests that Trp and/or related proteins may form mammalian CCC entry channels (Zhu, X. et al. (1996) *Cell* 85:661-671; Boulay, G. et al. (1997) *J. Biol. Chem.*

272:29672-29680). Melastatin is a gene isolated in both the mouse and human, and whose expression in melanoma cells is inversely correlated with melanoma aggressiveness in vivo. The human cDNA transcript corresponds to a 1533-amino acid protein having homology to members of the Trp family. It has been proposed that the combined use of malastatin mRNA expression status and tumor thickness might allow for the determination of subgroups of patients at both low and high risk for developing metastatic disease (Duncan, L.M. et al (2001) J. Clin. Oncol. 19:568-576).

Chloride channels are necessary in endocrine secretion and in regulation of cytosolic and organelle pH. In secretory epithelial cells, Cl^- enters the cell across a basolateral membrane through an Na^+ , K^+/Cl^- cotransporter, accumulating in the cell above its electrochemical equilibrium concentration. Secretion of Cl^- from the apical surface, in response to hormonal stimulation, leads to flow of Na^+ and water into the secretory lumen. The cystic fibrosis transmembrane conductance regulator (CFTR) is a chloride channel encoded by the gene for cystic fibrosis, a common fatal genetic disorder in humans. CFTR is a member of the ABC transporter family, and is composed of two domains each consisting of six transmembrane domains followed by a nucleotide-binding site. Loss of CFTR function decreases transepithelial water secretion and, as a result, the layers of mucus that coat the respiratory tree, pancreatic ducts, and intestine are dehydrated and difficult to clear. The resulting blockage of these sites leads to pancreatic insufficiency, "meconium ileus", and devastating "chronic obstructive pulmonary disease" (Al-Awqati, Q. et al. (1992) J. Exp. Biol. 172:245-266).

The voltage-gated chloride channels (CLC) are characterized by 10-12 transmembrane domains, as well as two small globular domains known as CBS domains. The CLC subunits probably function as homotetramers. CLC proteins are involved in regulation of cell volume, membrane potential stabilization, signal transduction, and transepithelial transport. Mutations in CLC-1, expressed predominantly in skeletal muscle, are responsible for autosomal recessive generalized myotonia and autosomal dominant myotonia congenita, while mutations in the kidney channel CLC-5 lead to kidney stones (Jentsch, T.J. (1996) Curr. Opin. Neurobiol. 6:303-310).

Ligand-gated channels open their pores when an extracellular or intracellular mediator binds to the channel. Neurotransmitter-gated channels are channels that open when a neurotransmitter binds to their extracellular domain. These channels exist in the postsynaptic membrane of nerve or muscle cells. There are two types of neurotransmitter-gated channels. Sodium channels open in response to excitatory neurotransmitters, such as acetylcholine, glutamate, and serotonin. This opening causes an influx of Na^+ and produces the initial localized depolarization that activates the voltage-gated channels and starts the action potential. Chloride channels open in response to inhibitory neurotransmitters, such as γ -aminobutyric acid (GABA) and glycine, leading to hyperpolarization of the membrane and the subsequent generation of an action potential. Neurotransmitter-gated ion channels have four

transmembrane domains and probably function as pentamers (Jentsch, supra). Amino acids in the second transmembrane domain appear to be important in determining channel permeation and selectivity (Sather, W.A. et al. (1994) *Curr. Opin. Neurobiol.* 4:313-323).

Ligand-gated channels can be regulated by intracellular second messengers. For example, calcium-activated K^+ channels are gated by internal calcium ions. In nerve cells, an influx of calcium during depolarization opens K^+ channels to modulate the magnitude of the action potential (Ishi et al., supra). The large conductance (BK) channel has been purified from brain and its subunit composition determined. The α subunit of the BK channel has seven rather than six transmembrane domains in contrast to voltage-gated K^+ channels. The extra transmembrane domain is located at the subunit N-terminus. A 28-amino-acid stretch in the C-terminal region of the subunit (the "calcium bowl" region) contains many negatively charged residues and is thought to be the region responsible for calcium binding. The β subunit consists of two transmembrane domains connected by a glycosylated extracellular loop, with intracellular N- and C-termini (Kaczorowski, supra; Vergara, C. et al. (1998) *Curr. Opin. Neurobiol.* 8:321-329).

Cyclic nucleotide-gated (CNG) channels are gated by cytosolic cyclic nucleotides. The best examples of these are the cAMP-gated Na^+ channels involved in olfaction and the cGMP-gated cation channels involved in vision. Both systems involve ligand-mediated activation of a G-protein coupled receptor which then alters the level of cyclic nucleotide within the cell. CNG channels also represent a major pathway for Ca^{2+} entry into neurons, and play roles in neuronal development and plasticity.

CNG channels are tetramers containing at least two types of subunits, an α subunit which can form functional homomeric channels, and a β subunit, which modulates the channel properties. All CNG subunits have six transmembrane domains and a pore forming region between the fifth and sixth transmembrane domains, similar to voltage-gated K^+ channels. A large C-terminal domain contains a cyclic nucleotide binding domain, while the N-terminal domain confers variation among channel subtypes (Zufall, F. et al. (1997) *Curr. Opin. Neurobiol.* 7:404-412).

The activity of other types of ion channel proteins may also be modulated by a variety of intracellular signalling proteins. Many channels have sites for phosphorylation by one or more protein kinases including protein kinase A, protein kinase C, tyrosine kinase, and casein kinase II, all of which regulate ion channel activity in cells. Kir channels are activated by the binding of the $G\beta\gamma$ subunits of heterotrimeric G-proteins (Reimann, F. and F.M. Ashcroft (1999) *Curr. Opin. Cell. Biol.* 11:503-508). Other proteins are involved in the localization of ion channels to specific sites in the cell membrane. Such proteins include the PDZ domain proteins known as MAGUKs (membrane-associated guanylate kinases) which regulate the clustering of ion channels at neuronal synapses (Craven, S.E. and D.S. Bredt (1998) *Cell* 93:495-498).

Disease Correlation

The etiology of numerous human diseases and disorders can be attributed to defects in the transport of molecules across membranes. Defects in the trafficking of membrane-bound transporters and ion channels are associated with several disorders, e.g., cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, von Gierke disease, and certain forms of diabetes mellitus. Single-gene defect diseases resulting in an inability to transport small molecules across membranes include, e.g., cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease (van't Hoff, W.G. (1996) *Exp. Nephrol.* 4:253-262; Talente, G.M. et al. (1994) *Ann. Intern. Med.* 120:218-226; and Chillon, M. et al. (1995) *New Engl. J. Med.* 332:1475-1480).

Human diseases caused by mutations in ion channel genes include disorders of skeletal muscle, cardiac muscle, and the central nervous system. Mutations in the pore-forming subunits of sodium and chloride channels cause myotonia, a muscle disorder in which relaxation after voluntary contraction is delayed. Sodium channel myotonias have been treated with channel blockers. Mutations in muscle sodium and calcium channels cause forms of periodic paralysis, while mutations in the sarcoplasmic calcium release channel, T-tubule calcium channel, and muscle sodium channel cause malignant hyperthermia. Cardiac arrhythmia disorders such as the long QT syndromes and idiopathic ventricular fibrillation are caused by mutations in potassium and sodium channels (Cooper, E.C. and L.Y. Jan (1998) *Proc. Natl. Acad. Sci. USA* 96:4759-4766). All four known human idiopathic epilepsy genes code for ion channel proteins (Berkovic, S.F. and I.E. Scheffer (1999) *Curr. Opin. Neurology* 12:177-182). Other neurological disorders such as ataxias, hemiplegic migraine and hereditary deafness can also result from mutations in ion channel genes (Jen, J. (1999) *Curr. Opin. Neurobiol.* 9:274-280; Cooper, *supra*).

Ion channels have been the target for many drug therapies. Neurotransmitter-gated channels have been targeted in therapies for treatment of insomnia, anxiety, depression, and schizophrenia. Voltage-gated channels have been targeted in therapies for arrhythmia, ischemic stroke, head trauma, and neurodegenerative disease (Taylor, C.P. and L.S. Narasimhan (1997) *Adv. Pharmacol.* 39:47-98). Various classes of ion channels also play an important role in the perception of pain, and thus are potential targets for new analgesics. These include the vanilloid-gated ion channels, which are activated by the vanilloid capsaicin, as well as by noxious heat. Local anesthetics such as lidocaine and mexiletine which blockade voltage-gated Na⁺ channels have been useful in the treatment of neuropathic pain (Eglen, *supra*).

Ion channels in the immune system have recently been suggested as targets for immunomodulation. T-cell activation depends upon calcium signaling, and a diverse set of T-cell specific ion channels has been characterized that affect this signaling process. Channel blocking agents

can inhibit secretion of lymphokines, cell proliferation, and killing of target cells. A peptide antagonist of the T-cell potassium channel Kv1.3 was found to suppress delayed-type hypersensitivity and allogenic responses in pigs, validating the idea of channel blockers as safe and efficacious immunosuppressants (Cahalan, M.D. and K.G. Chandy (1997) Curr. Opin. Biotechnol. 8:749-756).

5 The discovery of new transporters and ion channels, and the polynucleotides encoding them, satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of transport, neurological, muscle, immunological, and cell proliferative disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of transporters and ion channels.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, transporters and ion channels, referred to collectively as "TRICH" and individually as "TRICH-1," "TRICH-2," "TRICH-3," "TRICH-4," "TRICH-5," "TRICH-6," "TRICH-7," "TRICH-8," "TRICH-9," "TRICH-10," "TRICH-11,"
15 "TRICH-12," "TRICH-13," "TRICH-14," "TRICH-15," "TRICH-16," "TRICH-17," "TRICH-18," "TRICH-19," "TRICH-20," "TRICH-21," "TRICH-22," "TRICH-23," "TRICH-24," "TRICH-25," "TRICH-26," "TRICH-27," "TRICH-28," "TRICH-29," "TRICH-30," "TRICH-31," and "TRICH-32." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID
20 NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. In one alternative, the invention provides an
25 isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-32.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-
30 32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-32. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:33-64.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) combining the polypeptide with at least one test compound under

suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:33-64, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the

polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a

reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“TRICH” refers to the amino acid sequences of substantially purified TRICH obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term “agonist” refers to a molecule which intensifies or mimics the biological activity of TRICH. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TRICH either by directly interacting with TRICH or by acting on components of the biological pathway in which TRICH participates.

An “allelic variant” is an alternative form of the gene encoding TRICH. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding TRICH include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as TRICH or a polypeptide with at least one functional characteristic of TRICH. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding TRICH, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding TRICH. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent TRICH. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge,

solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of TRICH is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of TRICH. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TRICH either by directly interacting with TRICH or by acting on components of the biological pathway in which TRICH participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind TRICH polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic TRICH, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding TRICH or fragments of TRICH may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

“Conservative amino acid substitutions” are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
10	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
15	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
20	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
25	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A “deletion” refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term “derivative” refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A “detectable label” refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

“Differential expression” refers to increased or upregulated; or decreased, downregulated, or

absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

5 A "fragment" is a unique portion of TRICH or the polynucleotide encoding TRICH which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid
10 residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present
15 embodiments.

A fragment of SEQ ID NO:33-64 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:33-64, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:33-64 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish
20 SEQ ID NO:33-64 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:33-64 and the region of SEQ ID NO:33-64 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-32 is encoded by a fragment of SEQ ID NO:33-64. A fragment of SEQ ID NO:1-32 comprises a region of unique amino acid sequence that specifically identifies
25 SEQ ID NO:1-32. For example, a fragment of SEQ ID NO:1-32 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-32. The precise length of a fragment of SEQ ID NO:1-32 and the region of SEQ ID NO:1-32 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

30 A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

35 The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to

the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore, achieve a more meaningful comparison of the two sequences.

- 5 Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191.
- 10 For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

- Alternatively, a suite of commonly used and freely available sequence comparison algorithms is
- 15 provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other
- 20 polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to
- 25 compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

- 30 *Open Gap: 5 and Extension Gap: 2 penalties*

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases “percent identity” and “% identity,” as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and “diagonals saved”=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the “percent similarity” between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the “BLAST 2 Sequences” tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for

example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term “humanized antibody” refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour.

Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of TRICH which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of TRICH which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of TRICH. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of TRICH.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or

synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably
5 linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of
10 amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an TRICH may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the
15 art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of TRICH.

"Probe" refers to nucleic acid sequences encoding TRICH, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical
20 labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous
25 nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the
30 specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR

Protocols. A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

5 Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU
10 primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer
15 binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that
20 hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary
25 polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the
30 artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing TRICH, nucleic acids encoding TRICH, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers,

microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

5 "Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection,
10 electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to
15 animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in
20 vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention
25 into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999)
30 set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant

identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule.

Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human transporters and ion channels (TRICH), the polynucleotides encoding TRICH, and the use of these compositions for the diagnosis, treatment, or prevention of transport, neurological, muscle, immunological, and cell proliferative disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank

homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are transporters and ion channels. For example, SEQ ID NO:5 is 83% identical to rat GABA receptor rho-3 subunit precursor (GenBank ID g1060975) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $1.7e-206$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:5 also contains a neurotransmitter-gated ion channel domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:5 is a neurotransmitter-gated ion channel. In an alternate example, SEQ ID NO:16 is 57% identical to human Na⁺/glucose cotransporter (GenBank ID g338055) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $2.4e-181$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:16 also contains a sodium:solute symporter family domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:16 is a Na⁺/glucose cotransporter. In an alternate example, SEQ ID NO:27 is 53% identical to human ATP-binding cassette transporter-1 (ABC-1) (GenBank ID g4128033) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 0.0, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:27 also contains an ABC transporter domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from

BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:27 is an ABC transporter. In an alternate example, SEQ ID NO:12 is 45% identical to rat thyroid sodium/iodide symporter NIS (GenBank ID g1399954) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $3.0e-143$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:12 also contains a sodium:solute symporter family domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:12 is a sodium:solute symporter. SEQ ID NO:1-4, SEQ ID NO:6-11, SEQ ID NO:13-15, SEQ ID NO:17-26, and SEQ ID NO:28-32 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-32 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:33-64 or that distinguish between SEQ ID NO:33-64 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 6724643H1 is the identification number of an Incyte cDNA sequence, and LUNLTMT01 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71495515V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g5746200) which contributed to the assembly of the full length polynucleotide sequences. In addition, the identification numbers in column 5 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (*i.e.*, those sequences

including the designation "ENST"). Alternatively, the identification numbers in column 5 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (*i.e.*, those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (*i.e.*, those sequences including the designation "NP"). Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For example, FL_XXXXXX_N₁_N₂_YYYYY_N₃_N₄ represents a "stitched" sequence in which XXXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYY is the number of the prediction generated by the algorithm, and N_{1,2,3...}, if present, represent specific exons that may have been manually edited during analysis (See Example V).

Alternatively, the identification numbers in column 5 may refer to assemblages of exons brought together by an "exon-stretching" algorithm. For example, FLXXXXXX_gAAAAA_gBBBBB_1_N is the identification number of a "stretched" sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq sequence was used as a protein homolog for the "exon-stretching" algorithm, a RefSeq identifier (denoted by "NM," "NP," or "NT") may be used in place of the GenBank identifier (*i.e.*, gBBBBB).

Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis methods associated with the prefixes (see Example IV and Example V).

Prefix	Type of analysis and/or examples of programs
GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGENES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
GBI	Hand-edited analysis of genomic sequences.
FL	Stitched or stretched genomic sequences (see Example V).

In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte

cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses TRICH variants. A preferred TRICH variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the TRICH amino acid sequence, and which contains at least one functional or structural characteristic of TRICH.

The invention also encompasses polynucleotides which encode TRICH. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:33-64, which encodes TRICH. The polynucleotide sequences of SEQ ID NO:33-64, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding TRICH. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding TRICH. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:33-64 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:33-64. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of TRICH.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding TRICH, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring TRICH, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode TRICH and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring TRICH under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding TRICH or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide

occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding TRICH and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode TRICH and TRICH derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding TRICH or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:33-64 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding TRICH may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic

DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.)

Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids

5 Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences
10 are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060).

Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences,
15 Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include
20 sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary
25 sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer
30 controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode TRICH may be cloned in recombinant DNA molecules that direct expression of TRICH, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of

the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express TRICH.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter TRICH-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of TRICH, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding TRICH may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, TRICH itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of TRICH, or any part thereof, may be altered during direct synthesis and/or combined with sequences

from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.)

5 The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active TRICH, the nucleotide sequences encoding TRICH or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a
10 suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding TRICH. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding TRICH. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where
15 sequences encoding TRICH and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural
20 and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding TRICH and appropriate transcriptional and translational control
25 elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences
30 encoding TRICH. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or

animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO J. 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding TRICH. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding TRICH can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding TRICH into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of TRICH are needed, e.g. for the production of antibodies, vectors which direct high level expression of TRICH may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of TRICH. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of TRICH. Transcription of sequences encoding TRICH may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311).

Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding TRICH may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses TRICH in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of TRICH in cell lines is preferred. For example, sequences encoding TRICH can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*⁻ and *apr*⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980)

Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding TRICH is inserted within a marker gene sequence, transformed cells containing sequences encoding TRICH can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding TRICH under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding TRICH and that express TRICH may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of TRICH using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on TRICH is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding TRICH include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding TRICH, or any fragments thereof, may be cloned into a vector for

the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega
5 (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding TRICH may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein
10 produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode TRICH may be designed to contain signal sequences which direct secretion of TRICH through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the
15 inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities
20 (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding TRICH may be ligated to a heterologous sequence resulting in translation of a
25 fusion protein in any of the aforementioned host systems. For example, a chimeric TRICH protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of TRICH activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose
30 binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize

these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the TRICH encoding sequence and the heterologous protein sequence, so that TRICH may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled TRICH may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

TRICH of the present invention or fragments thereof may be used to screen for compounds that specifically bind to TRICH. At least one and up to a plurality of test compounds may be screened for specific binding to TRICH. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of TRICH, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which TRICH binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express TRICH, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing TRICH or cell membrane fractions which contain TRICH are then contacted with a test compound and binding, stimulation, or inhibition of activity of either TRICH or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with TRICH, either in solution or affixed to a solid support, and detecting the binding of TRICH to the compound.

Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

TRICH of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of TRICH. Such compounds may include agonists, antagonists, or partial

or inverse agonists. In one embodiment, an assay is performed under conditions permissive for TRICH activity, wherein TRICH is combined with at least one test compound, and the activity of TRICH in the presence of a test compound is compared with the activity of TRICH in the absence of the test compound. A change in the activity of TRICH in the presence of the test compound is indicative of a compound that modulates the activity of TRICH. Alternatively, a test compound is combined with an in vitro or cell-free system comprising TRICH under conditions suitable for TRICH activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of TRICH may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding TRICH or their mammalian homologs may be “knocked out” in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding TRICH may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding TRICH can also be used to create “knockin” humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding TRICH is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with

potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress TRICH, e.g., by secreting TRICH in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) *Biotechnol. Annu. Rev.* 4:55-74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of TRICH and transporters and ion channels. In addition, the expression of TRICH is closely associated with adrenal, testicular, and prostate tumors, Crohn's disease, teratocarcinoma and dendritic cells, brain, lung, ileum, small intestine, uterine myometrial, colon, and pancreatic tissues. Therefore, TRICH appears to play a role in transport, neurological, muscle, immunological, and cell proliferative disorders. In the treatment of disorders associated with increased TRICH expression or activity, it is desirable to decrease the expression or activity of TRICH. In the treatment of disorders associated with decreased TRICH expression or activity, it is desirable to increase the expression or activity of TRICH.

Therefore, in one embodiment, TRICH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, catatonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis, neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia, and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other

extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a muscle disorder such as cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, pheochromocytoma, and myopathies including encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, ophthalmoplegia, and acid maltase deficiency (AMD, also known as Pompe's disease); an immunological disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal

circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including
5 adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing TRICH or a fragment or derivative
10 thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified TRICH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those
15 provided above.

In still another embodiment, an agonist which modulates the activity of TRICH may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those listed above.

In a further embodiment, an antagonist of TRICH may be administered to a subject to treat or
20 prevent a disorder associated with increased expression or activity of TRICH. Examples of such disorders include, but are not limited to, those transport, neurological, muscle, immunological, and cell proliferative disorders described above. In one aspect, an antibody which specifically binds TRICH may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express TRICH.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding TRICH may be administered to a subject to treat or prevent a disorder associated with
25 increased expression or activity of TRICH including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate
30 therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of TRICH may be produced using methods which are generally known in the art. In particular, purified TRICH may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind TRICH. Antibodies to TRICH may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with TRICH or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to TRICH have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of TRICH amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to TRICH may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce TRICH-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be

generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for TRICH may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between TRICH and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering TRICH epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for TRICH. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of TRICH-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple TRICH epitopes, represents the average affinity, or avidity, of the antibodies for TRICH. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular TRICH epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the TRICH-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of TRICH, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of TRICH-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding TRICH, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding TRICH. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding TRICH. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding TRICH may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial

hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) Science 270:404-410; Verma, I.M. and N. Somia (1997) Nature 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) Nature 335:395-396; Poeschla, E. et al. (1996) Proc. Natl. Acad. Sci. USA. 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in TRICH expression or regulation causes disease, the expression of TRICH from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in TRICH are treated by constructing mammalian expression vectors encoding TRICH and introducing these vectors by mechanical means into TRICH-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) Annu. Rev. Biochem. 62:191-217; Ivics, Z. (1997) Cell 91:501-510; Boulay, J-L. and H. Récipon (1998) Curr. Opin. Biotechnol. 9:445-450).

Expression vectors that may be effective for the expression of TRICH include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). TRICH may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) Proc. Natl. Acad. Sci. USA 89:5547-5551; Gossen, M. et al. (1995) Science 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) Curr. Opin. Biotechnol. 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding TRICH from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver

polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to TRICH expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding TRICH under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding TRICH to cells which have one or more genetic abnormalities with respect to the expression of TRICH. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu.*

Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding TRICH to target cells which have one or more genetic abnormalities with respect to the expression of TRICH. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing TRICH to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding TRICH to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for TRICH into the alphavirus genome in place of the capsid-coding region results in the production of a large number of TRICH-coding RNAs and the synthesis of high levels of TRICH in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application

(Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of TRICH into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding TRICH.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding TRICH. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding TRICH.

Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased TRICH expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding TRICH may be therapeutically useful, and in the treatment of disorders associated with decreased TRICH expression or activity, a compound which specifically promotes expression of the polynucleotide encoding TRICH may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding TRICH is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding TRICH are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding TRICH. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of

the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of TRICH, antibodies to TRICH, and mimetics, agonists, antagonists, or inhibitors of TRICH.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form.

These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S.

Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising TRICH or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, TRICH or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example TRICH or fragments thereof, antibodies of TRICH, and agonists, antagonists or inhibitors of TRICH, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy.

Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind TRICH may be used for the diagnosis of disorders characterized by expression of TRICH, or in assays to monitor patients being treated with TRICH or agonists, antagonists, or inhibitors of TRICH. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for TRICH include methods which utilize the antibody and a label to detect TRICH in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring TRICH, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of TRICH expression. Normal or standard values for TRICH expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to TRICH under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of TRICH expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding TRICH may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of TRICH may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of TRICH, and to monitor regulation of TRICH levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding TRICH or closely related molecules may be used to identify nucleic acid sequences which encode TRICH. The specificity of the probe, whether it is made

from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding TRICH, allelic variants, or related sequences.

5 Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the TRICH encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:33-64 or from genomic sequences including promoters, enhancers, and introns of the TRICH gene.

Means for producing specific hybridization probes for DNAs encoding TRICH include the
 10 cloning of polynucleotide sequences encoding TRICH or TRICH derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as
 15 alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding TRICH may be used for the diagnosis of disorders associated with expression of TRICH. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes
 20 insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, catatonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis,
 25 cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis, neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia, and other disorders associated with transport, e.g., neurofibromatosis,
 30 postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and

Fanconi disease; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a muscle disorder such as cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, pheochromocytoma, and myopathies including encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, ophthalmoplegia, and acid maltase deficiency (AMD, also known as Pompe's disease); an immunological disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic

anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding TRICH may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered TRICH expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding TRICH may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding TRICH may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding TRICH in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of TRICH, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding TRICH, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several
5 days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ
10 preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding *TRICH* may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding
15 *TRICH*, or a fragment of a polynucleotide complementary to the polynucleotide encoding *TRICH*, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences
20 encoding *TRICH* may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding *TRICH* are used to amplify DNA using the
25 polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as
30 DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the

alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of TRICH include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, TRICH, fragments of TRICH, or antibodies specific for TRICH may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention
5 may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000)
10 Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested
15 compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for
20 example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample
25 containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples
30 are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are

analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for TRICH to quantify the levels of TRICH expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoz, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference

in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding TRICH may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding TRICH on a physical

map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps.

- 5 Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences
10 mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

- In another embodiment of the invention, TRICH, its catalytic or immunogenic fragments, or
15 oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between TRICH and the agent being tested may be measured.

- Another technique for drug screening provides for high throughput screening of compounds
20 having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with TRICH, or fragments thereof, and washed. Bound TRICH is then detected by methods well known in the art. Purified TRICH can also be coated directly onto plates for use in the aforementioned drug screening techniques.
25 Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

- In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding TRICH specifically compete with a test compound for binding TRICH. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more
30 antigenic determinants with TRICH.

In additional embodiments, the nucleotide sequences which encode TRICH may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

5 The disclosures of all patents, applications and publications, mentioned above and below including U.S. Ser. No. 60/216,547, U.S. Ser. No. 60/218,232, U.S. Ser. No. 60/220,112, and U.S. Ser. No. 60/221,839 are expressly incorporated by reference herein., are expressly incorporated by reference herein.

10 EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a
15 suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA
20 purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

25 In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic
30 oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,

PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX

5 DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC
10 Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a
15 high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

20 III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the
25 MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI
30 PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:33-64. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

5 IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative transporters and ion channels were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin
10 (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode transporters and ion channels, the encoded polypeptides were analyzed by querying
15 against PFAM models for transporters and ion channels. Potential transporters and ion channels were also identified by homology to Incyte cDNA sequences that had been annotated as transporters and ion channels. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by
20 Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using
25 the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification
30 program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence.

Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then “stitched” together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbprl public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

“Stretched” Sequences

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore “stretched” or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

VI. Chromosomal Mapping of TRICH Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:33-64 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:33-64 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences

had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For

example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

5 Alternatively, polynucleotide sequences encoding TRICH are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system;
10 embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer,
15 cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding TRICH. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

20 VIII. Extension of TRICH Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using
25 OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension
30 was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme

(Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 µl PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

IX. Labeling and Use of Individual Hybridization Probes

5 Hybridization probes derived from SEQ ID NO:33-64 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston
10 MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or
15 Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate.
20 Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra.), mechanical
25 microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), supra.). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be
30 produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37° C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85° C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and
5 coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 µl of the array element DNA, at an average concentration of 100 ng/µl, is loaded into the open capillary printing element by a high-speed robotic
10 apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in
15 0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with
20 an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide
30 containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477,
35 Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate

filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

5 The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different
10 fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC
15 computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

20 A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

XI. Complementary Polynucleotides

25 Sequences complementary to the TRICH-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring TRICH. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of TRICH. To inhibit transcription, a
30 complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the TRICH-encoding transcript.

XII. Expression of TRICH

Expression and purification of TRICH is achieved using bacterial or virus-based expression systems. For expression of TRICH in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express TRICH upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of TRICH in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding TRICH by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, TRICH is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from TRICH at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified TRICH obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII where applicable.

XIII. Functional Assays

TRICH function is assessed by expressing the sequences encoding TRICH at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which

contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of TRICH on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding TRICH and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding TRICH and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIV. Production of TRICH Specific Antibodies

TRICH substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the TRICH amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-TRICH activity by, for example, binding the peptide or TRICH to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XV. Purification of Naturally Occurring TRICH Using Specific Antibodies

Naturally occurring or recombinant TRICH is substantially purified by immunoaffinity chromatography using antibodies specific for TRICH. An immunoaffinity column is constructed by covalently coupling anti-TRICH antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing TRICH are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of TRICH (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/TRICH binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and TRICH is collected.

XVI. Identification of Molecules Which Interact with TRICH

Molecules which interact with TRICH may include transporter substrates, agonists or antagonists, modulatory proteins such as G β y proteins (Reimann, *supra*) or proteins involved in TRICH localization or clustering such as MAGUKs (Craven, *supra*). TRICH, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled TRICH, washed, and any wells with labeled TRICH complex are assayed. Data obtained using different concentrations of TRICH are used to calculate values for the number, affinity, and association of TRICH with the candidate molecules.

Alternatively, proteins that interact with TRICH are isolated using the yeast 2-hybrid system (Fields, S. and O. Song (1989) Nature 340:245-246). TRICH, or fragments thereof, are expressed as fusion proteins with the DNA binding domain of Gal4 or lexA, and potential interacting proteins are expressed as fusion proteins with an activation domain. Interactions between the TRICH fusion protein and the TRICH interacting proteins (fusion proteins with an activation domain) reconstitute a transactivation function that is observed by expression of a reporter gene. Yeast 2-hybrid systems are

commercially available, and methods for use of the yeast 2-hybrid system with ion channel proteins are discussed in Niethammer, M. and M. Sheng (1998, Meth. Enzymol. 293:104-122).

TRICH may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Potential TRICH agonists or antagonists may be tested for activation or inhibition of TRICH ion channel activity using the assays described in section XVIII.

XVII. Demonstration of TRICH Activity

Ion channel activity of TRICH is demonstrated using an electrophysiological assay for ion conductance. TRICH can be expressed by transforming a mammalian cell line such as COS7, HeLa or CHO with a eukaryotic expression vector encoding TRICH. Eukaryotic expression vectors are commercially available, and the techniques to introduce them into cells are well known to those skilled in the art. A second plasmid which expresses any one of a number of marker genes, such as β -galactosidase, is co-transformed into the cells to allow rapid identification of those cells which have taken up and expressed the foreign DNA. The cells are incubated for 48-72 hours after transformation under conditions appropriate for the cell line to allow expression and accumulation of TRICH and β -galactosidase.

Transformed cells expressing β -galactosidase are stained blue when a suitable colorimetric substrate is added to the culture media under conditions that are well known in the art. Stained cells are tested for differences in membrane conductance by electrophysiological techniques that are well known in the art. Untransformed cells, and/or cells transformed with either vector sequences alone or β -galactosidase sequences alone, are used as controls and tested in parallel. Cells expressing TRICH will have higher anion or cation conductance relative to control cells. The contribution of TRICH to conductance can be confirmed by incubating the cells using antibodies specific for TRICH. The antibodies will bind to the extracellular side of TRICH, thereby blocking the pore in the ion channel, and the associated conductance.

Alternatively, ion channel activity of TRICH is measured as current flow across a TRICH-containing *Xenopus laevis* oocyte membrane using the two-electrode voltage-clamp technique (Ishi et al., *supra*; Jegla, T. and L. Salkoff (1997) J. Neurosci. 17:32-44). TRICH is subcloned into an appropriate *Xenopus* oocyte expression vector, such as pBF, and 0.5-5 ng of mRNA is injected into mature stage IV oocytes. Injected oocytes are incubated at 18°C for 1-5 days. Inside-out macropatches are excised into an intracellular solution containing 116 mM K-gluconate, 4 mM KCl, and 10 mM Hepes (pH 7.2). The intracellular solution is supplemented with varying concentrations of the TRICH mediator, such as cAMP, cGMP, or Ca^{+2} (in the form of CaCl_2), where appropriate.

Electrode resistance is set at 2-5 M Ω and electrodes are filled with the intracellular solution lacking mediator. Experiments are performed at room temperature from a holding potential of 0 mV. Voltage ramps (2.5 s) from -100 to 100 mV are acquired at a sampling frequency of 500 Hz. Current measured is proportional to the activity of TRICH in the assay.

5 In particular, the activities of TRICH-1, TRICH-2, and TRICH-10, are measured as K⁺ conductance, the activities of TRICH-6 and TRICH-9 are measured as K⁺ conductance in the presence of membrane stretch or free fatty acids, the activities of TRICH-18, TRICH-25 and TRICH-31 are measured as voltage-gated K⁺ conductance, TRICH-5 activity is measured as Cl⁻ conductance in the presence of GABA, TRICH-11 activity is measured as cation conductance in the presence of heat, and
10 the activity of TRICH-9, TRICH-28 is measured as Ca²⁺ conductance.

Transport activity of TRICH is assayed by measuring uptake of labeled substrates into Xenopus laevis oocytes. Oocytes at stages V and VI are injected with TRICH mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM CaCl₂, 1mM MgCl₂, 1mM Na₂HPO₄, 5 mM Hepes, 3.8 mM NaOH, 50 μ g/ml gentamycin, pH 7.8) to allow
15 expression of TRICH. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM CaCl₂, 1mM MgCl₂, 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids, sugars, drugs, ions, and neurotransmitters) is initiated by adding labeled substrate (e.g. radiolabeled with ³H, fluorescently labeled with rhodamine, etc.) to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in Na⁺-free medium, measuring the
20 incorporated label, and comparing with controls. TRICH activity is proportional to the level of internalized labeled substrate. In particular, test substrates include pigment precursors and related molecules for TRICH-3, aminophospholipids for TRICH-4, fructose and glucose for TRICH-7 and TRICH-15, amino acids for TRICH-8, Na⁺ and iodide for TRICH-12, Na⁺ and H⁺ for TRICH-13 and TRICH-21, Na⁺ and glucose for TRICH-16 and TRICH-19, and glucose for TRICH-23, TRICH-26,
25 TRICH-29, TRICH-30, and TRICH-32.

ATPase activity associated with TRICH can be measured by hydrolysis of radiolabeled ATP-[γ -³²P], separation of the hydrolysis products by chromatographic methods, and quantitation of the recovered ³²P using a scintillation counter. The reaction mixture contains ATP-[γ -³²P] and varying amounts of TRICH in a suitable buffer incubated at 37°C for a suitable period of time. The reaction is
30 terminated by acid precipitation with trichloroacetic acid and then neutralized with base, and an aliquot of the reaction mixture is subjected to membrane or filter paper-based chromatography to separate the reaction products. The amount of ³²P liberated is counted in a scintillation counter. The amount of radioactivity recovered is proportional to the ATPase activity of TRICH in the assay.

XVIII. Identification of TRICH Agonists and Antagonists

TRICH is expressed in a eukaryotic cell line such as CHO (Chinese Hamster Ovary) or HEK (Human Embryonic Kidney) 293. Ion channel activity of the transformed cells is measured in the presence and absence of candidate agonists or antagonists. Ion channel activity is assayed using patch clamp methods well known in the art or as described in Example XVII. Alternatively, ion channel activity is assayed using fluorescent techniques that measure ion flux across the cell membrane (Velicelebi, G. et al. (1999) Meth. Enzymol. 294:20-47; West, M.R. and C.R. Molloy (1996) Anal. Biochem. 241:51-58). These assays may be adapted for high-throughput screening using microplates. Changes in internal ion concentration are measured using fluorescent dyes such as the Ca^{2+} indicator Fluo-4 AM, sodium-sensitive dyes such as SBFI and sodium green, or the Cl^- indicator MQAE (all available from Molecular Probes) in combination with the FLIPR fluorimetric plate reading system (Molecular Devices). In a more generic version of this assay, changes in membrane potential caused by ionic flux across the plasma membrane are measured using oxonyl dyes such as DiBAC₄ (Molecular Probes). DiBAC₄ equilibrates between the extracellular solution and cellular sites according to the cellular membrane potential. The dye's fluorescence intensity is 20-fold greater when bound to hydrophobic intracellular sites, allowing detection of DiBAC₄ entry into the cell (Gonzalez, J.E. and P.A. Negulescu (1998) Curr. Opin. Biotechnol. 9:624-631). Candidate agonists or antagonists may be selected from known ion channel agonists or antagonists, peptide libraries, or combinatorial chemical libraries.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
3474673	1	3474673CD1	33	3474673CB1
4588877	2	4588877CD1	34	4588877CB1
7472214	3	7472214CD1	35	7472214CB1
7473053	4	7473053CD1	36	7473053CB1
7473347	5	7473347CD1	37	7473347CB1
7474240	6	7474240CD1	38	7474240CB1
7475338	7	7475338CD1	39	7475338CB1
7476747	8	7476747CD1	40	7476747CB1
7477898	9	7477898CD1	41	7477898CB1
7472728	10	7472728CD1	42	7472728CB1
7474322	11	7474322CD1	43	7474322CB1
5455621	12	5455621CD1	44	5455621CB1
7477248	13	7477248CD1	45	7477248CB1
2944004	14	2944004CD1	46	2944004CB1
3046849	15	3046849CD1	47	3046849CB1
4538363	16	4538363CD1	48	4538363CB1
6427460	17	6427460CD1	49	6427460CB1
7474127	18	7474127CD1	50	7474127CB1
7476949	19	7476949CD1	51	7476949CB1
7477249	20	7477249CD1	52	7477249CB1
7477720	21	7477720CD1	53	7477720CB1
7477852	22	7477852CD1	54	7477852CB1
1471717	23	1471717CD1	55	1471717CB1
3874406	24	3874406CD1	56	3874406CB1
4599654	25	4599654CD1	57	4599654CB1
5047435	26	5047435CD1	58	5047435CB1
7475603	27	7475603CD1	59	7475603CB1
7477845	28	7477845CD1	60	7477845CB1
168827	29	168827CD1	61	168827CB1
7472734	30	7472734CD1	62	7472734CB1
7473473	31	7473473CD1	63	7473473CB1
7477725	32	7477725CD1	64	7477725CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	3474673CD1	g13507377	1.00E-151	[fl][Homo sapiens] potassium channel TASK-4 (Decher, N. et al. (2001) FEBS Lett. 492 (1-2), 84-89)
2	4588877CD1	g13926111	3.00E-96	[fl][Homo sapiens] (AF358910) 2P domain potassium channel Talk-2
3	7472214CD1	g1107730	1.70E-243	[Mus musculus] ABC8 (Savary, S. et al. (1996) Mamm. Genome 7 (9), 673-676)
4	7473053CD1	g11342541	0	[fl][Homo sapiens] putative white family ATP-binding cassette transporter
		g3850108	9.00E-209	[Schizosaccharomyces pombe] putative calcium- transporting atpase
		g3628757	0	[Homo sapiens] FIC1
5	7473347CD1	g1060975	1.70E-206	(Bull, L.N. et al. (1998) Nat. Genet. 18 (3), 219-224) [Rattus norvegicus] GABA receptor rho-3 subunit precursor
6	7474240CD1	g2745727	0	(Ogurusu, T. et al. (1996) Biochim. Biophys. Acta 1305 (1-2), 15-18) [Rattus norvegicus] potassium channel
7	7475338CD1	g183298	2.10E-158	(Shi, W. et al. (1997) J. Neurosci. 17 (24), 9423-9432) [Homo sapiens] GLUT5 protein (Kayano, T. et al. (1990) J. Biol. Chem. 265 (22), 13276-13282)
9	7477898CD1	g2745729	0	[Rattus norvegicus] potassium channel (Shi, W. et al. (1997) J. Neurosci. 17 (24), 9423-9432)
10	7472728CD1	g8452900	3.50E-261	[Rattus norvegicus] potassium channel TREK-2 (Bang, H. et al. (2000) J. Biol. Chem. 275 (23), 17412- 17419)
11	7474322CD1	g12003146	0	[fl][Homo sapiens] capsaicin receptor
12	5455621CD1	g1399954	3.00E-143	[Rattus norvegicus] thyroid sodium/iodide symporter NIS (Dai, G. et al. (1996) Nature 379 (6564), 458-460)
13	7477248CD1	g2944233	3.10E-195	[Homo sapiens] sodium-hydrogen exchanger 6 (Numata, M. et al. (1998) J. Biol. Chem. 273 (12), 6951- 6959)
14	2944004CD1	g3451312	1.40E-188	[Schizosaccharomyces pombe] membrane atpase
15	3046849CD1	g12802047	0	[fl][Homo sapiens] (AJ271290) facilitative glucose transporter GLUT11

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
16	4538363CD1	g338055	7.40E-181	[Homo sapiens] Na ⁺ /glucose cotransporter (Hediger, M.A. et al. (1989) Proc. Natl. Acad. Sci. U.S.A. 86 (15), 5748-5752)
17	6427460CD1	g6457274	0	[Mus musculus] putative EL-E2 ATPase (Halleck, M.S. et al. (1999) Physiol. Genomics (Online) 1 (3), 139-150)
18	7474127CD1	g206044	0	[Rattus norvegicus] potassium channel Kv3.2b (Wiedmann, R. et al. (1991) FEBS Lett. 288, 163-167)
19	7476949CD1	g9588428	0	[5' incm] [Homo sapiens] dj1024N4.1 (novel Sodium: solute symporter family member similar to SLC5A1 (SGLT1))
20	7477249CD1	g7715417	0	[Homo sapiens] Na ⁺ /glucose cotransporter (Hediger, M.A. et al. (1989) Proc. Natl. Acad. Sci. U.S.A. 86 (15), 5748-5752)
21	747720CD1	g205709	0	[Oryctolagus cuniculus] RING-finger binding protein (Mansharmani, M. et al. (2001) J. Biol. Chem. 276 (5), 3641-3649)
22	7477852CD1	g8920219	0	[Rattus norvegicus] sodium-hydrogen exchange protein-isoform 4 (Orlowski, J. et al. (1992) J. Biol. Chem. 267, 9331-9339)
23	1471717CD1	g529590	5.00E-36	[fl] [Homo sapiens] epithelial calcium channel (Muller, D. et al. (2000) Genomics 67 (1), 48-53)
24	3874406CD1	g1514530	1.90E-117	[Rattus norvegicus] liver-specific transport protein (Simonsen, G.D. et al. (1994) J. Cell. Sci. 107, 1065-1072)
25	4599654CD1	g3242244	0	[Homo sapiens] ABC-C transporter (Klugbauer, N. et al. (1996) FEBS Lett. 391 (1-2), 61-65)
				[Mus musculus] hyperpolarization-activated cation channel, HAC3 (Ludwig, A. et al. (1998) Nature 393 (6685), 587-591)

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
26	5047435CD1	g13445575	0	[fl][Homo sapiens] facilitative glucose transporter GLUT10 (McVie-Wyllie, A.J. et al. (2001) Genomics 72 (1), 113-117)
27	7475603CD1	g9211112	0	[fl][Homo sapiens] macrophage ABC transporter (Kaminski, W.E. et al. (2000) Biochem. Biophys. Res. Commun. 273 (2), 532-538)
28	7477845CD1	g3800830	0	[Rattus norvegicus] putative four repeat ion channel (Lee, J.H. et al. (1999) FEBS Lett. 445 (2-3), 231-236)
29	168827CD1	g7707622	1.20E-116	[Homo sapiens] organic anion transporter 4 (Cha, S.H. et al. (2000) J. Biol. Chem. 275 (6), 4507-4512)
30	7472734CD1	g3004482	0	[fl][Rattus norvegicus] putative integral membrane transport protein (Schomig, E. et al. (1998) FEBS Lett. 425 (1), 79-86)
		g7707622	4.50E-117	[Homo sapiens] organic anion transporter 4 (Cha, S.H. et al. (2000) J. Biol. Chem. 275 (6), 4507-4512)
31	7473473CD1	g3004482	0	[fl][Rattus norvegicus] putative integral membrane transport protein (Schomig, E. et al. (1998) FEBS Lett. 425 (1), 79-86)
		g6625694	0	[Rattus norvegicus] potassium channel Eag2 (Saganich, M.J. et al. (1999) J. Neurosci. 19 (24), 10789-10802)
32	7477725CD1	g3004482	1.00E-177	[fl][Rattus norvegicus] putative integral membrane transport protein (Schomig, E. et al. (1998) FEBS Lett. 425 (1), 79-86)
		g7707622	4.20E-130	[Homo sapiens] organic anion transporter 4 (Cha, S.H. et al. (2000) J. Biol. Chem. 275 (6), 4507-4512)

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	3474673CD1	332	S201 S207 S234 S265 S280 S281 S289 S51 T169 T67	N65 N94	Transmembrane domains: R130-M155, V245-L264 TASK K+ channel domain: V14-S332	HMMER
2	4588877CD1	226	S101 S128 S159 S174 S175 S183 S95		Transmembrane domain: V139-L158 CHANNEL PROTEIN IONIC POTASSIUM SUBUNIT K+ PUTATIVE SUBFAMILY K MEMBER PD021430: A78-E162	HMMER
3	7472214CD1	646	S143 S229 S261 S340 S341 S463 S554 S57 S644 S69 S89 T138 T157 T23 T472 T500 T591	N169 N422	Transmembrane domains: S430-M450, W564-D589, M618-V637 ABC transporter domain: R95-G277 ABC transporters family signature BL00211: I100-F111, L201-D232 ABC transporters family signature: V181-D232 PROTEIN TRANSMEMBRANE TRANSPORT ATPBINDING TRANSPORTER MEMBRANE ABC GLYCOPROTEIN INNER PUTATIVE PD000633: T365-Y583 do WHITE; FRUIT; FLY; SCARLET; DM05200 P45844 289-650: G277-L623 ABC TRANSPORTERS FAMILY DM00008 P45844 73-287: I61-Q276 ABC transporter motif: L201-L215 ATP/GTP binding site (P-loop): G102-S109	HMMER HMMER_PPFAM BLIMPS_BLOCKS PROFILESCAN BLAST_PRODUM BLAST_DOMO MOTIFS MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4	7473053CD1	1190	S153 S259 S268 S391 S413 S452 S493 S545 S573 S624 S631 S687 S723 S739 S744 S832 S1174 S1132 S1164 S1124 S1143 S1168 T267 T36 T370 T378 T514 T519 T580 T646 T705 T732 T899 T980 T1098 T1158 Y23 Y29 Y489 Y607	N579	Transmembrane domains: S77-V94, L276-W298, Y330-R350, L947-I971, Q991-I1009 E1-E2 ATPase domains: E381-V403, Q530-A562, Y633-G685, R788-D818 E1-E2 ATPases phosphorylation site proteins BL00154: G134-L151, V386-F404, D650-M690, T809-S832 E1-E2 ATPases phosphorylation site: A372-V417 P-type cation-transporting ATPase superfamily signature PR00119: F390-F404, A666-D676, I812-I831 ATPASE HYDROLASE TRANSMEMBRANE PHOSPHORYLATION ATPBINDING PROTEIN PROBABLE CALCIUMTRANSPORTING CALCIUM TRANSPORT PD004657: S846-P1093 FIC1 PROTEIN PD180313: H1039-W1165 do ATPASE; CALCIUM; TRANSPORTING; DM02405 P32660 318-1225: W128-F418, E466-N910 ATPase E1-E2 motif: D392-T398 Transmembrane domain: V332-V351	HMMER HMMER_PPFAM BLIMPS_BLOCKS PROFILES SCAN BLIMPS_PRINTS BLAST_PROD OM BLAST_DOMO MOTIFS HMMER
5	7473347CD1	467	S149 S175 S344 S37 S390 S411 S419 S427 S53 S96 T100 T136 T157 T355 T356 T366 T41	N126 N197 N220		

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
5					Neurotransmitter-gated ion-channel domain: P58-Q362, H441-W463 Neurotransmitter-gated ion channels signature BL00236: V85-P122, I139-H148, D169-Y207, Y254-A295 Neurotransmitter-gated ion-channels signature: L164-H218 Neurotransmitter-gated ion-channels signature PR00252: T105-F121, K138-S149, C184-C198, S261-P273 Gamma-aminobutyric acid A (GABA) receptor signature PR00253: F270-W290, V296-V317, V330-V351, Y446-Y466 CHANNEL IONIC TRANSMEMBRANE GLYCOPROTEIN POSTSYNAPTIC MEMBRANE RECEPTOR PRECURSOR SIGNAL PROTEIN PD000153: E62-S427 NEUROTRANSMITTER-GATED ION-CHANNELS DM00560 P50573 34-464: S37-V467 Neurotransmitter-gated ion channels motif: C184-C198	HMMEP_PFAM BLIMPS_BLOCKS PROFILESSCAN BLIMPS_PRINTS BLIMPS_PRINTS BLAST_PRODOR BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	7474240CD1	1196	S174 S187 S209 S211 S239 S269 S274 S275 S317 S349 S354 S514 S55 S609 S639 S821 S869 S879 S883 S896 S899 S906 S922 S923 S939 S940 S963 S974 S985 S1020 S1091 S1170 S1096 T133 T169 T344 T371 T392 T528 T582 T637 T673 T74 T829 T857 T916 T1022 T1027 T1134 T1099 Y248 Y446 Y98	N102 N230 N338 N369 N600 N661 N736 N881 N905 N1139	Transmembrane domain: V551-Y571 Transmembrane region cyclic nucleotide gated ion channel: Y492-I731 Cyclic nucleotide-binding domain: M759-E850 POTASSIUM CHANNEL IONIC CHANNEL PD104127: S852-Y1028 POTASSIUM CHANNEL IONIC CHANNEL PD104126: A1076-K1196 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 I38465 562-948: H564-A914 do POTASSIUM; CHANNEL; KST1; AKT1; DM02383 I38465 353-560: S353-A563 do CHANNEL; POTASSIUM; EAG; DM05484 I38465 1-351: M1-P351	HMMER HMMER_PPFAM HMMER_PPFAM BLAST_PRODOM BLAST_PRODOM BLAST_DOMO BLAST_DOMO BLAST_DOMO
7	7475338CD1	512	S222 S279 S412 S413 S438 T107 T170 T235 T247 T473 T59 T66 Y380	N41 N57	Signal peptide: M1-A35 Transmembrane domains: C79-G96, M171-L188, Y322-V342, F448-I466 Sugar (and other) transporter domain: A26-F481 Sugar transport proteins signatures: A119-I185, V323-S379 Sugar transporter signature PR00171: A35-V45, V135-M154, Q294-Y304, I383-V404, T406-F418 Glucose transporter signature PR00172: L284-Y305, Q321-V342, L352-Q372, I383-T406, A416-F434, Y446-I466	SPSCAN HMMER HMMER_PPFAM PROFILES SCAN BLIMPS_PRINTS BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7					SUGAR TRANSPORT PROTEINS DM00135 P22732 I32-466: R138-F473 Sugar transporter 1 motif: S338-A353 Sugar transporter 2 motif: V140-R165	BLAST_DOMO MOTIFS MOTIFS
8	7476747CD1	568	S143 S365 S4 S456 S46 S51 S55 T34 T430 Y45	N141 N205 N214 N256 N562 N62 N76	Transmembrane domains: I242-F269, Y289-P308, I322-Y342 Transmembrane amino acid transporter protein domain: A102-G543 ACID AMINO PROTEIN TRANSPORTER PERMEASE TRANSMEMBRANE INTERGENIC REGION PUTATIVE PROLINE PD001875: W80-I380	HMMER HMMER_PFAM BLAST_PRODUM
9	7477898CD1	958	S105 S140 S145 S200 S26 S283 S288 S458 S488 S55 S670 S706 S724 S751 S774 S788 S864 S872 S879 S897 S929 T13 T170 T202 T220 T301 T326 T363 T377 T486 T522 T678	N218 N449 N510 N742	Transmembrane domain: L300-N318 Transmembrane region cyclic nucleotide gated ion channel: Y341-I580 Cyclic nucleotide-binding domain: V608-A699 POTASSIUM CHANNEL IONIC CHANNEL PD118772: E702-S955 CHANNEL PROTEIN IONIC POTASSIUM NONPHOTOTROPIC HYPOCOTYL PUTATIVE SUBUNIT REPEAT EAG PD009483: M1-L86 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 I38465 562-948: H413-F738, do POTASSIUM; CHANNEL; KST1; AKT1; DM02383 I38465 353-560: T201-A412	HMMER HMMER_PFAM HMMER_PFAM BLAST_PRODUM BLAST_PRODUM BLAST_DOMO BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
10	7472728CD1	724	S229 S283 S303 S333 S512 S545 S597 S666 S718 T104 T19 T223 T444 T515 T540 T557 T591 T636 T640 T650 T661 T676	N327 N330 N331 N532 N664 N684 N716	Transmembrane domains: A370-L388, I419-F437, V486-M503 TASK K+ channel domain: M250-D646 TWIK1 RELATED POTASSIUM CHANNEL, SUBFAMILY K, MEMBER 2 TREK1 K+ CHANNEL SUBUNIT IONIC CHANNEL PD085853: P215-G326	HMMER HMMER_PFAM BLAST_PRODROM
11	7474322CD1	470	S134 S142 S245 S326 S355 S408 S411 S415 S432 S452 T15 T22 T229 T265 T337 T341 T36	N236 N256 N321 N380	Transmembrane domains: F62-Y87, F139-F163, F212-L230, I293-I312 VANILLOID RECEPTOR SUBTYPE 1 PD137334: C348-K470	HMMER BLAST_PRODROM
12	5455621CD1	618	S110 S265 S313 S373 S490 S550 S565 S576 S594 T154 T237 T268 T360 T37 T526 T567 T70	N219 N256 N480 N574	Transmembrane domains: D10-F28, F81-Y104, F278-M297, L439-Y459, I502-R528 Sodium:solute symporter family domain: F41-G445 Sodium:solute symporter signature BL00456: T154-G208 Sodium:solute symporter family signature: N151-T198 TRANSMEMBRANE TRANSPORT PERMEASE PROTEIN SODIUM SYMPORT PROLINE COTRANSPORTER SYMPORTER GLYCOPROTEIN PD000991: F41-C304 SYMPORTER SODIUM IODIDE THYROID SODIUM/IODIDE NIS PD024705: I446-L489, S490-G575 SODIUM:SOLUTE SYMPORTER FAMILY DM00745 P31636 24-561: D10-N219, G220-Y459	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLAST_PRODROM BLAST_PRODROM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
13	7477248CD1	631	S149 S212 S258 S522 S9 T518 T551 T73 T79 Y14	N352 N516 N96	Transmembrane domains: V22-F41, L159-M181, I391-A407 Sodium/hydrogen exchanger family domain: L25-V491 Na+/H+ exchanger isoform 6 signature PR01088: Y14-I38, W39-V57, Y58-V84, Q119-E132, A269-M288, T480-Q506, K515- D533, P539-Q567, P566-E593 Na+/H+ exchanger signature PR01084: I133-F144, G147-S161, I162- T170, G208-T218 + TRANSPORT EXCHANGER NA PD01672: I133-M181 Na+/H+ PROTEIN TRANSMEMBRANE TRANSPORT ANTIporter SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: G20-G63, E132-R490 SODIUMHYDROGEN EXCHANGER 6 MYELOBLAST KIAA0267 PD177855: G478-Y591 do BETA; EXCHANGER; NA; DM02572 P48764 10-734: L124-L541	HMMER HMMER_PFBAM BLIMPS_PRINTS BLIMPS_PRINTS BLIMPS_PRODUM BLAST_PRODUM BLAST_PRODUM BLAST_PRODUM BLAST_DOMO HMMER HMMER_PFBAM BLIMPS_BLOCKS PROFILES SCAN
14	2944004CD1	1256	S103 S130 S144 S170 S227 S252 S523 S802 S817 S899 S901 S98 S1055 T269 T353 T358 T387 T502 T549 T576 T74 T912 T1212 T1061 T1236 Y349 Y407	N150 N23 N300 N312 N318 N704 N1045 N1053 N1059 N1073 N1247	Transmembrane domains: Y231-Y251, L415-L434, I933-I959, F966- L985, I1002-F1020, N1104-M1122 E1-E2 ATPase domains: V274-V365, G490-D506, Q672-A785, L851- S899 E1-E2 ATPases phosphorylation site signature BL00154: V454-G490, L492-L510, K652- C662, N724-M764, V878-S901, A905-V938 E1-E2 ATPases phosphorylation site: I478-E526	HMMER HMMER_PFBAM BLIMPS_BLOCKS PROFILES SCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
14					P-type cation-transporting ATPase superfamily signature PR00119: N318-T332, C496-L510, A740-D750, C881-L900 ATPASE PROBABLE CALCIUMTRANSPORTING PROTEIN HYDROLASE CALCIUM TRANSPORT TRANSMEMBRANE PHOSPHORYLATION MAGNESIUM PD090368: Q995-Y1094, D1064-L1114 E1-E2 ATPASES PHOSPHORYLATION SITE DM00115 P22189 49-801: S202-K331, P401-E505, S556-A575, V623-P767, H800-S984 E1-E2 ATPase motif: D498-T504	BLIMPS_PRINTS BLAST_PRODUM BLAST_DOMO
15	3046849CD1	499	S100 S118 S215 S285 T466 T487	N292 N34 N50	Signal peptide: M1-G27 Transmembrane domains: M163-L181, T371-G389, M418-L440 Sugar (and other) transporter signature: L18-L474 Sugar transport proteins signature: A112-V178 Sugar transporter signature PR00171: T28-I38, M128-M147, M376-L397, T399-C411 Glucose transporter signature PR00172: Q314-I335, M376-T399, A409-L427 SUGAR TRANSPORT PROTEINS DM00135 P22732 132-466: R131-T466 Sugar transporter 2 motif: L133-R158	MOTIFS SPSCAN HMWER HMWER_PFAM PROFILES SCAN BLIMPS_PRINTS BLIMPS_PRINTS BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
16	4538363CD1	596	S17 S290 S39 S5 T119 T211	N239 N386 N4 N545 N96	Transmembrane domains: S73-W95, I185-I212, L356-A376, L410-V430, F473-F491, Y513-L533 Sodium:solute symporter family domain: Y50-G479 Sodium:solute symporter signature BL00456: Y27-G81, A103-R132, L165-G219, P452-G461 Sodium:solute symporter family signatures: H162-I209, V412-D502 TRANSMEMBRANE TRANSPORT PERMEASE PROTEIN SODIUM SYMPORT PROLINE COTRANSPORTER SYMPORTER GLYCOPROTEIN PD000991: Y50-G479 NA+/GLUCOSE COTRANSPORTERRELATED PROTEIN PD134393: L551-A596 NA+/GLUCOSE COTRANSPORTERRELATED PROTEIN PD166538: M1-G49 SODIUM:SOLUTE SYMPORTER FAMILY DM00745 P13866 24-561: S17-W548 Na solute symporter 2 motif: G461-V481	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLAST_PROD OM BLAST_PROD OM BLAST_PROD OM BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
17	6427460CD1	1192	S143 S169 S188 S283 S287 S335 S451 S507 S508 S52 S555 S561 S722 S933 T203 T255 T259 T269 T333 T380 T413 T418 T659 T708 T714 T715 T910 T1103 T1017 T1105 Y885 Y1026	N397 N745 N921 N989 N1001	Transmembrane domains: V299-Y316, F1004-L1022, I1030-W1049, A1075-L1092 E1-E2 ATPase domains: E403-E425 I550-C698 E1-E2 ATPases phosphorylation site signature BL00154: G149-F166, V408-F426, D663-L703 E1-E2 ATPases phosphorylation site: L395-C442 P-type cation-transporting ATPase superfamily signature PR00119: F412-F426, A679-D689 ATPASE HYDROLASE TRANSMEMBRANE PHOSPHORYLATION ATPBINDING PROTEIN PROBABLE CALCIUMTRANSPORTING CALCIUM TRANSPORT PD004657: A857-V1108 do ATPASE; CALCIUM; TRANSPORTING; DM02405 Q09891 206-1107: T105-Y436, F471-N921 E1-E2 ATPase motif: D414-T420	HMMER HMMER_PFBM BLIMPS_BLOCKS PROFILES CAN BLIMPS_PRINTS BLAST_PRODOM
18	7474127CD1	638	S205 S224 S336 S378 S414 S541 S553 S564 S86 T120 T146 T155 T17 T21 T25 T283 T374 T49 T520 T546 T579	N259 N266 N518 N536 N84	Transmembrane domains: I231-L248, F382-Y401, M451-V473 Ion transport protein domain: L240-I472 Potassium channel signature PR00169: E101-T120, P222-T250, Y284-K307, F310-V330, F352-S378, E381-E404, F421-M443, G450-F476	MOTIFS HMMER HMMER_PFBM BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
18					VOLTAGEGATED POTASSIUM CHANNEL PROTEIN KV3.2 KSHIIIA IONIC TRANSMEMBRANE ION TRANSPORT GLYCOPROTEIN MULTIGENE FAMILY ALTERNATIVE SPLICING PHOSPHORYLATION PD085814: K495-S538 do CHANNEL; POTASSIUM; CDRK; FORM; DM00436 P22462 189-350: R189-R351 do CHANNEL; POTASSIUM; CDRK; SHAW; DM00490 P22462 34-151: L34-C152	BLAST_PRODUM BLAST_DOMO BLAST_DOMO
19	7476949CD1	681	S307 S421 S56 S573 S582 S587 S638 S651 T422 T485 T650 Y510	N113 N251 N256 N403 N603	Transmembrane domains: I38-I57, S90-W112, I150-I167, L188-M207, L373-A393, V432-I448, Y530-L550 Sodium:solute symporter family domain: Y67-G496 Sodium:solute symporter signature BL00456: Y44-G98, A120-R149, L182-G236, P469-A478 Sodium:solute symporter family signatures: Q179-V226, D458-D519 TRANSMEMBRANE TRANSPORT PERMEASE PROTEIN SODIUM SYMPORT PROLINE COTRANSPORTER SYMPORTER GLYCOPROTEIN PD000991: Y67-G496 SODIUM:SOLUTE SYMPORTER FAMILY DM00745 P13866 24-561: H34-W565 Na solute symporter 1 motif: G183-A208	HMMER HMMER_PPFAM BLIMPS_BLOCKS PROFILES SCAN BLAST_PRODUM BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
20	7477249CD1	1096	S115 S163 S276 S280 S332 S333 S404 S454 S46 S461 S462 S508 S514 S671 S863 S891 S1084 T262 T340 T345 T347 T407 T570 T612 T687 T840 T948 T1034 T1036 Y322	N331 N383 N395 N411 N720 N932	Transmembrane domains: F289-L307, F935-L953, W967-V996, F1008-D1028 E1-E2 ATPase domains: T340-Q352, H502-V648 E1-E2 ATPases phosphorylation site signature BL00154: G143-L160, V335-F353, K529- C539, D616-H656 P-type cation-transporting ATPase superfamily signature PR00119: F339-F353, A632-D642 H+-transporting ATPase signatur PR00120: T547-A565 ATPASE HYDROLASE TRANSMEMBRANE PHOSPHORYLATION ATPBINDING PROTEIN PROBABLE CALCIUMTRANSPORTING CALCIUM TRANSPORT PD004657: A787-K1038 do ATPASE; CALCIUM; TRANSPORTING; DM02405 P39524 236-1049: T83-I306, F422-N851 E1-E2 ATPase motif: D341-T347 Signal peptide: M1-A26 Transmembrane domains: I155-Y178, I271-T292, Sodium/hydrogen exchanger family domain: V73-K482 Na+/H+ exchanger signature PR01084: I158-A166, G200-A210, I129- L140, G143-S157	HMMER HMMER_PFBM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS BLAST_PRODUM
21	7477720CD1	707	S204 S299 S360 S417 S488 S51 S58 S585 S591 S620 S638 S679 T334 T350 T483 T634 Y225 Y528	N297 N31 N342 N35		MOTIFS SPSCAN HMMER HMMER_PFBM BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
21					Na ⁺ /H ⁺ exchanger isoform 2 (NHE2) signature PR01086: F115-S128, K616-I627 + TRANSPORT EXCHANGER NA PD01672: A83-I113, I129-L177, Y178-L212, A213-F249, D262-I287, S288-Y321, L322-M355, S359-F405, Y406-F452, I489-K531, I532-G562, R593-R640 NA ⁺ /H ⁺ PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 14-716: L15-L687	BLIMPS_PRINTS BLIMPS_PRODUM BLAST_PRODUM
22	7477852CD1	729	S142 S144 S155 S285 S291 S299 S318 S654 S664 S669 S697 S719 T110 T138 T281 T379 T447 T532 T539	N208 N358 N717	Transmembrane domains: F493-F512, M554-M570 Ankyrin repeats: L78-E108, A116-T148, F162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419	BLAST_DOMO HMMER HMMER_PFAM BLAST_PRODUM MOTIFS
23	1471717CD1	492	S13 S18 S225 S314 S373 T323 T33 T351 T426	N229 N249	transmembrane domain: I48-V71, V86-F104, Y172-I199, I199-V217, F384-F402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152:R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344:I119-N249	HMMER HMMER_PFAM BLAST_DOMO BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
24	3874406CD1	1494	S30 S50 S134 S230 S368 S549 S638 S669 S686 S696 S792 S800 S831 S912 S1004 S1070 S1146 S1172 S1206 S1365 T111 T435 T449 T501 T520 T632 T649 T657 T729 T845 T1049 T1134 T1217 T1247 T1295 T1318 T1339 T1422 T1482 Y824	N109 N130 N313 N421 N453 N71 N788 N817 N84 N867 N91 N1182	transmembrane domain: L204-F221, T272-L290, L735-Y753, F896-S914, V941-I959, L975-R998, F1019-V1039 ABC transporter: G384-G566 G1190-G1366 ABC transporters family proteins BL00211: I389-L400, L492-D523 ABC transporters family signature: V472-D523 ABC TRANSPORTERS FAMILY DM00008 P41233 839-1045:I355-N565, K1177-M1363 DM00008 P34358 611-816:I355-N565, A1179-M1363 DM00008 P41233 1851-2058:K1173-S1365, I355-N565 DM00008 P23703 41-246:E1162-G1366, L377-G566 ATP/GTP-binding site motif A (P-loop): G391-S398, G1197-2004	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
25	4599654CD1	774	S355 S356 S40 S505 S552 S559 S597 S61 S67 S734 S736 T203 T418 T668 T764 Y490	N291 N416	transmembrane domain: Y95-F118, T203-L219, L327-L353 Transmembrane region cyclic Nucleotide G: Y168-I414 Cyclic nucleotide-binding domain: K443-M531 Cyclic nucleotide-binding domain proteins BL00888: G452-V475, G488-L497 cAMP-dependent protein kinase signature PR00103: F449-R463, S489-T498 HYPERPOLARIZATIONACTIVATED CATION CHANNEL, HAC3 PD180735: T538-M774 CHANNEL IONIC POTASSIUM K+ SUBUNIT HYPERPOLARIZATIONACTIVATED PROTEIN PUTATIVE EAG LONG PD001039: E74-R167 cAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 A55251 333-706:H263-P561 DM01165 P29973 311-684:H263-P561 DM01165 Q03041 286-658:H263-G548 DM01165 S52072 262-635:H263-Q595	HMMER HMMER_PFAM HMMER_PFAM BLIMPS_BLOCKS BLIMPS_PRINTS BLAST_PRODUM BLAST_PRODUM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
26	5047435CD1	614	S116 S210 S290 S538 S577 S606 T267 T432 T443 T591	N407 N599	transmembrane domain: V124-I142, A168-M190, A371-V390, W483-I511, S526-I543, F552-V570 Sugar (and other) transporter: L83-F585 Sugar transport proteins BL00216: L174-S223, G92-S103 Sugar transporter signature PR00171: G92-I102, V175-I194, L486-V507, S509-F521 Glucose transporter signature PR00172: V343-V364, L486-S509, R519-L537, W550-V570 Sugar_Transport_1: G138-G153 A360-A375 Sugar transport proteins signatures sugar_transport_1.pr: L344-S401 sugar_transport_2.pr: A160-A225 SUGAR TRANSPORT PROTEINS DM00135 S25015 122-478:A160-D417, L480-K574, DM00135 P09830 101-452:G161-V405, L481-K574 DM00135 Q01440 101-433:R178-G388, R178-G388, L486-G575 DM00135 P15729 242-463:A485-S577, R286-L414	HMER HMER_PFAM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS MOTIFS PROFILES SCAN BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
27	7475603CD1	2180	S181 S216 S233 S260 S409 S419 S842 S983 S1008 S1172 S1229 S1237 S1269 S1349 S1353 S1462 S1469 S1504 S1566 S1881 S1993 S2018 S2174 S2167 T120 T165 T338 T348 T510 T599 T614 T822 T931 T1079 T1086 T1094 T1171 T1181 T1209 T1219 T1417 T1439 T1822 T1870 T1917 T1988 T2057 T2125 Y656 Y1448	N112 N132 N346 N374 N1100 N1415 N1420 N1491 N1552 N1695 N1831	transmembrane domain: F630-L648, L664-L680, V1570-V1590, M1622-Q1641 ABC transporter: G1854-G2035 G868-G1048 ABC transporters family BL00211: F873-T884, L974-D1005 ABC transporters family signature: A1940-D1991, D955-D1005 Abc_Transporter: L974-F988 ATP/GTP-binding site motif A (P-loop): G875-T882, G1861-T1868 ATPBINDING TRANSPORTER CASSETTE ABC TRANSPORT PROTEIN GLYCOPROTEIN TRANSMEMBRANE RIM ABCR PD005939: L1563-N1740 ATPBINDING TRANSPORTER CASSETTE ABC GLYCOPROTEIN TRANSMEMBRANE TRANSPORT ABCR RIM PD010118: R238-R514, L95-R243 ATPBINDING TRANSPORTER CASSETTE ABC GLYCOPROTEIN TRANSMEMBRANE TRANSPORT ABCR RIM SIMILARITY PD008845: P1307-E1560 ATPBINDING TRANSPORTER CASSETTE ABC GLYCOPROTEIN TRANSMEMBRANE TRANSPORT RIM ABCR SIMILARITY PD006867: L540-S685, D515-Q541	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN MOTIFS MOTIFS BLAST_PRODROM BLAST_PRODROM BLAST_PRODROM BLAST_PRODROM

108

[illegible]

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
28					III REPEAT DM00079 A55138 1052-1268:V1020-L1227 DM00079 P35500 1424-1636:W1090-P1194, I1017-N1050 IV REPEAT DM00277 P27732 1363-1572:F1337-L1536 DM00277 P15381 1384-1595:F1337-L1536	BLAST_DOMO
29	168827CD1	547	S109 S167 S201 S282 S336 S404 S408 S526 T133 T323 T35 T432 T453 T58	N102 N107 N56	transmembrane domain: F16-T35, Y180-C200, S201-V222, M410- E429, T469-Y492, L496-L514 Sugar (and other) transporter: L13-Q528 ORGANIC TRANSPORTERLIKE TRANSPORT PROTEIN RENAL ANION TRANSPORTER CATIONIC KIDNEYSPECIFIC SOLUTE PD151320: N102-L144	HMMER HMMER_PPFAM BLAST_PRODROM
30	7472734CD1	547	S143 S167 S201 S282 S336 S404 S408 S46 S526 S60 S68 T133 T323 T432 T453 T58	N102 N39 N56 N62	transmembrane domain: I18-F32, M147-Y163, Y180-C200, S201- V222, M410-E429, T469-Y492, L496-L514 Sugar (and other) transporter: L18-Q528 SUGAR TRANSPORT PROTEINS DM00032 P46501 280-351:V121-K173 ORGANIC TRANSPORTERLIKE TRANSPORT PROTEIN RENAL ANION TRANSPORTER CATIONIC KIDNEYSPECIFIC SOLUTE PD151320: N102-K145	HMMER HMMER_PPFAM BLAST_DOMO BLAST_PRODROM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
31	7473473CD1	988	S142 S237 S24 S252 S322 S369 S502 S680 S773 S847 S883 S925 S943 S952 S974 S981 T127 T14 T215 T442 T478 T521 T634 T725 T73 T832 T869 T909 T929	N170 N235 N403 N466 N663 N830	transmembrane domain: L342-A360 Transmembrane cyclic Nucleotide G: Y288-I536 Cyclic nucleotide-binding domain: V564-A655 PAC motif PA: C92-T132 CHANNEL POTASSIUM IONIC EAG SUBUNIT HEAG LONG ELECTROCARDIOGRAPHIC QT SYNDROME PD017645: K809-D984 CHANNEL IONIC K+ SUBUNIT HYPERPO- LARIZATION ACTIVATED PUTATIVE EAG LONG PD001039: S179-I284 CHANNEL K+ IONIC EAG SUBUNIT TRANSMEMBRANE ION TRANSPORT VOLTAGEGATED PD011550: N658-E737 CHANNEL PROTEIN IONIC POTASSIUM NON PHOTOTROPIC HYPOCOTYL PUTATIVE SUBUNIT REPEAT EAG PD009483: M1-E89 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE- BINDING DOMAIN DM01165 I48912 391-786: H361-S756 DM01165 Q02280 384-776: H361-E737 DM01165 I38465 562-948: H361-R671, S974-E985 POTASSIUM; CHANNEL; KST1; AKT1; DM02383 I48912 164-389: V162-E314, E314-A360, W362-V455	HMME HMME_PPFAM HMME_PPFAM HMME_PPFAM BLAST_PRODUM BLAST_PRODUM BLAST_PRODUM BLAST_PRODUM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
32	7477725CD1	533	S107 S109 S143 S167 S282 S345 S408 S469 S60 T133 T289 T323 T336 T432 T526	N102 N216 N56 N62	transmembrane domain: F150-D168, L380-N401, I407-V426, L486- F504 Sugar (and other) transporter: A111-K528 ORGANIC TRANSPORTER LIKE TRANSPORT PROTEIN RENAL ANION TRANSPORTER CATIONIC KIDNEY SPECIFIC SOLUTE PD151320: N102-K145	HMMER HMMER_PFAM BLAST_PRODOR

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
33	3474673CB1	1775	1-391, 578-786, 1024-1301	GNFL.g7798848_00000 3_004.edit 6724643H1 (LUNLTMT01) 3474673H1 (LUNGN0T27) 71495515V1 71495515V1 FL135171_00001 71497982V1 GBI:g8117242_000054 _edit.8639-8803 GBI:g8117242_000054 _edit.4857-4997 GBI:g8117242_000054 _edit.10305-10463 6891360H1 (BRAITDR03) GBI:g8117242_000054 _edit.50-89 GBI:g8117242_000054 _edit.6950-7093 GBI:g8117242_000054 _edit.4345-4478 60124962D2 GBI:g8117242_000054 _edit.8313-8414 GBI:g8118985_000043 _edit.12301- 12444.comp GBI:g8117242_000054 _edit.4112-4228 GBI:g8117242_000054 _edit.10957-11181 5500380H1 (BRABDIR01) GBI:g8117242_000054 _edit.10616-10732	1 861 249 1205 975 539 1 1171 544 1441 1433 1 925 358 1735 1069 685 241 1717 907 1600	1156 1347 568 1775 1545 1534 662 1335 684 1599 1905 240 1068 492 1941 1170 810 357 1941 1119 1716
34	4588877CB1	1545	261-619, 1-193, 794-1071			
35	7472214CB1	1941	1483-1558, 1- 413, 495-616, 732-1149			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
35				GBI:g8117242_000054 _edit.8907-9011 GBI:g8117242_000054 _edit.6643-6756	1336 811	1440 924
36	7473053CB1	4971	3312-3482, 1- 1466, 4307-4971, 2184-2221	8035016H1 (SMCRUNE01) 6822202J1 (SINTNOR01) 6781747H1 (OVARDIR01) 8035016J1 (SMCRUNE01) 6824230H1 (SINTNOR01) 6894266H1 (BRAITDR03) 6777836H1 (OVARDIR01) 6908503H1 (PITUDIR01) 6908503J1 (PITUDIR01) 6823447H1 (SINTNOR01) 6823447J1 (SINTNOR01) 6006310F8 (FIBRUNT02) 4171959T6 (SINTNOT21) 5088860F6 (UTRSTMR01) GBI.lee4.edit	2315 2145 968 2979 2867 548 1601 1 1270 3525 4226 4501 3637 4461 1	2975 2877 1449 3643 3483 1157 2238 667 1830 4260 4829 4969 4287 4853 1404
37	7473347CB1	1404	126-633, 1013- 1404, 768-838			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
38	7474240CB1	4048	3023-4048, 1753- 2469, 1-920, 1593-1658, 2614- 2908, 1138-1367	71984804V1 GBI:g7656646_edit 71986624V1 1369 55055014H1 1 55037111J2 95 71983668V1 1371 GBI:g5923734_edit 2612 55037119J2 224 2502027F6 696 (ADRETUT05)	964 929 1369 1 95 1371 2612 224 696	1311 3418 1976 130 871 2043 4048 875 1235
39	7475338CB1	1539	1412-1539, 1- 328, 495-837, 922-1218	GBI:g7960701_000004 _edit.549-713 GBI:g7960701_000004 _edit.13381-13480 GBI:g7960701_000004 _edit.8755-8943 GBI:g7960701_000004 _edit.4292-4417 GBI:g7960701_000004 _edit.16237-16317 GBI:g7960701_000004 _edit.20107-20325 GBI:g7960701_000004 _edit.9989-10099 GBI:g7960701_000004 _edit.18748-18873 GBI:g7960701_000003 _edit.9783-9884 GBI:g7960701_000004 _edit.5251-5403 GBI:g7960701_000004 _edit.8384-8506 71906448V1 71753467V1 3351512F6 (PROSNOT28) 7761783J1 (THYMNOE02) 6934981R8 (SINTTMR02)	154 1015 715 313 1114 1321 904 1195 52 439 592 627 912 2185 1943 78	312 1113 903 438 1194 1539 1014 1320 153 591 714 1082 1539 2724 2570 860
40	7476747CB1	3114	1717-1870, 1- 503, 1468-1650	71906448V1 71753467V1 3351512F6 (PROSNOT28) 7761783J1 (THYMNOE02) 6934981R8 (SINTTMR02)	627 912 2185 1943 78	1082 1539 2724 2570 860

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
40				6389368H1 (PROSTMC01) 70536163V1 6934981F8 (SINTTMR02) GNN.g7712065_000012_002 7080657H1 (STOMTMR02) 5633289H1 (PLACFER01) g5746200 GBI.g2262095	1782 2575 1 452 838 639 1215 1	2075 3114 643 1922 1403 890 1473 2877
41	7477898CB1	2877	846-901, 1272-1378, 2319-2877			
42	7472728CB1	2820	1-1399, 2207-2229	55022826J1 55030210H1 4399366T6 (TESTTUT03) 55030274H1 g565876 55018149J1 FL203597_00001 GNN.g7263861_026.ed it	1138 403 2231 1482 2597 1907 712 1	1834 986 2777 2153 2820 2585 1807 1052
43	7474322CB1	1440	1-604, 714-768	GBI.g8081632_edit 71228887V1 70868623V1 3696546T6 (SININOT05) 70674954V1 1426382H1 (SINTBST01) 3696546F6 (SININOT05) 6828352H1 (SINTNOR01) 3699565H1 (SININOT05) 7700096H1 (KIDPTDE01) 70678552V1	1 1090 988 1833 1520 1224 799 530 1 250 1419	1440 1440 1385 2394 2091 1492 1381 1149 281 990 2055
44	5455621CB1	2394	1483-1686, 1-329, 838-1155, 2201-2235			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
45	7477248CB1	2890	1-58, 2739-2890, 2310-2349, 329- 1167	2777287H1 (OVRTUT03) 7977733H1 (LSUBDMC01) 7678168J1 (NOSETUE01) 7611941J1 (KIDCTME01) 6590507H1 (TLYMUNT03) 2701794F6 (OVRTUT10) 2544096F6 (UTRSNOT11) 60117044D2 5020832H1 (OVARNON03) 7662529H1 (UTRSTME01) 4762728F6 (PLACNOT05) g2264624 6264977H1 (MCLDTXN03) 2944004F6 (BRAITUT23) 6610392H2 (MUSTTMC01) GNN.g7328818_000024 _002.edit 7035078H1 (SINTFER03) 7620248J1 (HEARFEE03) 496537H1 (HNT2NOT01) 6264427T8 (MCLDTXN03) 6264427F8 (MCLDTXN03)	2250 841 1271 2273 179 1208 1732 1 2195 526 872 2268 1210 2790 3306 2145 1 2431 2329 453 170	2498 1427 1827 2890 672 1741 2252 431 2471 926 1387 2446 1797 3531 3926 2648 440 3039 2487 1174 842
46	2944004CB1	3926	3338-3365, 1- 687, 1222-2267			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
46				7673654H1 (FIBPFEC01)	1733	2239
47	3046849CB1	2135	2072-2135, 596- 711, 1014-1263	8262790U1 71896642V1 71247870V1 FL3046849_g6815043_ 000004_g183298	1383 1 1050 51	2135 592 1736 1520
48	4538363CB1	2637	1-183, 1575- 1680, 2094-2637	FL4538363_g3126781_ g520469 71401405V1 70857895V1 7727961J1 (UTRC DIE01) 70857789V1 g5689372_edit g3801917	1 1766 416 3284 566 1092 1	1917 2637 1035 3783 1109 3361 452
49	6427460CB1	3783	985-1833, 2687- 3204	GBI.g8568959_edit_3 g6140313 5819744F7 (PROSTUS23) g5920552 55049678J1 FL7476949_g6714723_ g338053 4669722H1 (SINTNOT24)	1119 482 168 1 862 1	2105 951 479 488 1359 2046
50	7474127CB1	2105	1078-2105	1233-1356, 1- 117, 2047-2069, 347-503, 1536- 1844 2833-3018, 1869- 2121, 3707-4245, 1-252, 982-1239, 289-357	1801	2069
51	7476949CB1	2069		71660072V1 71657569V1 7633968J1 (SINTDIE01) 6440145F8 (BRAENOT02) 71664080V1 GBI.g8567478.edit 71660176V1 71662066V1 2605539F6 (LUNG T07)	2404 3106 2579 938 3228 1 3773 1802 433	3156 3854 3175 1087 3891 2547 4245 2475 939
52	7477249CB1	4245		71659261V1 3825558H1 (BRAHCT02)	1690 1179	2437 1270

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
52				7765571H1 (UPRTTUE01)	1	693
53	7477720CB1	2124	1-936, 1200- 1488, 1982-2124, 1562-1745	5675861H1 FL7477720_g5836195_ g205709	1427 1	1716 2124
54	7477852CB1	2195	1-418, 1899-2195	GBI.g8748866.edit	1	2195
55	1471717CB1	2055	206-768, 881- 931, 1155-1323	70464956V1 72277206V1 70469664V1 GNN.g7109510_000068 _002.edit GBI.g8039708_50_63_ 62_56.edit 6540941H1 (LNODNON02)	492 1 939 772 238 1571	994 297 1582 1500 897 2055
56	3874406CB1	4727	1-1299, 1576- 1632, 2550-3619, 2014-2192	70466394V1 71793833V1 55052105J1 71798347V1 71798870V1 55058313J1 55051482J1 FL3874406_g3810670_ g4240130_3_3-4 55068154H1 3133035F6 (SMCCNOT01)	1035 4117 1673 3620 3575 1380 2475 482 2223 1	1616 4727 2128 4358 4244 2125 3134 744 2741 605
57	4599654CB1	3852	1-335, 2014-3231	55058329H1 55068182J1 71795307V1 8016331J1 (BMARTXE01) 71040001V1 8041905H1 (OVRTUE01) 55062505H1 g7959336_CD 6772024J1 (BRAUNOR01) 55064208J1	723 2048 2902 1778 3348 1666 660 349 1 1118	1528 2685 3593 2424 3852 2352 1233 2540 623 1718

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
57				6617183H2 (BRAXTDR14) 6195941H1 (PITUNON01) 71909238V1 2216896F6 (SINTFET03) 71042073V1 7431853H1 (UTPMTMR02) GNN:g4375937_004_ed it 6426880H1 (LUNGNON07) 6781142H1 (OVARDIR01) 2645767H1 (OVARNOT09) 71704421V1 7726210H1 (THYRDIE01) 7721710J2 (THYRDIE01) 6340173F8 (BRANDIN01) 71704256V1 7757131H1 (SPLNTUE01) GNN.g7711543_000002 _002.edit 7464813H1 (LIVRFEE04) 71703676V1 7760618H1 (THYMN0E02) 71970086V1 7462584H1 (LIVRFEE04) 7760618J1 (THYMN0E02) 71762287V1	2981 2823 1225 2474 2276 1211 1 814 224 128 6240 1885 2696 5516 3025 2408 198 544 3250 2183 5817 1 1251 4313	3530 3458 1747 2923 2745 1917 1845 1336 941 394 6791 2502 3232 6222 3734 3093 2751 696 3947 2676 6525 578 1983 4879
58	5047435CB1	1917	1-238, 1162-1474			
59	7475603CB1	6791	1-3283, 5952- 6101, 3793-4761			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
59				7724639H1 (THYRDIE01) 55052451J1 7739867H1 (THYMNOE01) 6879936H1 (UTRSTR02) 55058371H1 GBI.g8346195_edit GBI.g8052096_edit 8104845H1 (MIXDDIE02) GBI.g8518014_edit g1081430 168827H1 (LIVRNOT01) 55064792J1 55072770H1 GNN.g6498074_012.ed it 087510H1 (LIVRNOT01) g751568 55055559H1 55045003H2 g5361744 GBI.g8118965_000015 _000006_000001_0000 10_000003.edit g751568 55049235H1 GBI.g8018151_000001 .edit GBI.g6433826_000001 .edit 55063069J1 g669271	951 4792 5131 697 3850 1765 1132 2822 1 1036 65 1 495 1321 314 1336 16 1 908 602 10_000003.edit 1763 556 1799 1172 1 1799	1545 5698 5794 1054 4747 5214 1839 3367 1266 1525 406 209 1110 1818 574 1773 699 697 1109 2245 2000 2200 1287 3196 2052 850 2106
60	7477845CB1	5214	2390-4599, 645- 1796			
61	168827CB1	1818	1-281, 796-912			
62	7472734CB1	2245	1223-1339, 1-710			
63	7473473CB1	3196	1-376, 460-1796			

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
64	7477725CB1	1602	1072-1602	7455614H1 (LIVRTUE01)	416	835
				4288148H1 (LIVRDIR01)	112	257
				GBI.g8131631_000007 _000005.edit	1	1602
				g2656651	829	1084

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
33	3474673CB1	LJNLJMT01
34	4588877CB1	LJNLJMT01
35	7472214CB1	BRAENOT04
36	7473053CB1	SINTNOR01
38	7474240CB1	ADRETTU05
39	7475338CB1	SINTNOT18
40	7476747CB1	SINTTMR02
42	7472728CB1	TESITTU03
43	7474322CB1	SINTBST01
44	5455621CB1	SININOT05
45	7477248CB1	UTRSNOT11
46	2944004CB1	MCLDTXN03
47	3046849CB1	HNT2AGT01
48	4538363CB1	PANCNOT07
49	6427460CB1	BRAUNOR01
50	7474127CB1	PROSTUS23
51	7476949CB1	COLNPMC01
52	7477249CB1	COLNPOT01
55	1471717CB1	OVARDIT01
56	3874406CB1	LIVRDIR01
57	4599654CB1	LUNGNOT23
58	5047435CB1	OVARDIR01
59	7475603CB1	THYRDIE01
60	7477845CB1	MIXDDIE02
61	168827CB1	LIVRNOT01
64	7477725CB1	LIVRTUE01

Table 6

Library	Vector	Library Description
ADRETUT05	pINCY	Library was constructed using RNA isolated from adrenal tumor tissue removed from a 52-year-old Caucasian female during a unilateral adrenalectomy. Pathology indicated a pheochromocytoma.
BRAENOT04	pINCY	Library was constructed using RNA isolated from inferior parietal cortex tissue removed from the brain of a 35-year-old Caucasian male who died from cardiac failure. Pathology indicated moderate leptomeningeal fibrosis and multiple microinfarctions of the cerebral neocortex. Patient history included dilated cardiomyopathy, congestive heart failure, cardiomegaly and an enlarged spleen and liver.
BRAUNOR01	pINCY	This random primed library was constructed using RNA isolated from striatum, globus pallidus and posterior putamen tissue removed from an 81-year-old Caucasian female who died from a hemorrhage and ruptured thoracic aorta due to atherosclerosis. Pathology indicated moderate atherosclerosis involving the internal carotids, bilaterally; microscopic infarcts of the frontal cortex and hippocampus; and scattered diffuse amyloid plaques and neurofibrillary tangles, consistent with age. Grossly, the leptomeninges showed only mild thickening and hyalinization along the superior sagittal sinus. The remainder of the leptomeninges was thin and contained some congested blood vessels. Mild atrophy was found mostly in the frontal poles and lobes, and temporal lobes, bilaterally. Microscopically, there were pairs of Alzheimer type II astrocytes within the deep layers of the neocortex. There was increased satellitosis around neurons in the deep gray matter in the middle frontal cortex. The amygdala contained rare diffuse plaques and neurofibrillary tangles. The posterior hippocampus contained a microscopic area of cystic cavitation with hemosiderin-laden macrophages surrounded by reactive gliosis. Patient history included sepsis, cholangitis, post-operative atelectasis, pneumonia CAD, cardiomegaly due to left ventricular hypertrophy, splenomegaly, arteriolonephrosclerosis, nodular colloidal goiter, emphysema, CHF, hypothyroidism, and peripheral vascular disease.
COLNPOT01	pINCY	Library was constructed using RNA isolated from colon polyp tissue removed from a 40-year-old Caucasian female during a total colectomy. Pathology indicated an inflammatory pseudopolyp; this tissue was associated with a focally invasive grade 2 adenocarcinoma and multiple tubovillous adenomas. Patient history included a benign neoplasm of the bowel.
COLNTMC01	pINCY	This large size-fractionated library was constructed using pooled cDNA from three different donors. cDNA was generated using mRNA isolated from colon epithelium tissue removed from a 13-year-old Caucasian female (donor A) who died from a motor vehicle accident; from ascending colon removed from a 29-year-old female (donor

Table 6 (cont.)

Library	Vector	Library Description
		B); and from colon tissue removed from the appendix of a 37-year-old Black female (donor C) during myomectomy, dilation and curettage, right fimbrial region biopsy, and incidental appendectomy. Pathology for donor B indicated the proximal and distal resection margins of small bowel and colon away from the mass lesion were uninvolved by lymphoma. Pathology for donor C indicated an unremarkable appendix. Pathology for the matched tumor tissue (donor B) indicated malignant lymphoma, small cell, non-cleaved (Burkitt's lymphoma, B-cell phenotype), forming a polypoid mass in the region of the ileocecal valve, associated with intussusception and obstruction clinically. The liver and multiple (3 of 12) ileocecal region lymph nodes were also involved by lymphoma. Pathology for the associated tumor tissue (donor C) indicated multiple uterine leiomyomata. Donor C presented with deficiency anemia, an umbilical hernia, and premenopausal menorrhagia. Patient history included sarcoidosis of the lung.
HNT2AGT01	PBLUESCRIPT	Library was constructed at Stratagene (STR937233), using RNA isolated from the hNT2 cell line derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor. Cells were treated with retinoic acid for 5 weeks and with mitotic inhibitors for two weeks and allowed to mature for an additional 4 weeks in conditioned medium.
LIVRDIR01	pINCY	The library was constructed using RNA isolated from diseased liver tissue removed from a 63-year-old Caucasian female during a liver transplant. Patient history included primary biliary cirrhosis diagnosed in 1989. Serology was positive for anti-mitochondrial antibody.
LIVRNOT01	PBLUESCRIPT	Library was constructed at Stratagene, using RNA isolated from the liver tissue of a 49-year-old male.
LIVRTUE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from liver tumor tissue removed from a 72-year-old Caucasian male during partial hepatectomy. Pathology indicated metastatic grade 2 (of 4) neuroendocrine carcinoma forming a mass. The patient presented with metastatic liver cancer. Patient history included benign hypertension, type I diabetes, prostatic hyperplasia, prostate cancer, alcohol abuse in remission, and tobacco abuse in remission. Previous surgeries included destruction of a pancreatic lesion, closed prostatic biopsy, transurethral prostatectomy, removal of bilateral testes and total splenectomy. Patient medications included Eulexin, Hytrin, Proscar, Ecotrin, and insulin. Family history included atherosclerotic coronary artery disease and acute myocardial infarction in the mother; atherosclerotic coronary artery disease and type II diabetes in the father.
LUNGNOT23	pINCY	Library was constructed using RNA isolated from left lobe lung tissue removed from

Table 6 (cont.)

Library	Vector	Library Description
		a 58-year-old Caucasian male. Pathology for the associated tumor tissue indicated metastatic grade 3 (of 4) osteosarcoma. Patient history included soft tissue cancer, secondary cancer of the lung, prostate cancer, and an acute duodenal ulcer with hemorrhage. Family history included prostate cancer, breast cancer, and acute leukemia.
LUNLTMT01	pINCY	The library was constructed using RNA isolated from right middle lobe lung tissue removed from a 63-year-old Caucasian female during a segmental lung resection. Pathology for the associated tumor tissue indicated grade 3 adenocarcinoma in the right lower lobe and right middle lobe that infiltrated the parietal pleural surface. Metastatic grade 3 adenocarcinoma was found in the diaphragm. The lymph nodes contained metastatic grade 3 adenocarcinoma and involved the superior mediastinal and inferior mediastinal lymph nodes. Patient history included hyperlipidemia. Family history included benign hypertension, cerebrovascular disease, breast cancer, and hyperlipidemia.
MCLDTXN03	pINCY	This normalized dendritic cell library was constructed from one million independent clones from a pool of two derived dendritic cell libraries. Starting libraries were constructed using RNA isolated from untreated and treated derived dendritic cells from umbilical cord blood CD34+ precursor cells removed from a male. The cells were derived with granulocyte/macrophage colony stimulating factor (GM-CSF), tumor necrosis factor alpha (TNF alpha), and stem cell factor (SCF). The GM-CSF was added at time 0 at 100 ng/ml, the TNF alpha was added at time 0 at 2.5 ng/ml, and the SCF was added at time 0 at 25 ng/ml. Incubation time was 13 days. The treated cells were then exposed to phorbol myristate acetate (PMA), and Ionomycin. The PMA and Ionomycin were added at 13 days for five hours. The library was normalized in two rounds using conditions adapted from Soares et al., PNAS (1994) 91:9228-9232 and Bonaldo et al., Genome Research (1996) 6:791, except that a significantly longer (48 hours/round) reannealing hybridization was used.
MIXDDIE02	PBK-CMV	This 5' biased random primed library was constructed using pooled cDNA from seven donors. cDNA was generated using mRNA isolated from brain tissue removed from two Caucasian male fetuses who died after 23 weeks gestation from hypoplastic left heart (A) and prematurity (B); from posterior hippocampus from a 55-year-old male who died from COPD (C); from cerebellum, corpus callosum, thalamus and temporal lobe tissue from a 57-year-old Caucasian male who died from a CVA (D); from dentate nucleus and vermis from an 82-year-old Caucasian male who died from a myocardial infarction (E); from pituitary gland from a 74-year-old Caucasian female who died from a myocardial infarction (F) and vermis tissue from a 77-year-old Caucasian female who died from pneumonia (G). For donor C, pathology indicated

Table 6 (cont.)

Library	Vector	Library Description
OVARDIR01	PCDNA2.1	<p>mild lateral ventricular enlargement. For donor F, pathology indicated moderate Alzheimer's disease, recent multiple infarctions involving left thalamus, left parietal and occipital lobes (microscopic) and right cerebellum (gross), mild atherosclerosis involving middle cerebral arteries bilaterally and mild cerebral amyloid angiopathy. For donor G, pathology indicated severe Alzheimer's disease, mild atherosclerosis involving the middle cerebral and basilar arteries, and cerebral atrophy consistent with Alzheimer's disease. For donor D, patient history included Huntington's chorea. Donor E was taking nitroglycerin and dopamine; donor F was taking Lopressor, heparin, ceftriaxone, captopril, Isordil, nitroglycerin, Clinoril, Ecotrin and tacrine; and donor G was taking insulin.</p> <p>This random primed library was constructed using RNA isolated from right ovary tissue removed from a 45-year-old Caucasian female during total abdominal hysterectomy, bilateral salpingo-oophorectomy, vaginal suspension and fixation, and incidental appendectomy. Pathology indicated stromal hyperthecosis of the right and left ovaries. Pathology for the matched tumor tissue indicated a dermoid cyst (benign cystic teratoma) in the left ovary. Multiple (3) intramural leiomyomata were identified. The cervix showed squamous metaplasia. Patient history included metrorrhagia, female stress incontinence, alopecia, depressive disorder, pneumonia, normal delivery, and deficiency anemia. Family history included benign hypertension, atherosclerotic coronary artery disease, hyperlipidemia, and primary tuberculous complex.</p>
OVARDIT01	pINCY	<p>Library was constructed using RNA isolated from diseased ovary tissue removed from a 39-year-old Caucasian female during total abdominal hysterectomy, bilateral salpingo-oophorectomy, dilation and curettage, partial colectomy, incidental appendectomy, and temporary colostomy. Pathology indicated the right and left adnexa were extensively involved by endometriosis. Endometriosis also involved the anterior and posterior serosal surfaces of the uterus and the cul-de-sac and the mesentery and muscularis propria of the sigmoid colon. Pathology for the associated tumor tissue indicated multiple (3) intramural, 1 subserosal) leiomyomata. Family history included hyperlipidemia, benign hypertension, atherosclerotic coronary artery disease, depressive disorder, brain cancer, and type II diabetes.</p>
PANCN07	pINCY	Library was constructed using RNA isolated from the pancreatic tissue of a Caucasian male fetus, who died at 23 weeks' gestation.
PROSTUS23	pINCY	This subtracted prostate tumor library was constructed using 10 million clones from a pooled prostate tumor library that was subjected to 2 rounds of subtractive hybridization with 10 million clones from a pooled prostate tissue library. The

Table 6 (cont.)

Library	Vector	Library Description
SININOT05	pINCY	starting library for subtraction was constructed by pooling equal numbers of clones from 4 prostate tumor libraries using mRNA isolated from prostate tumor removed from Caucasian males at ages 58 (A), 61 (B), 66 (C), and 68 (D) during prostatectomy with lymph node excision. Pathology indicated adenocarcinoma in all donors. History included elevated PSA, induration and tobacco abuse in donor A; elevated PSA, induration, prostate hyperplasia, renal failure, osteoarthritis, renal artery stenosis, benign HTN, thrombocytopenia, hyperlipidemia, tobacco/alcohol abuse and hepatitis C (carrier) in donor B; elevated PSA, induration, and tobacco abuse in donor C; and elevated PSA, induration, hypercholesterolemia, and kidney calculus in donor D. The hybridization probe for subtraction was constructed by pooling equal numbers of cDNA clones from 3 prostate tissue libraries derived from prostate tissue, prostate epithelial cells, and fibroblasts from prostate stroma from 3 different donors. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR 19 (1991):1954 and Bonaldo, et al. Genome Research 6 (1996):791.
SINTBST01	pINCY	Library was constructed using RNA isolated from ileum tissue obtained from a 30-year-old Caucasian female during partial colectomy, open liver biopsy, incidental appendectomy, and permanent colostomy. Patient history included endometriosis. Family history included hyperlipidemia, anxiety, and upper lobe lung cancer, stomach cancer, liver cancer, and cirrhosis.
SINTNOR01	PCDNA2.1	Library was constructed using RNA isolated from the ileum tissue of an 18-year-old Caucasian female. The ileum tissue, along with the cecum and appendix, were removed during bowel anastomosis. Pathology indicated Crohn's disease of the ileum, involving 15 cm of the small bowel. The cecum and appendix were unremarkable, and the margins were uninvolved. The patient presented with abdominal pain and regional enteritis. Patient history included osteoporosis of the vertebra and abnormal blood chemistry. Patient medications included Prilosec (omeprazole), Pentasa (mesalamine), amoxicillin, and multivitamins. Family history included cerebrovascular disease and atherosclerotic coronary artery disease.
SINTNOT18	pINCY	This random primed library was constructed using RNA isolated from small intestine tissue removed from a 31-year-old Caucasian female during Roux-en-Y gastric bypass. Patient history included clinical obesity.
SINTTMR02	PCDNA2.1	Library was constructed using RNA isolated from small intestine tissue obtained from a 59-year-old male.
		This random primed library was constructed using RNA isolated from small intestine tissue removed from a 59-year-old male. Pathology for the matched tumor tissue

Table 6 (cont.)

Library	Vector	Library Description
TESTTUT03	pINCY	indicated multiple (9) carcinoid tumors, grade 1, in the small bowel. The largest tumor was associated with a large mesenteric mass. Multiple convoluted segments of bowel were adhered to the tumor. A single (1 of 13) regional lymph node was positive for malignancy. The peritoneal biopsy indicated focal fat necrosis. Library was constructed using RNA isolated from right testicular tumor tissue removed from a 45-year-old Caucasian male during a unilateral orchiectomy. Pathology indicated seminoma. Patient history included hyperlipidemia and stomach ulcer. Family history included cerebrovascular disease, skin cancer, hyperlipidemia, acute myocardial infarction, and atherosclerotic coronary artery disease.
THYRDIE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from diseased thyroid tissue removed from a 22-year-old Caucasian female during closed thyroid biopsy, partial thyroidectomy, and regional lymph node excision. Pathology indicated adenomatous hyperplasia. The patient presented with malignant neoplasm of the thyroid. Patient history included normal delivery, alcohol abuse, and tobacco abuse. Previous surgeries included myringotomy. Patient medications included an unspecified type of birth control pills. Family history included hyperlipidemia and depressive disorder in the mother; and benign hypertension, congestive heart failure, and chronic leukemia in the grandparent(s).
UTRSNOT11	pINCY	Library was constructed using RNA isolated from uterine myometrial tissue removed from a 43-year-old female during a vaginal hysterectomy and removal of the fallopian tubes and ovaries. Pathology for the associated tumor tissue indicated that the myometrium contained an intramural and a submucosal leiomyoma. Family history included benign hypertension, hyperlipidemia, colon cancer, type II diabetes, and atherosclerotic coronary artery disease.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) <i>J. Mol. Biol.</i> 215:403-410; Altschul, S.F. et al. (1997) <i>Nucleic Acids Res.</i> 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less <i>Full Length sequences</i> : Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) <i>Proc. Natl. Acad. Sci. USA</i> 85:2444-2448; Pearson, W.R. (1990) <i>Methods Enzymol.</i> 183:63-98; and Smith, T.F. and M.S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489.	ESTs: fasta E value=1.06E-6 <i>Assembled ESTs</i> : fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less <i>Full Length sequences</i> : fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) <i>Nucleic Acids Res.</i> 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) <i>Methods Enzymol.</i> 266:88-105; and Attwood, T.K. et al. (1997) <i>J. Chem. Inf. Comput. Sci.</i> 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) <i>J. Mol. Biol.</i> 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) <i>Nucleic Acids Res.</i> 26:320-322; Durbin, R. et al. (1998) <i>Our World View</i> , in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less <i>Signal peptide hits</i> : Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score ₂ GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:

a) a polypeptide comprising an amino acid sequence selected from the group consisting of

5 SEQ ID NO:1-32,

b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32,

c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and

10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-

32.

15

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:33-64.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

25

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

30 9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

35 b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

11. An isolated polynucleotide selected from the group consisting of:

a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting

5 of SEQ ID NO:33-64,

b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64,

c) a polynucleotide complementary to a polynucleotide of a),

d) a polynucleotide complementary to a polynucleotide of b), and

10 e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization
20 complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

18. A method for treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional TRICH, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,

b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and

c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,

b) detecting altered expression of the target polynucleotide, and

c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

a) treating a biological sample containing nucleic acids with the test compound;

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;

c) quantifying the amount of hybridization complex; and

d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29. A diagnostic test for a condition or disease associated with the expression of TRICH in a biological sample comprising the steps of:

- 5 a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

- 10 a) a chimeric antibody,
b) a single chain antibody,
c) a Fab fragment,
d) a F(ab')₂ fragment, or
e) a humanized antibody.

15 31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of TRICH in a subject, comprising administering to said subject an effective amount of the composition of claim 31.

33. A composition of claim 31, wherein the antibody is labeled.

25 34. A method of diagnosing a condition or disease associated with the expression of TRICH in a subject, comprising administering to said subject an effective amount of the composition of claim 33.

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

- 30 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, or an immunogenic fragment thereof, under conditions to elicit an antibody response;
b) isolating antibodies from said animal; and

c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

5 36. An antibody produced by a method of claim 35.

37. A composition comprising the antibody of claim 36 and a suitable carrier.

10 38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10 comprising:

a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, or an immunogenic fragment thereof, under conditions to elicit an antibody response;

b) isolating antibody producing cells from the animal;

15 c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells;

d) culturing the hybridoma cells; and

e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

20

39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

25 41. The antibody of claim 10, wherein the antibody is produced by screening a Fab expression library.

42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

30

43. A method for detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32 in a sample, comprising the steps of:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32 in the sample.

5 44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32 from a sample, the method comprising:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

10 b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

15 46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

20 49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

25 51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.

30 54. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.

55. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.

57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.
58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.
- 5 59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.
60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.
61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.
- 10 62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.
63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.
64. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.
- 15 65. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.
66. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:22.
- 20 67. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:23.
68. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:24.
- 25 69. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:25.
70. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:26.
71. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:27.
- 30 72. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:28.
73. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:29.

74. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:30.

75. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:31.

5 76. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:32.

77. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:33.

10 78. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:34.

79. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:35.

15 80. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:36.

20 81. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:37.

82. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:38.

25 83. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:39.

84. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:40.

30 85. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:41.

86. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:42.

5 87. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:43.

88. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:44.

10 89. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:45.

90. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:46.

15 91. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:47.

20 92. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:48.

93. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:49.

25 94. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:50.

95. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:51.

30 96. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:52.

97. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:53.

5 98. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:54.

99. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:55.

10 100. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:56.

101. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:57.

15 102. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:58.

20 103. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:59.

104. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:60.

25 105. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:61.

106. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:62.

30 107. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:63.

108. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:64.

<110> INCYTE GENOMICS, INC.
 RAUMANN, Brigitte E.
 THORNTON, Michael
 DING, Li
 YUE, Henry
 TANG, Y.Tom
 HARLAND, Lee
 BURFORD, Neil
 GREENE, Barrie D.
 SANJANWALA, Madhu S.
 BAUGHN, Mariah R.
 YAO, Monique G.
 YANG, Junming
 PATTERSON, Chandra
 GANDHI, Ameena R.
 HAFALIA, April J.A.
 TRIBOULEY, Catherine M.
 WALIA, Narinder K.
 AU-YOUNG, Janice
 WALSH, Roderick T.
 RAMKUMAR, Jayalaxmi
 LU, Yan
 LU, Dyung Aina M.
 AZIMZAI, Yalda
 LAL, Preeti
 ELLIOTT, Vicki S.
 NGUYEN, Danniel B.
 XU, Yuming
 SEILHAMER, Jeffrey J.
 BOROWSKY, Mark L.
 KHAN, Farrah A.
 KEARNEY, Liam
 THANGAVELU, Kavitha
 DAS, Debopriya
 POLICKY, Jennifer L.

<120> TRANSPORTERS AND ION CHANNELS

<130> PI-0149 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/216,547; 60/218,232; 60/220,112; 60/221,839

<151> 2000-07-07; 2000-07-14; 2000-07-21; 2000-07-28

<160> 64

<170> PERL Program

<210> 1

<211> 332

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3474673CD1

<400> 1

Met	Tyr	Arg	Pro	Arg	Ala	Arg	Ala	Ala	Pro	Glu	Gly	Arg	Val	Arg
1				5					10					15
Gly	Cys	Ala	Val	Pro	Ser	Thr	Val	Leu	Leu	Leu	Leu	Ala	Tyr	Leu
				20					25					30
Ala	Tyr	Leu	Ala	Leu	Gly	Thr	Gly	Val	Phe	Trp	Thr	Leu	Glu	Gly
				35					40					45
Arg	Ala	Ala	Gln	Asp	Ser	Ser	Arg	Ser	Phe	Gln	Arg	Asp	Lys	Trp

	50		55		60
Glu Leu Leu Gln Asn	Phe Thr Cys Leu Asp	Arg Pro Ala Leu Asp			
65	70	75			
Ser Leu Ile Arg Asp	Val Val Gln Ala Tyr	Lys Asn Gly Ala Ser			
80	85	90			
Leu Leu Ser Asn Thr	Thr Ser Met Gly Arg	Trp Glu Leu Val Gly			
95	100	105			
Ser Phe Phe Phe Ser	Val Ser Thr Ile Thr	Thr Ile Gly Tyr Gly			
110	115	120			
Asn Leu Ser Pro Asn	Thr Met Ala Ala Arg	Leu Phe Cys Ile Phe			
125	130	135			
Phe Ala Leu Val Gly	Ile Pro Leu Asn Leu	Val Val Leu Asn Arg			
140	145	150			
Leu Gly His Leu Met	Gln Gln Gly Val Asn	His Trp Ala Ser Arg			
155	160	165			
Leu Gly Gly Thr Trp	Gln Asp Pro Asp Lys	Ala Arg Trp Leu Ala			
170	175	180			
Gly Ser Gly Ala Leu	Leu Ser Gly Leu Leu	Leu Phe Leu Leu Leu			
185	190	195			
Pro Pro Leu Leu Phe	Ser His Met Glu Gly	Trp Ser Tyr Thr Glu			
200	205	210			
Gly Phe Tyr Phe Ala	Phe Ile Thr Leu Ser	Thr Val Gly Phe Gly			
215	220	225			
Asp Tyr Val Ile Gly	Met Asn Pro Ser Gln	Arg Tyr Pro Leu Trp			
230	235	240			
Tyr Lys Asn Met Val	Ser Leu Trp Ile Leu	Phe Gly Met Ala Trp			
245	250	255			
Leu Ala Leu Ile Ile	Lys Leu Ile Leu Ser	Gln Leu Glu Thr Pro			
260	265	270			
Gly Arg Val Cys Ser	Cys Cys His His Ser	Ser Lys Glu Asp Phe			
275	280	285			
Lys Ser Gln Ser Trp	Arg Gln Gly Pro Asp	Arg Glu Pro Glu Ser			
290	295	300			
His Ser Pro Gln Gln	Gly Cys Tyr Pro Glu	Gly Pro Met Gly Ile			
305	310	315			
Ile Gln His Leu Glu	Pro Ser Ala His Ala	Ala Gly Cys Gly Lys			
320	325	330			
Asp Ser					

<210> 2

<211> 226

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4588877CD1

<400> 2

Met Val Glu Met Gly	Trp Asp Trp Ala Asp	Arg Lys Asp Met Arg			
1	5	10	15		
His Arg Leu Gln Ala	Gly Asn Leu Glu Asn	Thr Asp Gln Val Lys			
20	25	30			
Ser Pro Leu Leu Thr	Gly Asp Ser Ser Gly	Leu Pro Pro Ala Pro			
35	40	45			
Ser Ala Pro Thr His	Gly Val Lys Ala Ser	Gly Gly Leu Gly Thr			
50	55	60			
Ile Leu His Pro Gln	Asp Pro Asp Lys Ala	Arg Trp Leu Ala Gly			
65	70	75			
Ser Gly Ala Leu Leu	Ser Gly Leu Leu Leu	Phe Leu Leu Leu Pro			
80	85	90			
Pro Leu Leu Phe Ser	His Met Glu Gly Trp	Ser Tyr Thr Glu Gly			
95	100	105			
Phe Tyr Phe Ala Phe	Ile Thr Leu Ser Thr	Val Gly Phe Gly Asp			
110	115	120			
Tyr Val Ile Gly Met	Asn Pro Ser Gln Arg	Tyr Pro Leu Trp Tyr			

Lys	Asn	Met	Val	125	Ser	Leu	Trp	Ile	Leu	130	Phe	Gly	Met	Ala	Trp	Leu	135
				140						145							150
Ala	Leu	Ile	Ile	155	Lys	Leu	Ile	Leu	Ser	160	Gln	Leu	Glu	Thr	Pro	Gly	165
Arg	Val	Cys	Ser	170	Cys	Cys	His	His	Ser	175	Ser	Lys	Glu	Asp	Phe	Lys	180
Ser	Gln	Ser	Trp	185	Arg	Gln	Gly	Pro	Asp	190	Arg	Glu	Pro	Glu	Ser	His	195
Ser	Pro	Gln	Gln	200	Gly	Cys	Tyr	Pro	Glu	205	Gly	Pro	Met	Gly	Ile	Ile	210
Gln	His	Leu	Glu	215	Pro	Ser	Ala	His	Ala	220	Ala	Gly	Cys	Gly	Lys	Asp	225
Ser																	

<210> 3

<211> 646

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472214CD1

<400> 3

Met	Ala	Glu	Lys	Ala	Leu	Glu	Ala	Val	Gly	Cys	Gly	Leu	Gly	Pro			
1				5					10					15			
Gly	Ala	Val	Ala	Met	Ala	Val	Thr	Leu	Glu	Asp	Gly	Ala	Glu	Pro			
				20					25					30			
Pro	Val	Leu	Thr	Thr	His	Leu	Lys	Lys	Val	Glu	Asn	His	Ile	Thr			
				35					40					45			
Glu	Ala	Gln	Arg	Phe	Ser	His	Leu	Pro	Lys	Arg	Ser	Ala	Val	Asp			
				50					55					60			
Ile	Glu	Phe	Val	Glu	Leu	Ser	Tyr	Ser	Val	Arg	Glu	Gly	Pro	Cys			
				65					70					75			
Trp	Arg	Lys	Arg	Gly	Tyr	Lys	Thr	Leu	Leu	Lys	Cys	Leu	Ser	Gly			
				80					85					90			
Lys	Phe	Cys	Arg	Arg	Glu	Leu	Ile	Gly	Ile	Met	Gly	Pro	Ser	Gly			
				95					100					105			
Ala	Gly	Lys	Ser	Thr	Phe	Met	Asn	Ile	Leu	Ala	Gly	Tyr	Arg	Glu			
				110					115					120			
Ser	Gly	Met	Lys	Gly	Gln	Ile	Leu	Val	Asn	Gly	Arg	Pro	Arg	Glu			
				125					130					135			
Leu	Arg	Thr	Phe	Arg	Lys	Met	Ser	Cys	Tyr	Ile	Met	Gln	Asp	Asp			
				140					145					150			
Met	Leu	Leu	Pro	His	Leu	Thr	Val	Leu	Glu	Ala	Met	Met	Val	Ser			
				155					160					165			
Ala	Asn	Leu	Asn	Leu	Thr	Glu	Asn	Pro	Asp	Val	Lys	Asn	Asp	Leu			
				170					175					180			
Val	Thr	Glu	Ile	Leu	Thr	Ala	Leu	Gly	Leu	Met	Ser	Cys	Ser	His			
				185					190					195			
Thr	Arg	Thr	Ala	Leu	Leu	Ser	Gly	Gly	Gln	Arg	Lys	Arg	Leu	Ala			
				200					205					210			
Ile	Ala	Leu	Glu	Leu	Val	Asn	Asn	Pro	Pro	Val	Met	Phe	Phe	Asp			
				215					220					225			
Glu	Pro	Thr	Ser	Gly	Leu	Asp	Ser	Ala	Ser	Cys	Phe	Gln	Val	Val			
				230					235					240			
Ser	Leu	Met	Lys	Ser	Leu	Ala	Gln	Gly	Gly	Arg	Thr	Ile	Ile	Cys			
				245					250					255			
Thr	Ile	His	Gln	Pro	Ser	Ala	Lys	Leu	Phe	Glu	Met	Phe	Asp	Lys			
				260					265					270			
Leu	Tyr	Ile	Leu	Ser	Gln	Gly	Gln	Cys	Ile	Phe	Lys	Gly	Val	Val			
				275					280					285			
Thr	Asn	Leu	Ile	Pro	Tyr	Leu	Lys	Gly	Leu	Gly	Leu	His	Cys	Pro			
				290					295					300			
Thr	Tyr	His	Asn	Pro	Ala	Asp	Phe	Val	Ile	Glu	Val	Ala	Ser	Gly			

Glu Tyr Gly Asp	305	Leu Asn Pro Met Leu	310	Phe Arg Ala Val Gln	315
Gly Leu Cys Ala	320	Met Ala Glu Lys Lys	325	Ser Ser Pro Glu Lys	330
Glu Val Pro Ala	335	Pro Cys Pro Pro Cys	340	Pro Pro Glu Val Asp	345
Ile Glu Ser His	350	Thr Phe Ala Thr Ser	355	Thr Leu Thr Gln Phe	360
Ile Leu Phe Lys	365	Arg Thr Phe Leu Ser	370	Ile Leu Arg Asp Thr	375
Leu Thr His Leu	380	Arg Phe Met Ser His	385	Val Val Ile Gly Val	390
Ile Gly Leu Leu	395	Tyr Leu His Ile Gly	400	Asp Asp Ala Ser Lys	405
Phe Asn Asn Thr	410	Gly Cys Leu Phe Phe	415	Ser Met Leu Phe Leu	420
Phe Ala Ala Leu	425	Met Pro Thr Val Leu	430	Thr Val Pro Leu Glu	435
Ala Val Phe Met	440	Arg Glu His Leu Asn	445	Tyr Trp Tyr Ser Leu	450
Ala Tyr Tyr Leu	455	Ala Lys Thr Met Ala	460	Asp Val Pro Phe Gln	465
Val Cys Pro Val	470	Val Tyr Cys Ser Ile	475	Val Tyr Trp Met Thr	480
Gln Pro Ala Glu	485	Thr Ser Arg Phe Leu	490	Leu Phe Ser Ala Leu	495
Thr Ala Thr Ala	500	Leu Val Ala Gln Ser	505	Leu Gly Leu Leu Ile	510
Ala Ala Ser Asn	515	Ser Leu Gln Val Ala	520	Thr Phe Val Gly Pro	525
Thr Ala Ile Pro	530	Val Leu Leu Phe Ser	535	Gly Phe Phe Val Ser	540
Lys Thr Ile Pro	545	Thr Tyr Leu Gln Trp	550	Ser Ser Tyr Leu Ser	555
Val Arg Tyr Gly	560	Phe Glu Gly Val Ile	565	Leu Thr Ile Tyr Gly	570
Glu Arg Gly Asp	575	Leu Thr Cys Leu Glu	580	Glu Arg Cys Pro Phe	585
Glu Pro Gln Ser	590	Ile Leu Arg Ala Leu	595	Asp Val Glu Asp Ala	600
Leu Tyr Met Asp	605	Phe Leu Val Leu Gly	610	Ile Phe Phe Leu Ala	615
Arg Leu Leu Ala	620	Tyr Leu Val Leu Arg	625	Arg Val Lys Ser Glu	630
Arg	635		640		645

<210> 4
 <211> 1190
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7473053CD1

<400> 4
 Met Ala Val Cys Ala Lys Lys Arg Pro Pro Glu Glu Glu Arg Arg
 1 5 10 15
 Ala Arg Ala Asn Asp Arg Glu Tyr Asn Glu Lys Phe Gln Tyr Ala
 20 25 30
 Ser Asn Cys Ile Lys Thr Ser Lys Tyr Asn Ile Leu Thr Phe Leu
 35 40 45
 Pro Val Asn Leu Phe Glu Gln Phe Gln Glu Val Ala Asn Thr Tyr
 50 55 60
 Phe Leu Phe Leu Leu Ile Leu Gln Leu Ile Pro Gln Ile Ser Ser

Leu	Ser	Trp	Phe	65	Thr	Ile	Val	Pro	70	Leu	Val	Leu	Val	Leu	Thr	75
				80					85							90
Ile	Thr	Ala	Val	Lys	Asp	Ala	Thr	Asp	100	Asp	Tyr	Phe	Arg	His	Lys	105
Ser	Asp	Asn	Gln	Val	Asn	Asn	Arg	Gln	115	Ser	Gln	Val	Leu	Ile	Asn	120
Gly	Ile	Leu	Gln	Gln	Glu	Gln	Trp	Met	130	Asn	Val	Cys	Val	Gly	Asp	135
Ile	Ile	Lys	Leu	Glu	Asn	Asn	Gln	Phe	145	Val	Ala	Ala	Asp	Leu	Leu	150
Leu	Leu	Ser	Ser	Ser	Glu	Pro	His	Gly	160	Leu	Cys	Tyr	Ile	Glu	Thr	165
Ala	Glu	Leu	Asp	Gly	Glu	Thr	Asn	Met	175	Lys	Val	Arg	Gln	Ala	Ile	180
Pro	Val	Thr	Ser	Glu	Leu	Gly	Asp	Ile	190	Ser	Lys	Leu	Ala	Lys	Phe	195
Asp	Gly	Glu	Val	Ile	Cys	Glu	Pro	Pro	205	Asn	Asn	Lys	Leu	Asp	Lys	210
Phe	Ser	Gly	Thr	Leu	Tyr	Trp	Lys	Glu	220	Asn	Lys	Phe	Pro	Leu	Ser	225
Asn	Gln	Asn	Met	Leu	Leu	Arg	Gly	Cys	235	Val	Leu	Arg	Asn	Thr	Glu	240
Trp	Cys	Phe	Gly	Leu	Val	Ile	Phe	Ala	250	Gly	Pro	Asp	Thr	Lys	Leu	255
Met	Gln	Asn	Ser	Gly	Arg	Thr	Lys	Phe	265	Lys	Arg	Thr	Ser	Ile	Asp	270
Arg	Leu	Met	Asn	Thr	Leu	Val	Leu	Trp	280	Ile	Phe	Gly	Phe	Leu	Val	285
Cys	Met	Gly	Val	Ile	Leu	Ala	Ile	Gly	295	Asn	Ala	Ile	Trp	Glu	His	300
Glu	Val	Gly	Met	Arg	Phe	Gln	Val	Tyr	310	Leu	Pro	Trp	Asp	Glu	Ala	315
Val	Asp	Ser	Ala	Phe	Phe	Ser	Gly	Phe	325	Leu	Ser	Phe	Trp	Ser	Tyr	330
Ile	Ile	Ile	Leu	Asn	Thr	Val	Val	Pro	340	Ile	Ser	Leu	Tyr	Val	Ser	345
Val	Glu	Val	Ile	Arg	Leu	Gly	His	Ser	355	Tyr	Phe	Ile	Asn	Trp	Asp	360
Lys	Lys	Met	Phe	Cys	Met	Lys	Lys	Arg	370	Thr	Pro	Ala	Glu	Ala	Arg	375
Thr	Thr	Thr	Leu	Asn	Glu	Glu	Leu	Gly	385	Gln	Val	Glu	Tyr	Ile	Phe	390
Ser	Asp	Lys	Thr	Gly	Thr	Leu	Thr	Gln	400	Asn	Ile	Met	Val	Phe	Asn	405
Lys	Cys	Ser	Ile	Asn	Gly	His	Ser	Tyr	415	Gly	Asp	Val	Phe	Asp	Val	420
Leu	Gly	His	Lys	Ala	Glu	Leu	Gly	Glu	430	Arg	Pro	Glu	Pro	Val	Asp	435
Phe	Ser	Phe	Asn	Pro	Leu	Ala	Asp	Lys	445	Lys	Phe	Leu	Phe	Trp	Asp	450
Pro	Ser	Leu	Leu	Glu	Ala	Val	Lys	Ile	460	Gly	Asp	Pro	His	Thr	His	465
Glu	Phe	Phe	Arg	Leu	Leu	Ser	Leu	Cys	475	His	Thr	Val	Met	Ser	Glu	480
Glu	Lys	Asn	Glu	Gly	Glu	Leu	Tyr	Tyr	490	Lys	Ala	Gln	Ser	Pro	Asp	495
Glu	Gly	Ala	Leu	Val	Thr	Ala	Ala	Arg	505	Asn	Phe	Gly	Phe	Val	Phe	510
Arg	Ser	Arg	Thr	Pro	Lys	Thr	Ile	Thr	520	Val	His	Glu	Met	Gly	Thr	525
Ala	Ile	Thr	Tyr	Gln	Leu	Leu	Ala	Ile	535	Leu	Asp	Phe	Asn	Asn	Ile	540
Arg	Lys	Arg	Met	Ser	Val	Ile	Val	Arg	550	Asn	Pro	Glu	Gly	Lys	Ile	555
Arg	Leu	Tyr	Cys	Lys	Gly	Ala	Asp	Thr	565	Ile	Leu	Leu	Asp	Arg	Leu	570

His	His	Ser	Thr	Gln	Glu	Leu	Leu	Asn	Thr	Thr	Met	Asp	His	Leu
				575					580					585
Asn	Glu	Tyr	Ala	Gly	Glu	Gly	Leu	Arg	Thr	Leu	Val	Leu	Ala	Tyr
				590					595					600
Lys	Asp	Leu	Asp	Glu	Glu	Tyr	Tyr	Glu	Glu	Trp	Ala	Glu	Arg	Arg
				605					610					615
Leu	Gln	Ala	Ser	Leu	Ala	Gln	Asp	Ser	Arg	Glu	Asp	Arg	Leu	Ala
				620					625					630
Ser	Ile	Tyr	Glu	Glu	Val	Glu	Asn	Asn	Met	Met	Leu	Leu	Gly	Ala
				635					640					645
Thr	Ala	Ile	Glu	Asp	Lys	Leu	Gln	Gln	Gly	Val	Pro	Glu	Thr	Ile
				650					655					660
Ala	Leu	Leu	Thr	Leu	Ala	Asn	Ile	Lys	Ile	Trp	Val	Leu	Thr	Gly
				665					670					675
Asp	Lys	Gln	Glu	Thr	Ala	Val	Asn	Ile	Gly	Tyr	Ser	Cys	Lys	Met
				680					685					690
Leu	Thr	Asp	Asp	Met	Thr	Glu	Val	Phe	Ile	Val	Thr	Gly	His	Thr
				695					700					705
Val	Leu	Glu	Val	Arg	Glu	Glu	Leu	Arg	Lys	Ala	Arg	Glu	Lys	Met
				710					715					720
Met	Asp	Ser	Ser	Arg	Ser	Val	Gly	Asn	Gly	Phe	Thr	Tyr	Gln	Asp
				725					730					735
Lys	Leu	Ser	Ser	Ser	Lys	Leu	Thr	Ser	Val	Leu	Glu	Ala	Val	Ala
				740					745					750
Gly	Glu	Tyr	Ala	Leu	Val	Ile	Asn	Gly	His	Ser	Leu	Ala	His	Ala
				755					760					765
Leu	Glu	Ala	Asp	Met	Glu	Leu	Glu	Phe	Leu	Glu	Thr	Ala	Cys	Ala
				770					775					780
Cys	Lys	Ala	Val	Ile	Cys	Cys	Arg	Val	Thr	Pro	Leu	Gln	Lys	Ala
				785					790					795
Gln	Val	Val	Glu	Leu	Val	Lys	Lys	Tyr	Lys	Lys	Ala	Val	Thr	Leu
				800					805					810
Ala	Ile	Gly	Asp	Gly	Ala	Asn	Asp	Val	Ser	Met	Ile	Lys	Thr	Ala
				815					820					825
His	Ile	Gly	Val	Gly	Ile	Ser	Gly	Gln	Glu	Gly	Ile	Gln	Ala	Val
				830					835					840
Leu	Ala	Ser	Asp	Tyr	Ser	Phe	Ser	Gln	Phe	Lys	Phe	Leu	Gln	Arg
				845					850					855
Leu	Leu	Leu	Val	His	Gly	Arg	Trp	Ser	Tyr	Leu	Arg	Met	Cys	Lys
				860					865					870
Phe	Leu	Cys	Tyr	Phe	Phe	Tyr	Lys	Asn	Phe	Ala	Phe	Thr	Met	Val
				875					880					885
His	Phe	Trp	Phe	Gly	Phe	Phe	Cys	Gly	Phe	Ser	Ala	Gln	Thr	Val
				890					895					900
Tyr	Asp	Gln	Tyr	Phe	Ile	Thr	Leu	Tyr	Asn	Ile	Val	Tyr	Thr	Ser
				905					910					915
Leu	Pro	Val	Leu	Ala	Met	Gly	Val	Phe	Asp	Gln	Asp	Val	Pro	Glu
				920					925					930
Gln	Arg	Ser	Met	Glu	Tyr	Pro	Lys	Leu	Tyr	Glu	Pro	Gly	Gln	Leu
				935					940					945
Asn	Leu	Leu	Phe	Asn	Lys	Arg	Glu	Phe	Phe	Ile	Cys	Ile	Ala	Gln
				950					955					960
Gly	Ile	Tyr	Thr	Ser	Val	Leu	Met	Phe	Phe	Ile	Pro	Tyr	Gly	Val
				965					970					975
Phe	Ala	Asp	Ala	Thr	Arg	Asp	Asp	Gly	Thr	Gln	Leu	Ala	Asp	Tyr
				980					985					990
Gln	Ser	Phe	Ala	Val	Thr	Val	Ala	Thr	Ser	Leu	Val	Ile	Val	Val
				995					1000					1005
Ser	Val	Gln	Ile	Gly	Leu	Asp	Thr	Gly	Tyr	Trp	Thr	Ala	Ile	Asn
				1010					1015					1020
His	Phe	Phe	Ile	Trp	Gly	Ser	Leu	Ala	Val	Tyr	Phe	Ala	Ile	Leu
				1025					1030					1035
Phe	Ala	Met	His	Ser	Asn	Gly	Leu	Phe	Asp	Met	Phe	Pro	Asn	Gln
				1040					1045					1050
Phe	Arg	Phe	Val	Gly	Asn	Ala	Gln	Asn	Thr	Leu	Ala	Gln	Pro	Thr
				1055					1060					1065
Val	Trp	Leu	Thr	Ile	Val	Leu	Thr	Thr	Val	Val	Cys	Ile	Met	Pro

Val	Val	Ala	Phe	Arg	Phe	Leu	Arg	Leu	Asn	Leu	Lys	Pro	Asp	Leu	1070	1075	1080
Ser	Asp	Thr	Val	Arg	Tyr	Thr	Gln	Leu	Val	Arg	Lys	Lys	Gln	Lys	1085	1090	1095
Ala	Gln	His	Arg	Cys	Met	Arg	Arg	Val	Gly	Arg	Thr	Gly	Ser	Arg	1100	1105	1110
Arg	Ser	Gly	Tyr	Ala	Phe	Ser	His	Gln	Glu	Gly	Phe	Gly	Glu	Leu	1115	1120	1125
Ile	Met	Ser	Gly	Lys	Asn	Met	Arg	Leu	Ser	Ser	Leu	Ala	Leu	Ser	1130	1135	1140
Ser	Phe	Thr	Thr	Arg	Ser	Ser	Ser	Ser	Trp	Ile	Glu	Ser	Leu	Arg	1145	1150	1155
Arg	Lys	Lys	Ser	Asp	Ser	Ala	Ser	Ser	Pro	Ser	Gly	Gly	Ala	Asp	1160	1165	1170
Lys	Pro	Leu	Lys	Gly											1175	1180	1185
															1190		

<210> 5

<211> 467

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473347CD1

<400> 5

Met	Val	Leu	Ala	Phe	Gln	Leu	Val	Ser	Phe	Thr	Tyr	Ile	Trp	Ile	1	5	10	15
Ile	Leu	Lys	Pro	Asn	Val	Cys	Ala	Ala	Ser	Asn	Ile	Lys	Met	Thr	20	25	30	35
His	Gln	Arg	Cys	Ser	Ser	Ser	Met	Lys	Gln	Thr	Cys	Lys	Gln	Glu	40	45	50	55
Thr	Arg	Met	Lys	Lys	Asp	Asp	Ser	Thr	Lys	Ala	Arg	Pro	Gln	Lys	60	65	70	75
Tyr	Glu	Gln	Leu	Leu	His	Ile	Glu	Asp	Asn	Asp	Phe	Ala	Met	Arg	80	85	90	95
Pro	Gly	Phe	Gly	Gly	Ser	Pro	Val	Pro	Val	Gly	Ile	Asp	Val	His	100	105	110	115
Val	Glu	Ser	Ile	Asp	Ser	Ile	Ser	Glu	Thr	Asn	Met	Asp	Phe	Thr	120	125	130	135
Met	Thr	Phe	Tyr	Leu	Arg	His	Tyr	Trp	Lys	Asp	Glu	Arg	Leu	Ser	140	145	150	155
Phe	Pro	Ser	Thr	Ala	Asn	Lys	Ser	Met	Thr	Phe	Asp	His	Arg	Leu	160	165	170	175
Thr	Arg	Lys	Ile	Trp	Val	Pro	Asp	Ile	Phe	Phe	Val	His	Ser	Lys	180	185	190	195
Arg	Ser	Phe	Ile	His	Asp	Thr	Thr	Met	Glu	Asn	Ile	Met	Leu	Arg	200	205	210	215
Val	His	Pro	Asp	Gly	Asn	Val	Leu	Leu	Ser	Leu	Arg	Ile	Thr	Val	220	225	230	235
Ser	Ala	Met	Cys	Phe	Met	Asp	Phe	Ser	Arg	Phe	Pro	Leu	Asp	Thr	240	245	250	255
Gln	Asn	Cys	Ser	Leu	Glu	Leu	Glu	Ser	Tyr	Ala	Tyr	Asn	Glu	Asp	260	265	270	275
Asp	Leu	Met	Leu	Tyr	Trp	Lys	His	Gly	Asn	Lys	Ser	Leu	Asn	Thr	280	285	290	295
Glu	Glu	His	Met	Ser	Leu	Ser	Gln	Phe	Phe	Ile	Glu	Asp	Phe	Ser	300	305	310	315
Ala	Ser	Ser	Gly	Leu	Ala	Phe	Tyr	Ser	Ser	Thr	Gly	Trp	Tyr	Asn	320	325	330	335
Arg	Leu	Phe	Ile	Ile	Ser	Val	Leu	Arg	Arg	His	Val	Phe	Phe	Phe	340	345	350	355
Val	Leu	Pro	Thr	Tyr	Tyr	Pro	Ala	Ile	Leu	Met	Val	Met	Leu	Ser	360	365	370	375
Trp	Val	Ser	Phe	Trp	Ile	Asp	Arg	Arg	Ala	Val	Pro	Ala	Arg	Val	380	385	390	395

Ser	Leu	Gly	Ile	290	Thr	Val	Leu	Thr	295	Met	Ser	Thr	Ile	Ile	300
				305					310						315
Ala	Val	Ser	Ala	320	Ser	Met	Pro	Gln	325	Ser	Tyr	Leu	Lys	Ala	330
Asp	Val	Tyr	Leu	335	Trp	Val	Ser	Ser	340	Phe	Val	Phe	Leu	Ser	345
Ile	Glu	Tyr	Ala	350	Ala	Val	Asn	Tyr	355	Thr	Thr	Val	Glu	Glu	360
Lys	Gln	Phe	Lys	365	Lys	Thr	Gly	Lys	370	Ser	Arg	Met	Tyr	Asn	375
Asp	Ala	Val	Gln	380	Ala	Met	Ala	Phe	385	Gly	Cys	Tyr	His	Asp	390
Glu	Ile	Asp	Met	395	Asp	Gln	Thr	Ser	400	Ser	Leu	Asn	Ser	Glu	405
Phe	Met	Arg	Arg	410	Lys	Ser	Ile	Cys	415	Pro	Ser	Thr	Asp	Ser	420
Arg	Ile	Lys	Arg	425	Arg	Lys	Ser	Leu	430	Gly	Gly	His	Val	Gly	435
Ile	Leu	Glu	Asn	440	Asn	His	Val	Ile	445	Thr	Tyr	Ser	Arg	Ile	450
Phe	Pro	Ile	Val	455	Tyr	Ile	Leu	Phe	460	Leu	Phe	Tyr	Trp	Gly	465
Tyr	Val														

<210> 6

<211> 1196

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474240CD1

<400> 6

Met	Pro	Val	Arg	Arg	Gly	His	Val	Ala	Pro	Gln	Asn	Thr	Phe	Leu	
1				5					10					15	
Gly	Thr	Ile	Ile	Arg	Lys	Phe	Glu	Gly	Gln	Asn	Lys	Lys	Phe	Ile	
				20					25					30	
Ile	Ala	Asn	Ala	Arg	Val	Gln	Asn	Cys	Ala	Ile	Ile	Tyr	Cys	Asn	
				35					40					45	
Asp	Gly	Phe	Cys	Glu	Met	Thr	Gly	Phe	Ser	Arg	Pro	Asp	Val	Met	
				50					55					60	
Gln	Lys	Pro	Cys	Thr	Cys	Asp	Phe	Leu	His	Gly	Pro	Glu	Thr	Lys	
				65					70					75	
Arg	His	Asp	Ile	Ala	Gln	Ile	Ala	Gln	Ala	Leu	Leu	Gly	Ser	Glu	
				80					85					90	
Glu	Arg	Lys	Val	Glu	Val	Thr	Tyr	Tyr	His	Lys	Asn	Gly	Ser	Thr	
				95					100					105	
Phe	Ile	Cys	Asn	Thr	His	Ile	Ile	Pro	Val	Lys	Asn	Gln	Glu	Gly	
				110					115					120	
Val	Ala	Met	Met	Phe	Ile	Ile	Asn	Phe	Glu	Tyr	Val	Thr	Asp	Asn	
				125					130					135	
Glu	Asn	Ala	Ala	Thr	Pro	Glu	Arg	Val	Asn	Pro	Ile	Leu	Pro	Ile	
				140					145					150	
Lys	Thr	Val	Asn	Arg	Lys	Phe	Phe	Gly	Phe	Lys	Phe	Pro	Gly	Leu	
				155					160					165	
Arg	Val	Leu	Thr	Tyr	Arg	Lys	Gln	Ser	Leu	Pro	Gln	Glu	Asp	Pro	
				170					175					180	
Asp	Val	Val	Val	Ile	Asp	Ser	Ser	Lys	His	Ser	Asp	Asp	Ser	Val	
				185					190					195	
Ala	Met	Lys	His	Phe	Lys	Ser	Pro	Thr	Lys	Glu	Ser	Cys	Ser	Pro	
				200					205					210	
Ser	Glu	Ala	Asp	Asp	Thr	Lys	Ala	Leu	Ile	Gln	Pro	Ser	Lys	Cys	
				215					220					225	
Ser	Pro	Leu	Val	Asn	Ile	Ser	Gly	Pro	Leu	Asp	His	Ser	Ser	Pro	

Lys Arg Gln Trp	230	Arg Leu Tyr Pro	235	Asp Met Leu Gln Ser	240
Ser Gln Leu Ser	245	His Ser Arg Ser Arg	250	Glu Ser Leu Cys Ser	255
Arg Arg Ala Ser	260	Ser Val His Asp Ile	265	Glu Gly Phe Gly Val	270
Pro Lys Asn Ile	275	Phe Arg Asp Arg His	280	Ala Ser Glu Asp Asn	285
Arg Asn Val Lys	290	Gly Pro Phe Asn His	295	Ile Lys Ser Ser Leu	300
Gly Ser Thr Ser	305	Asp Ser Asn Leu Asn	310	Lys Tyr Ser Thr Ile	315
Lys Ile Pro Gln	320	Leu Thr Leu Asn Phe	325	Ser Glu Val Lys Thr	330
Lys Lys Asn Ser	335	Ser Pro Pro Ser Ser	340	Asp Lys Thr Ile Ile	345
Pro Lys Val Lys	350	Asp Arg Thr His Asn	355	Val Thr Glu Lys Val	360
Gln Val Leu Ser	365	Leu Gly Ala Asp Val	370	Leu Pro Glu Tyr Lys	375
Gln Thr Pro Arg	380	Ile Asn Lys Phe Thr	385	Leu Leu His Tyr Ser	390
Phe Lys Ala Val	395	Trp Asp Trp Leu Ile	400	Leu Leu Leu Val Ile	405
Thr Ala Ile Phe	410	Thr Pro Tyr Ser Ala	415	Ala Phe Leu Leu Asn	420
Arg Glu Glu Gln	425	Lys Arg Arg Glu Cys	430	Gly Tyr Ser Cys Ser	435
Leu Asn Val Val	440	Asp Leu Ile Val Asp	445	Ile Met Phe Ile Ile	450
Ile Leu Ile Asn	455	Phe Arg Thr Thr Tyr	460	Val Asn Gln Asn Glu	465
Val Val Ser Asp	470	Pro Ala Lys Ile Ala	475	Ile His Tyr Phe Lys	480
Trp Phe Leu Ile	485	Asp Met Val Ala Ala	490	Ile Pro Phe Asp Leu	495
Ile Phe Gly Ser	500	Gly Ser Asp Glu Thr	505	Thr Thr Leu Ile Gly	510
Leu Lys Thr Ala	515	Arg Leu Leu Arg Leu	520	Val Arg Val Ala Arg	525
Leu Asp Arg Tyr	530	Ser Glu Tyr Gly Ala	535	Ala Val Leu Met Leu	540
Met Cys Ile Phe	545	Ala Leu Ile Ala His	550	Trp Leu Ala Cys Ile	555
Tyr Ala Ile Gly	560	Asn Val Glu Arg Pro	565	Tyr Leu Thr Asp Lys	570
Gly Trp Leu Asp	575	Ser Leu Gly Gln Gln	580	Ile Gly Lys Arg Tyr	585
Asp Ser Asp Ser	590	Ser Ser Gly Pro Ser	595	Ile Lys Asp Lys Tyr	600
Thr Ala Leu Tyr	605	Phe Thr Phe Ser Ser	610	Leu Thr Ser Val Gly	615
Gly Asn Val Ser	620	Pro Asn Thr Asn Ser	625	Glu Lys Ile Phe Ser	630
Cys Val Met Leu	635	Ile Gly Ser Leu Met	640	Tyr Ala Ser Ile Phe	645
Asn Val Ser Ala	650	Ile Ile Gln Arg Leu	655	Tyr Ser Gly Thr Ala	660
Tyr His Met Gln	665	Met Leu Arg Val Lys	670	Glu Phe Ile Arg Phe	675
Gln Ile Pro Asn	680	Pro Leu Arg Gln Arg	685	Leu Glu Glu Tyr Phe	690
His Ala Trp Thr	695	Tyr Thr Asn Gly Ile	700	Asp Met Asn Met Val	705
Lys Gly Phe Pro	710	Glu Cys Leu Gln Ala	715	Asp Ile Cys Leu His	720
	725		730		735

Asn	Gln	Thr	Leu	Leu	Gln	Asn	Cys	Lys	Ala	Phe	Arg	Gly	Ala	Ser
				740					745					750
Lys	Gly	Cys	Leu	Arg	Ala	Leu	Ala	Met	Lys	Phe	Lys	Thr	Thr	His
				755					760					765
Ala	Pro	Pro	Gly	Asp	Thr	Leu	Val	His	Cys	Gly	Asp	Val	Leu	Thr
				770					775					780
Ala	Leu	Tyr	Phe	Leu	Ser	Arg	Gly	Ser	Ile	Glu	Ile	Leu	Lys	Asp
				785					790					795
Asp	Ile	Val	Val	Ala	Ile	Leu	Gly	Lys	Asn	Asp	Ile	Phe	Gly	Glu
				800					805					810
Met	Val	His	Leu	Tyr	Ala	Lys	Pro	Gly	Lys	Ser	Asn	Ala	Asp	Val
				815					820					825
Arg	Ala	Leu	Thr	Tyr	Cys	Asp	Leu	His	Lys	Ile	Gln	Arg	Glu	Asp
				830					835					840
Leu	Leu	Glu	Val	Leu	Asp	Met	Tyr	Pro	Glu	Phe	Ser	Asp	His	Phe
				845					850					855
Leu	Thr	Asn	Leu	Glu	Leu	Thr	Phe	Asn	Leu	Arg	His	Glu	Ser	Ala
				860					865					870
Lys	Ala	Asp	Leu	Leu	Arg	Ser	Gln	Ser	Met	Asn	Asp	Ser	Glu	Gly
				875					880					885
Asp	Asn	Cys	Lys	Leu	Arg	Arg	Arg	Lys	Leu	Ser	Phe	Glu	Ser	Glu
				890					895					900
Gly	Glu	Lys	Glu	Asn	Ser	Thr	Asn	Asp	Pro	Glu	Asp	Ser	Ala	Asp
				905					910					915
Thr	Ile	Arg	His	Tyr	Gln	Ser	Ser	Lys	Arg	His	Phe	Glu	Glu	Lys
				920					925					930
Lys	Ser	Arg	Ser	Ser	Ser	Phe	Ile	Ser	Ser	Ile	Asp	Asp	Glu	Gln
				935					940					945
Lys	Pro	Leu	Phe	Ser	Gly	Ile	Val	Asp	Ser	Ser	Pro	Gly	Ile	Gly
				950					955					960
Lys	Ala	Ser	Gly	Leu	Asp	Phe	Glu	Glu	Thr	Val	Pro	Thr	Ser	Gly
				965					970					975
Arg	Met	His	Ile	Asp	Lys	Arg	Ser	His	Ser	Cys	Lys	Asp	Ile	Thr
				980					985					990
Asp	Met	Arg	Ser	Trp	Glu	Arg	Glu	Asn	Ala	His	Pro	Gln	Pro	Glu
				995					1000					1005
Asp	Ser	Ser	Pro	Ser	Ala	Leu	Gln	Arg	Ala	Ala	Trp	Gly	Ile	Ser
				1010					1015					1020
Glu	Thr	Glu	Ser	Asp	Leu	Thr	Tyr	Gly	Glu	Val	Glu	Gln	Arg	Leu
				1025					1030					1035
Asp	Leu	Leu	Gln	Glu	Gln	Leu	Asn	Arg	Leu	Glu	Ser	Gln	Met	Thr
				1040					1045					1050
Thr	Asp	Ile	Gln	Thr	Ile	Leu	Gln	Leu	Leu	Gln	Lys	Gln	Thr	Thr
				1055					1060					1065
Val	Val	Pro	Pro	Ala	Tyr	Ser	Met	Val	Thr	Ala	Gly	Ser	Glu	Tyr
				1070					1075					1080
Gln	Arg	Pro	Ile	Ile	Gln	Leu	Met	Arg	Thr	Ser	Gln	Pro	Glu	Ala
				1085					1090					1095
Ser	Ile	Lys	Thr	Asp	Arg	Ser	Phe	Ser	Pro	Ser	Ser	Gln	Cys	Pro
				1100					1105					1110
Glu	Phe	Leu	Asp	Leu	Glu	Lys	Ser	Lys	Leu	Lys	Ser	Lys	Glu	Ser
				1115					1120					1125
Leu	Ser	Ser	Gly	Val	His	Leu	Asn	Thr	Ala	Ser	Glu	Asp	Asn	Leu
				1130					1135					1140
Thr	Ser	Leu	Leu	Lys	Gln	Asp	Ser	Asp	Leu	Ser	Leu	Glu	Leu	His
				1145					1150					1155
Leu	Arg	Gln	Arg	Lys	Thr	Tyr	Val	His	Pro	Ile	Arg	His	Pro	Ser
				1160					1165					1170
Leu	Pro	Asp	Ser	Ser	Leu	Ser	Thr	Val	Gly	Ile	Val	Gly	Leu	His
				1175					1180					1185
Arg	His	Val	Ser	Asp	Pro	Gly	Leu	Pro	Gly	Lys				
				1190					1195					

<210> 7

<211> 512

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475338CD1

<400> 7

Met	Glu	Asn	Lys	Glu	Ala	Gly	Thr	Pro	Pro	Pro	Ile	Pro	Ser	Arg	15
1				5					10						
Glu	Gly	Arg	Leu	Gln	Pro	Thr	Leu	Leu	Leu	Ala	Thr	Leu	Ser	Ala	30
				20					25						
Ala	Phe	Gly	Ser	Ala	Phe	Gln	Tyr	Gly	Tyr	Asn	Leu	Ser	Val	Val	45
				35					40						
Asn	Thr	Pro	His	Lys	Val	Phe	Lys	Ser	Phe	Tyr	Asn	Glu	Thr	Tyr	60
				50					55						
Phe	Glu	Arg	His	Ala	Thr	Phe	Met	Asp	Gly	Lys	Leu	Met	Leu	Leu	75
				65					70						
Leu	Trp	Ser	Cys	Thr	Val	Ser	Met	Phe	Pro	Leu	Gly	Gly	Leu	Leu	90
				80					85						
Gly	Ser	Leu	Leu	Val	Gly	Leu	Leu	Val	Asp	Ser	Cys	Gly	Arg	Lys	105
				95					100						
Gly	Thr	Leu	Leu	Ile	Asn	Asn	Ile	Phe	Ala	Ile	Ile	Pro	Ala	Ile	120
				110					115						
Leu	Met	Gly	Val	Ser	Lys	Val	Ala	Lys	Ala	Phe	Glu	Leu	Ile	Val	135
				125					130						
Phe	Ser	Arg	Val	Val	Leu	Gly	Val	Cys	Ala	Gly	Ile	Ser	Tyr	Ser	150
				140					145						
Ala	Leu	Pro	Met	Tyr	Leu	Gly	Glu	Leu	Ala	Pro	Lys	Asn	Leu	Arg	165
				155					160						
Gly	Met	Val	Gly	Thr	Met	Thr	Glu	Val	Phe	Val	Ile	Val	Gly	Val	180
				170					175						
Phe	Leu	Ala	Gln	Ile	Phe	Ser	Leu	Gln	Ala	Ile	Leu	Gly	Asn	Pro	195
				185					190						
Ala	Gly	Trp	Pro	Val	Leu	Leu	Ala	Leu	Thr	Gly	Val	Pro	Ala	Leu	210
				200					205						
Leu	Gln	Leu	Leu	Thr	Leu	Pro	Phe	Phe	Pro	Glu	Ser	Pro	Arg	Tyr	225
				215					220						
Ser	Leu	Ile	Gln	Lys	Gly	Asp	Glu	Ala	Thr	Ala	Arg	Gln	Ala	Leu	240
				230					235						
Arg	Arg	Leu	Arg	Gly	His	Thr	Asp	Met	Glu	Ala	Glu	Leu	Glu	Asp	255
				245					250						
Met	Arg	Ala	Glu	Ala	Arg	Ala	Glu	Arg	Ala	Glu	Gly	His	Leu	Ser	270
				260					265						
Val	Leu	His	Leu	Cys	Ala	Leu	Arg	Ser	Leu	Arg	Trp	Gln	Leu	Leu	285
				275					280						
Ser	Ile	Ile	Val	Leu	Met	Ala	Gly	Gln	Gln	Leu	Ser	Gly	Ile	Asn	300
				290					295						
Ala	Ile	Asn	Tyr	Tyr	Ala	Asp	Thr	Ile	Tyr	Thr	Ser	Ala	Gly	Val	315
				305					310						
Glu	Ala	Ala	His	Ser	Gln	Tyr	Val	Thr	Val	Gly	Ser	Gly	Val	Val	330
				320					325						
Asn	Ile	Val	Met	Thr	Ile	Thr	Ser	Ala	Val	Leu	Val	Glu	Arg	Leu	345
				335					340						
Gly	Arg	Arg	His	Leu	Leu	Leu	Ala	Gly	Tyr	Gly	Ile	Cys	Gly	Ser	360
				350					355						
Ala	Cys	Leu	Val	Leu	Thr	Val	Val	Leu	Leu	Phe	Gln	Asn	Arg	Val	375
				365					370						
Pro	Glu	Leu	Ser	Tyr	Leu	Gly	Ile	Ile	Cys	Val	Phe	Ala	Tyr	Ile	390
				380					385						
Ala	Gly	His	Ser	Ile	Gly	Pro	Ser	Pro	Val	Pro	Ser	Val	Val	Arg	405
				395					400						
Thr	Glu	Ile	Phe	Leu	Gln	Ser	Ser	Arg	Arg	Ala	Ala	Phe	Met	Val	420
				410					415						
Asp	Gly	Ala	Val	His	Trp	Leu	Thr	Asn	Phe	Ile	Ile	Gly	Phe	Leu	435
				425					430						
Phe	Pro	Ser	Ile	Gln	Glu	Ala	Ile	Gly	Ala	Tyr	Ser	Phe	Ile	Ile	450
				440					445						
Phe	Ala	Gly	Ile	Cys	Leu	Leu	Thr	Ala	Ile	Tyr	Ile	Tyr	Val	Val	465
				455					460						

Ile	Pro	Glu	Thr	Lys	Gly	Lys	Thr	Phe	Val	Glu	Ile	Asn	Arg	Ile
				470					475					480
Phe	Ala	Lys	Arg	Asn	Arg	Val	Lys	Leu	Pro	Glu	Glu	Lys	Glu	Glu
				485					490					495
Thr	Ile	Asp	Ala	Gly	Pro	Pro	Thr	Ala	Ser	Pro	Ala	Lys	Glu	Thr
				500					505					510
Ser	Phe													

<210> 8

<211> 568

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7476747CD1

<400> 8

Met	Thr	Ala	Ser	Thr	Pro	Glu	Ala	Thr	Pro	Asn	Met	Glu	Leu	Lys
1				5					10					15
Ala	Pro	Ala	Ala	Gly	Gly	Leu	Asn	Ala	Gly	Pro	Val	Pro	Pro	Ala
				20					25					30
Ala	Met	Ser	Thr	Gln	Arg	Leu	Arg	Asn	Glu	Asp	Tyr	His	Asp	Tyr
				35					40					45
Ser	Ser	Thr	Asp	Val	Ser	Pro	Glu	Glu	Ser	Pro	Ser	Glu	Gly	Leu
				50					55					60
Asn	Asn	Leu	Ser	Ser	Pro	Gly	Ser	Tyr	Gln	Arg	Phe	Gly	Gln	Ser
				65					70					75
Asn	Ser	Thr	Thr	Trp	Phe	Gln	Thr	Leu	Ile	His	Leu	Leu	Lys	Gly
				80					85					90
Asn	Ile	Gly	Thr	Gly	Leu	Leu	Gly	Leu	Pro	Leu	Ala	Val	Lys	Asn
				95					100					105
Ala	Gly	Ile	Val	Met	Gly	Pro	Ile	Ser	Leu	Leu	Ile	Ile	Gly	Ile
				110					115					120
Val	Ala	Val	His	Cys	Met	Gly	Ile	Leu	Val	Lys	Cys	Ala	His	His
				125					130					135
Phe	Cys	Arg	Arg	Leu	Asn	Lys	Ser	Phe	Val	Asp	Tyr	Gly	Asp	Thr
				140					145					150
Val	Met	Tyr	Gly	Leu	Glu	Ser	Ser	Pro	Cys	Ser	Trp	Leu	Arg	Asn
				155					160					165
His	Ala	His	Trp	Gly	Arg	Arg	Val	Val	Asp	Phe	Phe	Leu	Ile	Val
				170					175					180
Thr	Gln	Leu	Gly	Phe	Cys	Cys	Val	Tyr	Phe	Val	Phe	Leu	Ala	Asp
				185					190					195
Asn	Phe	Lys	Gln	Val	Ile	Glu	Ala	Ala	Asn	Gly	Thr	Thr	Asn	Asn
				200					205					210
Cys	His	Asn	Asn	Glu	Thr	Val	Ile	Leu	Thr	Pro	Thr	Met	Asp	Ser
				215					220					225
Arg	Leu	Tyr	Met	Leu	Ser	Phe	Leu	Pro	Phe	Leu	Val	Leu	Leu	Val
				230					235					240
Phe	Ile	Arg	Asn	Leu	Arg	Ala	Leu	Ser	Ile	Phe	Ser	Leu	Leu	Ala
				245					250					255
Asn	Ile	Thr	Met	Leu	Val	Ser	Leu	Val	Met	Ile	Tyr	Gln	Phe	Ile
				260					265					270
Val	Gln	Arg	Ile	Pro	Asp	Pro	Ser	His	Leu	Pro	Leu	Val	Ala	Pro
				275					280					285
Trp	Lys	Thr	Tyr	Pro	Leu	Phe	Phe	Gly	Thr	Ala	Ile	Phe	Ser	Phe
				290					295					300
Glu	Gly	Ile	Gly	Met	Val	Leu	Pro	Leu	Glu	Asn	Lys	Met	Lys	Asp
				305					310					315
Pro	Arg	Lys	Phe	Pro	Leu	Ile	Leu	Tyr	Leu	Gly	Met	Val	Ile	Val
				320					325					330
Thr	Ile	Leu	Tyr	Ile	Ser	Leu	Gly	Cys	Leu	Gly	Tyr	Leu	Gln	Phe
				335					340					345
Gly	Ala	Asn	Ile	Gln	Gly	Ser	Ile	Thr	Leu	Asn	Leu	Pro	Asn	Cys
				350					355					360

Trp	Leu	Tyr	Gln	Ser	Val	Lys	Leu	Leu	Tyr	Ser	Ile	Gly	Ile	Phe
				365					370					375
Phe	Thr	Tyr	Ala	Leu	Gln	Phe	Tyr	Val	Pro	Ala	Glu	Ile	Ile	Ile
				380					385					390
Pro	Phe	Phe	Val	Ser	Arg	Ala	Pro	Glu	Pro	Cys	Glu	Leu	Val	Val
				395					400					405
Asp	Leu	Phe	Val	Arg	Pro	Val	Leu	Val	Cys	Leu	Thr	Ser	Leu	Ser
				410					415					420
Gly	Ser	Val	Asp	Asn	Gly	Trp	Tyr	Gly	Thr	Glu	Ala	Asp	Gly	Thr
				425					430					435
Ser	Cys	Gly	Ser	Ala	Pro	Leu	Val	Phe	Val	Ser	Ser	Ser	Phe	Leu
				440					445					450
Ala	His	Pro	Trp	Leu	Ser	Phe	Arg	Cys	Glu	Ser	Gln	Trp	Val	Ser
				455					460					465
Cys	His	Arg	Asp	Thr	Val	Val	Val	Trp	Gly	Phe	Ala	Arg	Gly	Ile
				470					475					480
Leu	Ala	Ile	Leu	Ile	Pro	Arg	Leu	Asp	Leu	Val	Ile	Ser	Leu	Val
				485					490					495
Gly	Ser	Val	Ser	Ser	Ser	Ala	Leu	Ala	Leu	Ile	Ile	Pro	Pro	Leu
				500					505					510
Leu	Glu	Val	Thr	Thr	Phe	Tyr	Ser	Glu	Gly	Met	Ser	Pro	Leu	Thr
				515					520					525
Ile	Phe	Lys	Asp	Ala	Leu	Ile	Ser	Ile	Leu	Gly	Phe	Val	Gly	Phe
				530					535					540
Val	Val	Gly	Thr	Tyr	Glu	Ala	Leu	Tyr	Glu	Leu	Ile	Gln	Pro	Ser
				545					550					555
Asn	Ala	Pro	Ile	Phe	Ile	Asn	Ser	Thr	Cys	Ala	Phe	Ile		
				560					565					

<210> 9

<211> 958

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477898CD1

<400> 9

Met	Pro	Val	Arg	Arg	Gly	His	Val	Ala	Pro	Gln	Asn	Thr	Tyr	Leu
1				5					10					15
Asp	Thr	Ile	Ile	Arg	Lys	Phe	Glu	Gly	Gln	Ser	Arg	Lys	Phe	Leu
				20					25					30
Ile	Ala	Asn	Ala	Gln	Met	Glu	Asn	Cys	Ala	Ile	Ile	Tyr	Cys	Asn
				35					40					45
Asp	Gly	Phe	Cys	Glu	Leu	Phe	Gly	Tyr	Ser	Arg	Val	Glu	Val	Met
				50					55					60
Gln	Gln	Pro	Cys	Thr	Cys	Asp	Phe	Leu	Thr	Gly	Pro	Asn	Thr	Pro
				65					70					75
Ser	Ser	Ala	Val	Ser	Arg	Leu	Ala	Gln	Ala	Leu	Leu	Gly	Ala	Glu
				80					85					90
Glu	Cys	Lys	Val	Asp	Ile	Leu	Tyr	Tyr	Arg	Lys	Asp	Ala	Ser	Ser
				95					100					105
Phe	Arg	Cys	Leu	Val	Asp	Val	Val	Pro	Val	Lys	Asn	Glu	Asp	Gly
				110					115					120
Ala	Val	Ile	Met	Phe	Ile	Leu	Asn	Phe	Glu	Asp	Leu	Ala	Gln	Leu
				125					130					135
Leu	Ala	Lys	Cys	Ser	Ser	Arg	Ser	Leu	Ser	Gln	Arg	Leu	Leu	Ser
				140					145					150
Gln	Ser	Phe	Leu	Gly	Ser	Glu	Gly	Ser	His	Gly	Arg	Pro	Gly	Gly
				155					160					165
Pro	Gly	Pro	Gly	Thr	Gly	Arg	Gly	Lys	Tyr	Arg	Thr	Ile	Ser	Gln
				170					175					180
Ile	Pro	Gln	Phe	Thr	Leu	Asn	Phe	Val	Glu	Phe	Asn	Leu	Glu	Lys
				185					190					195
His	Arg	Ser	Ser	Ser	Thr	Thr	Glu	Ile	Glu	Ile	Ile	Ala	Pro	His
				200					205					210

Lys	Val	Val	Glu	Arg	Thr	Gln	Asn	Val	Thr	Glu	Lys	Val	Thr	Gln
				215					220					225
Val	Leu	Ser	Leu	Gly	Ala	Asp	Val	Leu	Pro	Glu	Tyr	Lys	Leu	Gln
				230					235					240
Ala	Pro	Arg	Ile	His	Arg	Trp	Thr	Ile	Leu	His	Tyr	Ser	Pro	Phe
				245					250					255
Lys	Ala	Val	Trp	Asp	Trp	Leu	Ile	Leu	Leu	Leu	Val	Ile	Tyr	Thr
				260					265					270
Ala	Val	Phe	Thr	Pro	Tyr	Ser	Ala	Ala	Phe	Leu	Leu	Ser	Asp	Gln
				275					280					285
Asp	Glu	Ser	Arg	Arg	Gly	Ala	Cys	Ser	Tyr	Thr	Cys	Ser	Pro	Leu
				290					295					300
Thr	Val	Val	Asp	Leu	Ile	Val	Asp	Ile	Met	Phe	Val	Val	Asp	Ile
				305					310					315
Val	Ile	Asn	Phe	Arg	Thr	Thr	Tyr	Val	Asn	Thr	Asn	Asp	Glu	Val
				320					325					330
Val	Ser	His	Pro	Arg	Arg	Ile	Ala	Val	His	Tyr	Phe	Lys	Gly	Trp
				335					340					345
Phe	Leu	Ile	Asp	Met	Val	Ala	Ala	Ile	Pro	Phe	Asp	Leu	Leu	Ile
				350					355					360
Phe	Arg	Thr	Gly	Ser	Asp	Glu	Thr	Thr	Thr	Leu	Ile	Gly	Leu	Leu
				365					370					375
Lys	Thr	Ala	Arg	Leu	Leu	Arg	Leu	Val	Arg	Val	Ala	Arg	Lys	Leu
				380					385					390
Asp	Arg	Tyr	Ser	Glu	Tyr	Gly	Ala	Ala	Val	Leu	Phe	Leu	Leu	Met
				395					400					405
Cys	Thr	Phe	Pro	Leu	Ile	Ala	His	Trp	Leu	Ala	Cys	Ile	Trp	Tyr
				410					415					420
Ala	Ile	Gly	Asn	Val	Glu	Arg	Pro	Tyr	Leu	Glu	His	Lys	Ile	Gly
				425					430					435
Trp	Leu	Asp	Ser	Leu	Gly	Val	Gln	Leu	Gly	Lys	Arg	Tyr	Asn	Gly
				440					445					450
Ser	Asp	Pro	Ala	Ser	Gly	Pro	Ser	Val	Gln	Asp	Lys	Tyr	Val	Thr
				455					460					465
Ala	Leu	Tyr	Phe	Thr	Phe	Ser	Ser	Leu	Thr	Ser	Val	Gly	Phe	Gly
				470					475					480
Asn	Val	Ser	Pro	Asn	Thr	Asn	Ser	Glu	Lys	Val	Phe	Ser	Ile	Cys
				485					490					495
Val	Met	Leu	Ile	Gly	Ser	Leu	Met	Tyr	Ala	Ser	Ile	Phe	Gly	Asn
				500					505					510
Val	Ser	Ala	Ile	Ile	Gln	Arg	Leu	Tyr	Ser	Gly	Thr	Ala	Arg	Tyr
				515					520					525
His	Thr	Gln	Met	Leu	Arg	Val	Lys	Glu	Phe	Ile	Arg	Phe	His	Gln
				530					535					540
Ile	Pro	Asn	Pro	Leu	Arg	Gln	Arg	Leu	Glu	Glu	Tyr	Phe	Gln	His
				545					550					555
Ala	Trp	Ser	Tyr	Thr	Asn	Gly	Ile	Asp	Met	Asn	Ala	Val	Leu	Lys
				560					565					570
Gly	Phe	Pro	Glu	Cys	Leu	Gln	Ala	Asp	Ile	Cys	Leu	His	Leu	His
				575					580					585
Arg	Ala	Leu	Leu	Gln	His	Cys	Pro	Ala	Phe	Ser	Gly	Ala	Gly	Lys
				590					595					600
Gly	Cys	Leu	Arg	Ala	Leu	Ala	Val	Lys	Phe	Lys	Thr	Thr	His	Ala
				605					610					615
Pro	Pro	Gly	Asp	Thr	Leu	Val	His	Leu	Gly	Asp	Val	Leu	Ser	Thr
				620					625					630
Leu	Tyr	Phe	Ile	Ser	Arg	Gly	Ser	Ile	Glu	Ile	Leu	Arg	Asp	Asp
				635					640					645
Val	Val	Val	Ala	Ile	Leu	Gly	Lys	Asn	Asp	Ile	Phe	Gly	Glu	Pro
				650					655					660
Val	Ser	Leu	His	Ala	Gln	Pro	Gly	Lys	Ser	Ser	Ala	Asp	Val	Arg
				665					670					675
Ala	Leu	Thr	Tyr	Cys	Asp	Leu	His	Lys	Ile	Gln	Arg	Ala	Asp	Leu
				680					685					690
Leu	Glu	Val	Leu	Asp	Met	Tyr	Pro	Ala	Phe	Ala	Glu	Ser	Phe	Trp
				695					700					705
Ser	Lys	Leu	Glu	Val	Thr	Phe	Asn	Leu	Arg	Asp	Val	Thr	Gly	Gly

Leu	His	Ser	Ser	710	Pro	Arg	Gln	Ala	Pro	715	Gly	Ser	Gln	Asp	His	Gln	720
				725	Pro					730							735
Gly	Phe	Phe	Leu	Ser	Asp	Asn	Gln	Ser	Asp	Ala	Ala	Pro	Pro	Pro	Pro	Leu	740
				740						745							750
Ser	Ile	Ser	Asp	Ala	Phe	Trp	Leu	Trp	Pro	Glu	Leu	Leu	Gln	Glu			755
				755						760							765
Met	Pro	Pro	Lys	His	Ser	Pro	Gln	Ser	Pro	Gln	Glu	Asp	Pro	Asp			770
				770						775							780
Cys	Trp	Pro	Leu	Lys	Leu	Gly	Ser	Arg	Leu	Glu	Gln	Leu	Gln	Ala			785
				785						790							795
Gln	Met	Asn	Arg	Leu	Glu	Ser	Arg	Val	Ser	Ser	Asp	Leu	Ser	Arg			800
				800						805							810
Ile	Leu	Gln	Leu	Leu	Gln	Lys	Pro	Met	Pro	Gln	Gly	His	Ala	Ser			815
				815						820							825
Tyr	Ile	Leu	Glu	Ala	Pro	Ala	Ser	Asn	Asp	Leu	Ala	Leu	Val	Pro			830
				830						835							840
Ile	Ala	Ser	Glu	Thr	Thr	Ser	Pro	Gly	Pro	Arg	Leu	Pro	Gln	Gly			845
				845						850							855
Phe	Leu	Pro	Pro	Ala	Gln	Thr	Pro	Ser	Tyr	Gly	Asp	Leu	Asp	Asp			860
				860						865							870
Cys	Ser	Pro	Lys	His	Arg	Asn	Ser	Ser	Pro	Arg	Met	Pro	His	Leu			875
				875						880							885
Ala	Val	Ala	Met	Asp	Lys	Thr	Leu	Ala	Pro	Ser	Ser	Glu	Gln	Glu			890
				890						895							900
Gln	Pro	Glu	Gly	Leu	Trp	Pro	Pro	Leu	Ala	Ser	Pro	Leu	His	Pro			905
				905						910							915
Leu	Glu	Val	Gln	Gly	Leu	Ile	Cys	Gly	Pro	Cys	Phe	Ser	Ser	Leu			920
				920						925							930
Pro	Glu	His	Leu	Gly	Ser	Val	Pro	Lys	Gln	Leu	Asp	Phe	Gln	Arg			935
				935						940							945
His	Gly	Ser	Asp	Pro	Gly	Phe	Ala	Gly	Ser	Trp	Gly	His					950
				950						955							

<210> 10

<211> 724

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472728CD1

<400> 10

Met	Gly	His	Gln	Gly	Pro	Phe	Glu	Glu	Gly	Asn	Gly	Gly	Leu	Arg			
1				5					10					15			
Val	Ile	Ala	Thr	Trp	Arg	Arg	Lys	Glu	Ala	Trp	Arg	Arg	Asp	Cys			
				20					25					30			
Leu	Leu	Gly	Ala	Leu	Pro	Ser	Val	Ser	Cys	Gly	Gly	Trp	Gly	His			
				35					40					45			
Arg	Gly	Arg	Gln	Thr	Tyr	Gly	Arg	Ala	Cys	Gly	Val	Lys	Glu	Lys			
				50					55					60			
Pro	Phe	Ser	Leu	Leu	Gly	Pro	Gln	Ile	Thr	Val	Tyr	Ala	Val	Trp			
				65					70					75			
Pro	Gln	Ser	Glu	Gly	Pro	Gln	Glu	Gly	Arg	Leu	Arg	Val	Asn	Ser			
				80					85					90			
Ala	Cys	Leu	Pro	Pro	Glu	Arg	Gly	Leu	Thr	Asn	Ala	Cys	Thr	Asn			
				95					100					105			
His	Glu	Glu	Leu	Ser	Leu	Asp	Cys	Leu	Leu	Phe	Glu	Asn	Val	Asn			
				110					115					120			
Thr	Leu	Thr	Leu	Asp	Phe	Cys	Leu	Trp	Glu	Lys	Thr	Thr	Ile	Val			
				125					130					135			
Pro	Gly	Val	Leu	Pro	Tyr	Ala	Gly	Leu	Thr	Leu	Gln	Ser	Lys	Phe			
				140					145					150			
Leu	Leu	Gly	Arg	Ala	Leu	Leu	Ala	Gly	Val	His	Val	Ile	Thr	Leu			
				155					160					165			
Thr	Pro	Glu	Arg	Val	Thr	His	His	Val	His	Gly	Trp	Tyr	Met	Glu			

Asp Gly Phe Lys	170	Asp Arg Thr Glu	175	Cys Arg Ser Asp	180
Val Ala Val Pro	185	Ala Ala Pro Val	190	Gln Pro Lys Ser	195
Thr Asn Gly Gln	200	Pro Ala Pro Ala	205	Thr Pro Thr Pro	210
Leu Ser Ile Ser	215	Ser Arg Ala Thr	220	Val Ala Arg Met	225
Thr Ser Gln Gly	230	Gly Leu Gln Thr	235	Met Lys Trp Lys	240
Val Ala Ile Phe	245	Val Val Val Val	250	Leu Val Thr Gly	255
Leu Val Phe Arg	260	Ala Leu Glu Gln	265	Pro Phe Glu Ser	270
Asn Thr Ile Ala	275	Leu Glu Lys Ala	280	Ser Ser Gln Lys	285
Cys Val Ser Pro	290	Gln Glu Leu Glu	295	Thr Leu Ile Gln	300
Asp Ala Asp Asn	305	Ala Gly Val Ser	310	Pro Ile Gly Asn	315
Asn Ser Ser His	320	Trp Asp Leu Gly	325	Ser Ser Asn	330
Thr Val Ile Thr	335	Thr Met Tyr Gly	340	Ala Phe Phe Phe	345
Gly Gly Lys Ile	350	Thr Cys Ile Leu	355	Asn Ile Ala Pro	360
Leu Phe Gly Phe	365	Thr Phe Cys Ile	370	Thr Ala Ile Phe	375
Ile Phe Gly Lys	380	Leu Leu Ala Gly	385	Gly Asp Gln Leu	390
Lys Gln Val Ser	395	Ser Ile Ala Arg	400	Val Glu Lys Val	405
Phe Ile Leu Ala	410	Gly Thr Lys Ile	415	Arg Val Ile Ser	420
Ile Phe Lys Tyr	425	Thr Cys Ile Val	430	Phe Val Thr Ile	435
Phe Val Val Val	440	Gly Glu Gly Trp	445	Thr Ala Leu Glu	450
Ala Val Val Val	455	Thr Leu Thr Thr	460	Val Gly Phe Gly	465
Trp Tyr Lys Pro	470	Val Phe Arg Gly	475	Asp Ile Asn Tyr	480
Tyr Phe Ala Ala	485	Leu Val Trp Phe	490	Trp Ile Leu Val	495
Leu Ser Lys Lys	500	Val Leu Ser Met	505	Gly Asp Trp Leu	510
Ala Ala Glu Trp	515	Thr Lys Glu Glu	520	Val Gly Glu Ile	525
Arg Arg Arg Leu	530	Lys Ala Asn Val	535	Thr Ala Glu Phe	540
Ala Thr Ile Arg	545	Ser Val Glu Ile	550	Arg Asp Lys Leu	555
Arg Ala His Ser	560	Met Glu Arg Arg	565	Gly Leu Gly Leu	570
Phe Ala Ala Leu	575	Asp Met Leu Ser	580	Pro Glu Lys Arg	585
Ser Ile Asn Asn	590	Thr Asp Gly Arg	595	Ser Phe Ala Ser	600
Gln Leu Asn Lys	605	Arg Pro Asn Asn	610	Gly Leu Lys Gly	615
Asn Lys Phe Gly	620	His Gly Gln Gly	625	Ala Ser Glu Asp	630
Asp Leu Lys Lys	635	Ser Thr Ser Arg	640	Leu Thr Lys Arg	645
Thr Phe Arg Asn	650	Thr Leu Pro Glu	655	Asp Val Gln Lys	660
	665	Tyr Ser Leu Asp	670	Glu Lys Lys Glu	675

Thr	Glu	Lys	Met	Cys	Asn	Ser	Asp	Asn	Ser	Ser	Thr	Ala	Met	Leu
				680					685					690
Thr	Asp	Cys	Ile	Gln	Gln	His	Ala	Glu	Leu	Glu	Asn	Gly	Met	Ile
				695					700					705
Pro	Thr	Asp	Thr	Lys	Asp	Arg	Glu	Pro	Glu	Asn	Asn	Ser	Leu	Leu
				710					715					720
Glu	Asp	Arg	Asn											

<210> 11
 <211> 470
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7474322CD1

<220>
 <221> unsure
 <222> 253
 <223> unknown or other

<400> 11

Met	Tyr	Asn	Glu	Ile	Leu	Met	Leu	Gly	Ala	Lys	Leu	His	Pro	Thr
1				5					10					15
Leu	Lys	Leu	Glu	Glu	Leu	Thr	Asn	Lys	Lys	Gly	Met	Thr	Pro	Leu
				20					25					30
Ala	Leu	Ala	Ala	Gly	Thr	Gly	Lys	Ile	Gly	Asn	Arg	His	Asp	Met
				35					40					45
Leu	Leu	Val	Glu	Pro	Leu	Asn	Arg	Leu	Leu	Gln	Asp	Lys	Trp	Asp
				50					55					60
Arg	Phe	Val	Lys	Arg	Ile	Phe	Tyr	Phe	Asn	Phe	Leu	Val	Tyr	Cys
				65					70					75
Leu	Tyr	Met	Ile	Ile	Phe	Thr	Met	Ala	Ala	Tyr	Tyr	Arg	Pro	Val
				80					85					90
Asp	Gly	Leu	Pro	Pro	Phe	Lys	Met	Glu	Lys	Thr	Gly	Asp	Tyr	Phe
				95					100					105
Arg	Val	Thr	Gly	Glu	Ile	Leu	Ser	Val	Leu	Gly	Gly	Val	Tyr	Phe
				110					115					120
Phe	Phe	Arg	Gly	Ile	Gln	Tyr	Phe	Leu	Gln	Arg	Arg	Pro	Ser	Met
				125					130					135
Lys	Thr	Leu	Phe	Val	Asp	Ser	Tyr	Ser	Glu	Met	Leu	Leu	Phe	Leu
				140					145					150
Gln	Ser	Leu	Phe	Met	Leu	Ala	Thr	Val	Val	Leu	Tyr	Phe	Ser	His
				155					160					165
Leu	Lys	Glu	Tyr	Val	Ala	Ser	Met	Val	Phe	Ser	Leu	Ala	Leu	Gly
				170					175					180
Trp	Thr	Asn	Met	Leu	Tyr	Tyr	Thr	Arg	Gly	Phe	Gln	Gln	Met	Gly
				185					190					195
Ile	Tyr	Ala	Val	Met	Ile	Glu	Lys	Met	Ile	Leu	Arg	Asp	Leu	Cys
				200					205					210
Arg	Phe	Met	Phe	Val	Tyr	Ile	Val	Phe	Leu	Phe	Gly	Phe	Ser	Thr
				215					220					225
Ala	Val	Val	Thr	Leu	Ile	Glu	Asp	Gly	Lys	Asn	Asp	Ser	Leu	Pro
				230					235					240
Ser	Glu	Ser	Thr	Ser	His	Arg	Trp	Arg	Gly	Pro	Ala	Xaa	Arg	Pro
				245					250					255
Asn	Ser	Ser	Tyr	Asn	Ser	Leu	Tyr	Ser	Thr	Cys	Leu	Glu	Leu	Phe
				260					265					270
Lys	Phe	Thr	Ile	Gly	Met	Gly	Asp	Leu	Glu	Phe	Thr	Glu	Asn	Tyr
				275					280					285
Asp	Phe	Lys	Ala	Val	Phe	Ile	Ile	Leu	Leu	Leu	Ala	Tyr	Val	Ile
				290					295					300
Leu	Thr	Tyr	Ile	Val	Leu	Leu	Leu	Asn	Met	Leu	Ile	Ala	Leu	Met
				305					310					315
Gly	Glu	Thr	Val	Glu	Asn	Val	Ser	Lys	Glu	Ser	Glu	Arg	Ile	Trp

Arg	Leu	Gln	Arg	320	Ile	Thr	Ile	Leu	325	Thr	Glu	Lys	Ser	330
				335	Ala				340	Asp				345
Leu	Lys	Cys	Met	350	Arg	Lys	Ala	Phe	355	Ser	Gly	Lys	Leu	360
Val	Gly	Tyr	Thr	365	Pro	Asp	Gly	Lys	370	Asp	Tyr	Arg	Trp	375
Val	Asp	Glu	Val	380	Asn	Trp	Thr	Thr	385	Asn	Thr	Asn	Val	390
Ile	Asn	Glu	Asp	395	Pro	Gly	Asn	Cys	400	Gly	Val	Lys	Arg	405
Ser	Phe	Ser	Leu	410	Arg	Ser	Ser	Arg	415	Ser	Gly	Arg	His	420
Asn	Phe	Ala	Leu	425	Val	Pro	Leu	Leu	430	Glu	Ala	Ser	Ala	435
Arg	Gln	Ser	Ala	440	Gln	Pro	Glu	Glu	445	Tyr	Leu	Arg	Gln	450
Gly	Ser	Leu	Lys	455	Pro	Glu	Asp	Ala	460	Val	Phe	Lys	Ser	465
Ala	Ser	Gly	Glu	470	Lys									

<210> 12

<211> 618

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5455621CD1

<400> 12

Met	Glu	Val	Lys	Asn	Phe	Ala	Val	Trp	Asp	Tyr	Val	Val	Phe	Ala
1				5					10					15
Ala	Leu	Phe	Phe	Ile	Ser	Ser	Gly	Ile	Gly	Val	Phe	Phe	Ala	Ile
				20					25					30
Lys	Glu	Arg	Lys	Lys	Ala	Thr	Ser	Arg	Glu	Phe	Leu	Val	Gly	Gly
				35					40					45
Arg	Gln	Met	Ser	Phe	Gly	Pro	Val	Gly	Leu	Ser	Leu	Thr	Ala	Ser
				50					55					60
Phe	Met	Ser	Ala	Val	Thr	Val	Leu	Gly	Thr	Pro	Ser	Glu	Val	Tyr
				65					70					75
Arg	Phe	Gly	Ala	Ser	Phe	Leu	Val	Phe	Phe	Ile	Ala	Tyr	Leu	Phe
				80					85					90
Val	Ile	Leu	Leu	Thr	Ser	Glu	Leu	Phe	Leu	Pro	Val	Phe	Tyr	Arg
				95					100					105
Ser	Gly	Ile	Thr	Ser	Thr	Tyr	Glu	Tyr	Leu	Gln	Leu	Arg	Phe	Asn
				110					115					120
Lys	Pro	Val	Arg	Tyr	Ala	Ala	Thr	Val	Ile	Tyr	Ile	Val	Gln	Thr
				125					130					135
Ile	Leu	Tyr	Thr	Gly	Val	Val	Val	Tyr	Ala	Pro	Ala	Leu	Ala	Leu
				140					145					150
Asn	Gln	Val	Thr	Gly	Phe	Asp	Leu	Trp	Gly	Ser	Val	Phe	Ala	Thr
				155					160					165
Gly	Ile	Val	Cys	Thr	Phe	Tyr	Cys	Thr	Leu	Gly	Gly	Leu	Lys	Ala
				170					175					180
Val	Val	Trp	Thr	Asp	Ala	Phe	Gln	Met	Val	Val	Met	Ile	Val	Gly
				185					190					195
Phe	Leu	Thr	Val	Leu	Ile	Gln	Gly	Ser	Thr	His	Ala	Gly	Gly	Phe
				200					205					210
His	Asn	Val	Leu	Glu	Gln	Ser	Thr	Asn	Gly	Ser	Arg	Leu	His	Ile
				215					220					225
Phe	Asp	Phe	Asp	Val	Asp	Pro	Leu	Arg	Arg	His	Thr	Phe	Trp	Thr
				230					235					240
Ile	Thr	Val	Gly	Gly	Thr	Phe	Thr	Trp	Leu	Gly	Ile	Tyr	Gly	Val
				245					250					255
Asn	Gln	Ser	Thr	Ile	Gln	Arg	Cys	Ile	Ser	Cys	Lys	Thr	Glu	Lys

His	Ala	Lys	Leu	260	Ala	Leu	Tyr	Phe	Asn	265	Leu	Leu	Gly	Leu	Trp	270	Ile
Ile	Leu	Val	Cys	275	Ala	Val	Phe	Ser	Gly	280	Leu	Ile	Met	Tyr	Ser	285	His
Phe	Lys	Asp	Cys	290	Asp	Pro	Trp	Thr	Ser	295	Gly	Ile	Ile	Ser	Ala	300	Pro
Asp	Gln	Leu	Met	305	Pro	Tyr	Phe	Val	Met	310	Glu	Ile	Phe	Ala	Thr	315	Met
Pro	Gly	Leu	Pro	320	Gly	Leu	Phe	Val	Ala	325	Cys	Ala	Phe	Ser	Gly	330	Thr
Leu	Ser	Thr	Val	335	Ala	Ser	Ser	Ile	Asn	340	Ala	Leu	Ala	Thr	Val	345	Thr
Phe	Glu	Asp	Phe	350	Val	Lys	Ser	Cys	Phe	355	Pro	His	Leu	Ser	Asp	360	Lys
Leu	Ser	Thr	Trp	365	Ile	Ser	Lys	Gly	Leu	370	Cys	Leu	Leu	Phe	Gly	375	Val
Met	Cys	Thr	Ser	380	Met	Ala	Val	Ala	Ala	385	Ser	Val	Met	Gly	Gly	390	Val
Val	Gln	Ala	Ser	395	Leu	Ser	Ile	His	Gly	400	Met	Cys	Gly	Gly	Pro	405	Met
Leu	Gly	Leu	Phe	410	Ser	Leu	Gly	Ile	Val	415	Phe	Pro	Phe	Val	Asn	420	Trp
Lys	Gly	Ala	Leu	425	Gly	Gly	Leu	Leu	Thr	430	Gly	Ile	Thr	Leu	Ser	435	Phe
Trp	Val	Ala	Ile	440	Gly	Ala	Phe	Ile	Tyr	445	Pro	Ala	Pro	Ala	Ser	450	Lys
Thr	Trp	Pro	Leu	455	Pro	Leu	Ser	Thr	Asp	460	Gln	Cys	Ile	Lys	Ser	465	Asn
Val	Thr	Ala	Thr	470	Gly	Pro	Pro	Val	Leu	475	Ser	Ser	Arg	Pro	Gly	480	Ile
Ala	Asp	Thr	Trp	485	Tyr	Ser	Ile	Ser	Tyr	490	Leu	Tyr	Tyr	Ser	Ala	495	Leu
Gly	Cys	Leu	Gly	500	Cys	Ile	Val	Ala	Gly	505	Val	Ile	Ile	Ser	Leu	510	Ile
Thr	Gly	Arg	Gln	515	Arg	Gly	Glu	Asp	Ile	520	Gln	Pro	Leu	Leu	Ile	525	Arg
Pro	Val	Cys	Asn	530	Leu	Phe	Cys	Phe	Trp	535	Ser	Lys	Lys	Tyr	Lys	540	Thr
Leu	Cys	Trp	Cys	545	Gly	Val	Gln	His	Asp	550	Ser	Gly	Thr	Glu	Gln	555	Glu
Asn	Leu	Glu	Asn	560	Gly	Ser	Ala	Arg	Lys	565	Gln	Gly	Ala	Glu	Ser	570	Val
Leu	Gln	Asn	Gly	575	Leu	Arg	Arg	Glu	Ser	580	Leu	Val	His	Val	Pro	585	Gly
Tyr	Asp	Pro	Lys	590	Asp	Lys	Ser	Tyr	Asn	595	Asn	Met	Ala	Phe	Glu	600	Thr
Thr	His	Phe		605						610						615	

<210> 13
 <211> 631
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7477248CD1

<400> 13
 Met Glu Arg Gln Ser Arg Val Met Ser Glu Lys Asp Glu Tyr Gln
 1 5 10 15
 Phe Gln His Gln Gly Ala Val Glu Leu Leu Val Phe Asn Phe Leu
 20 25 30
 Leu Ile Leu Thr Ile Leu Thr Ile Trp Leu Phe Lys Asn His Arg
 35 40 45
 Phe Arg Phe Leu His Glu Thr Gly Gly Ala Met Val Tyr Gly Leu

Ile Met Gly Leu	Ile 50	Leu Arg Tyr Ala	Thr 55	Ala Pro Thr Asp	Ile 60
Glu Ser Gly Thr	Val 65	Tyr Asp Cys Val	Lys 70	Leu Thr Phe Ser	Pro 75
Ser Thr Leu Leu	Val 80	Asn Ile Thr Asp	Gln 85	Val Tyr Glu Tyr	Lys 90
Tyr Lys Arg Glu	Ile 95	Ser Gln His Asn	Ile 100	Asn Pro His Gln	Gly 105
Asn Ala Ile Leu	Glu 110	Lys Met Thr Phe	Asp 115	Pro Glu Ile Phe	Phe 120
Asn Val Leu Leu	Pro 125	Pro Ile Ile Phe	His 130	Ala Gly Tyr Ser	Leu 135
Lys Lys Arg His	Phe 140	Phe Gln Asn Leu	Gly 145	Ser Ile Leu Thr	Tyr 150
Ala Phe Leu Gly	Thr 155	Ala Ile Ser Cys	Ile 160	Val Ile Gly Leu	Ile 165
Met Tyr Gly Phe	Val 170	Lys Ala Met Ile	His 175	Ala Gly Gln Leu	Lys 180
Asn Gly Asp Phe	His 185	Phe Thr Asp Cys	Leu 190	Phe Phe Gly Ser	Leu 195
Met Ser Ala Thr	Asp 200	Pro Val Thr Val	Leu 205	Ala Ile Phe His	Glu 210
Leu His Val Asp	Pro 215	Asp Leu Tyr Thr	Leu 220	Leu Phe Gly Glu	Ser 225
Val Leu Asn Asp	Ala 230	Val Ala Ile Val	Leu 235	Thr Tyr Ser Ile	Ser 240
Ile Tyr Ser Pro	Lys 245	Glu Asn Pro Asn	Ala 250	Phe Asp Ala Ala	Ala 255
Phe Phe Gln Ser	Val 260	Gly Asn Phe Leu	Gly 265	Ile Phe Ala Gly	Ser 270
Phe Ala Met Gly	Ser 275	Ala Tyr Ala Ile	Ile 280	Thr Ala Leu Leu	Thr 285
Lys Phe Thr Lys	Leu 290	Cys Glu Phe Pro	Met 295	Leu Glu Thr Gly	Leu 300
Phe Phe Leu Leu	Ser 305	Trp Ser Ala Phe	Leu 310	Ser Ala Glu Ala	Ala 315
Gly Leu Thr Gly	Ile 320	Val Ala Val Leu	Phe 325	Cys Gly Val Thr	Gln 330
Ala His Tyr Thr	Tyr 335	Asn Asn Leu Ser	Ser 340	Asp Ser Lys Ile	Arg 345
Thr Lys Gln Leu	Phe 350	Glu Phe Met Asn	Phe 355	Leu Ala Glu Asn	Val 360
Ile Phe Cys Tyr	Met 365	Gly Leu Ala Leu	Phe 370	Thr Phe Gln Asn	His 375
Ile Phe Asn Ala	Leu 380	Phe Ile Leu Gly	Ala 385	Phe Leu Ala Ile	Phe 390
Val Ala Arg Ala	Cys 395	Asn Ile Tyr Pro	Leu 400	Ser Phe Leu Leu	Asn 405
Leu Gly Arg Lys	Gln 410	Lys Ile Pro Trp	Asn 415	Phe Gln His Met	Met 420
Met Phe Ser Gly	Leu 425	Arg Gly Ala Ile	Ala 430	Phe Ala Leu Ala	Ile 435
Arg Asn Thr Glu	Ser 440	Gln Pro Lys Gln	Met 445	Met Phe Thr Thr	Thr 450
Leu Leu Leu Val	Phe 455	Phe Thr Val Trp	Val 460	Phe Gly Gly Gly	Thr 465
Thr Pro Met Leu	Thr 470	Trp Leu Gln Ile	Arg 475	Val Gly Val Asp	Leu 480
Asp Glu Asn Leu	Lys 485	Glu Asp Pro Ser	Ser 490	Gln His Gln Glu	Ala 495
Asn Asn Leu Asp	Lys 500	Asn Met Thr Lys	Ala 505	Glu Ser Ala Arg	Leu 510
Phe Arg Met Trp	Tyr 515	Ser Phe Asp His	Lys 520	Tyr Leu Lys Pro	Ile 525
Leu Thr His Ser	Gly 530	Pro Pro Leu Thr	Thr 535	Thr Leu Pro Glu	Trp 540
	545		550		555

Cys	Gly	Pro	Ile	Ser	Arg	Leu	Leu	Thr	Ser	Pro	Gln	Ala	Tyr	Gly
				560					565					570
Glu	Gln	Leu	Lys	Glu	Asp	Asp	Val	Glu	Cys	Ile	Val	Asn	Gln	Asp
				575					580					585
Glu	Leu	Ala	Ile	Asn	Tyr	Gln	Glu	Gln	Ala	Ser	Ser	Pro	Cys	Ser
				590					595					600
Pro	Pro	Ala	Arg	Leu	Gly	Leu	Asp	Gln	Lys	Ala	Ser	Pro	Gln	Thr
				605					610					615
Pro	Gly	Lys	Glu	Asn	Ile	Tyr	Glu	Gly	Asp	Leu	Gly	Pro	Gly	Arg
				620					625					630
Leu														

<210> 14
 <211> 1256
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2944004CD1

<400> 14

Met	Asp	Arg	Glu	Glu	Arg	Lys	Thr	Ile	Asn	Gln	Gly	Gln	Glu	Asp
1				5					10					15
Glu	Met	Glu	Ile	Tyr	Gly	Tyr	Asn	Leu	Ser	Arg	Trp	Lys	Leu	Ala
				20					25					30
Ile	Val	Ser	Leu	Gly	Val	Ile	Cys	Ser	Gly	Gly	Val	Ser	Pro	Pro
				35					40					45
Pro	Leu	Tyr	Trp	Met	Pro	Glu	Trp	Arg	Val	Lys	Ala	Thr	Cys	Val
				50					55					60
Arg	Ala	Ala	Ile	Lys	Asp	Cys	Glu	Val	Val	Leu	Leu	Arg	Thr	Thr
				65					70					75
Asp	Glu	Phe	Lys	Met	Trp	Phe	Cys	Ala	Lys	Ile	Arg	Val	Leu	Ser
				80					85					90
Leu	Glu	Thr	Tyr	Pro	Val	Ser	Ser	Pro	Lys	Ser	Met	Ser	Asn	Lys
				95					100					105
Leu	Ser	Asn	Gly	His	Ala	Val	Cys	Leu	Ile	Glu	Asn	Pro	Thr	Glu
				110					115					120
Glu	Asn	Arg	His	Arg	Ile	Ser	Lys	Tyr	Ser	Gln	Thr	Glu	Ser	Gln
				125					130					135
Gln	Ile	Arg	Tyr	Phe	Thr	His	His	Ser	Val	Lys	Tyr	Phe	Trp	Asn
				140					145					150
Asp	Thr	Ile	His	Asn	Phe	Asp	Phe	Leu	Lys	Gly	Leu	Asp	Glu	Gly
				155					160					165
Val	Ser	Cys	Thr	Ser	Ile	Tyr	Glu	Lys	His	Ser	Ala	Gly	Leu	Thr
				170					175					180
Lys	Gly	Met	His	Ala	Tyr	Arg	Lys	Leu	Leu	Tyr	Gly	Val	Asn	Glu
				185					190					195
Ile	Ala	Val	Lys	Val	Pro	Ser	Val	Phe	Lys	Leu	Leu	Ile	Lys	Glu
				200					205					210
Val	Leu	Asn	Pro	Phe	Tyr	Ile	Phe	Gln	Leu	Phe	Ser	Val	Ile	Leu
				215					220					225
Trp	Ser	Thr	Asp	Glu	Tyr	Tyr	Tyr	Tyr	Ala	Leu	Ala	Ile	Val	Val
				230					235					240
Met	Ser	Ile	Val	Ser	Ile	Val	Ser	Ser	Leu	Tyr	Ser	Ile	Arg	Lys
				245					250					255
Gln	Tyr	Val	Met	Leu	His	Asp	Met	Val	Ala	Thr	His	Ser	Thr	Val
				260					265					270
Arg	Val	Ser	Val	Cys	Arg	Val	Asn	Glu	Glu	Ile	Glu	Glu	Ile	Phe
				275					280					285
Ser	Thr	Asp	Leu	Val	Pro	Gly	Asp	Val	Met	Val	Ile	Pro	Leu	Asn
				290					295					300
Gly	Thr	Ile	Met	Pro	Cys	Asp	Ala	Val	Leu	Ile	Asn	Gly	Thr	Cys
				305					310					315
Ile	Val	Asn	Glu	Ser	Met	Leu	Thr	Gly	Glu	Ser	Val	Pro	Val	Thr
				320					325					330

Lys	Thr	Asn	Leu	Pro	Asn	Pro	Ser	Val	Asp	Val	Lys	Gly	Ile	Gly
				335					340					345
Asp	Glu	Leu	Tyr	Asn	Pro	Glu	Thr	His	Lys	Arg	His	Thr	Leu	Phe
				350					355					360
Cys	Gly	Thr	Thr	Val	Ile	Gln	Thr	Arg	Phe	Tyr	Thr	Gly	Glu	Leu
				365					370					375
Val	Lys	Ala	Ile	Val	Val	Arg	Thr	Gly	Phe	Ser	Thr	Ser	Lys	Gly
				380					385					390
Gln	Leu	Val	Arg	Ser	Ile	Leu	Tyr	Pro	Lys	Pro	Thr	Asp	Phe	Lys
				395					400					405
Leu	Tyr	Arg	Asp	Ala	Tyr	Leu	Phe	Leu	Leu	Cys	Leu	Val	Ala	Val
				410					415					420
Ala	Gly	Ile	Gly	Phe	Ile	Tyr	Thr	Ile	Ile	Asn	Ser	Ile	Leu	Asn
				425					430					435
Glu	Val	Gln	Val	Gly	Val	Ile	Ile	Ile	Glu	Ser	Leu	Asp	Ile	Ile
				440					445					450
Thr	Ile	Thr	Val	Pro	Pro	Ala	Leu	Pro	Ala	Ala	Met	Thr	Ala	Gly
				455					460					465
Ile	Val	Tyr	Ala	Gln	Arg	Arg	Leu	Lys	Lys	Ile	Gly	Ile	Phe	Cys
				470					475					480
Ile	Ser	Pro	Gln	Arg	Ile	Asn	Ile	Cys	Gly	Gln	Leu	Asn	Leu	Val
				485					490					495
Cys	Phe	Asp	Lys	Thr	Gly	Thr	Leu	Thr	Glu	Asp	Gly	Leu	Asp	Leu
				500					505					510
Trp	Gly	Ile	Gln	Arg	Val	Glu	Asn	Ala	Arg	Phe	Leu	Ser	Pro	Glu
				515					520					525
Glu	Asn	Val	Cys	Asn	Glu	Met	Leu	Val	Lys	Ser	Gln	Phe	Val	Ala
				530					535					540
Cys	Met	Ala	Thr	Cys	His	Ser	Leu	Thr	Lys	Ile	Glu	Gly	Val	Leu
				545					550					555
Ser	Gly	Asp	Pro	Leu	Asp	Leu	Lys	Met	Phe	Glu	Ala	Ile	Gly	Trp
				560					565					570
Ile	Leu	Glu	Glu	Ala	Thr	Glu	Glu	Glu	Thr	Ala	Leu	His	Asn	Arg
				575					580					585
Ile	Met	Pro	Thr	Val	Val	Arg	Pro	Pro	Lys	Gln	Leu	Leu	Pro	Glu
				590					595					600
Ser	Thr	Pro	Ala	Gly	Asn	Gln	Glu	Met	Glu	Leu	Phe	Glu	Leu	Pro
				605					610					615
Ala	Thr	Tyr	Glu	Ile	Gly	Ile	Val	Arg	Gln	Phe	Pro	Phe	Ser	Ser
				620					625					630
Ala	Leu	Gln	Arg	Met	Ser	Val	Val	Ala	Arg	Val	Leu	Gly	Asp	Arg
				635					640					645
Lys	Met	Asp	Ala	Tyr	Met	Lys	Gly	Ala	Pro	Glu	Ala	Ile	Ala	Gly
				650					655					660
Leu	Cys	Lys	Pro	Glu	Thr	Val	Pro	Val	Asp	Phe	Gln	Asn	Val	Leu
				665					670					675
Glu	Asp	Phe	Thr	Lys	Gln	Gly	Phe	Arg	Val	Ile	Ala	Leu	Ala	His
				680					685					690
Arg	Lys	Leu	Glu	Ser	Lys	Leu	Thr	Trp	His	Lys	Val	Gln	Asn	Ile
				695					700					705
Ser	Arg	Asp	Ala	Ile	Glu	Asn	Asn	Met	Asp	Phe	Met	Gly	Leu	Ile
				710					715					720
Ile	Met	Gln	Asn	Lys	Leu	Lys	Gln	Lys	Thr	Pro	Ala	Val	Leu	Glu
				725					730					735
Asp	Leu	His	Lys	Ala	Asn	Ile	Arg	Thr	Val	Met	Val	Thr	Gly	Asp
				740					745					750
Ser	Met	Leu	Thr	Ala	Val	Ser	Val	Ala	Arg	Asp	Cys	Gly	Met	Ile
				755					760					765
Leu	Pro	Gln	Asp	Lys	Val	Ile	Ile	Ala	Glu	Ala	Leu	Pro	Pro	Lys
				770					775					780
Asp	Gly	Lys	Val	Ala	Lys	Ile	Asn	Trp	His	Tyr	Ala	Asp	Ser	Leu
				785					790					795
Thr	Gln	Cys	Ser	His	Pro	Ser	Ala	Ile	Asp	Pro	Glu	Ala	Ile	Pro
				800					805					810
Val	Lys	Leu	Val	His	Asp	Ser	Leu	Glu	Asp	Leu	Gln	Met	Thr	Arg
				815					820					825
Tyr	His	Phe	Ala	Met	Asn	Gly	Lys	Ser	Phe	Ser	Val	Ile	Leu	Glu

His Phe Gln Asp	830	Val Pro Lys Leu	835	Leu His Gly Thr	840
	845		850		855
Phe Ala Arg Met	860	Ala Pro Asp Gln Lys	865	Gln Leu Ile Glu	870
	875		880		885
Leu Gln Asn Val	890	Asp Tyr Phe Val Gly	895	Met Cys Gly Asp Gly	900
	905		910		915
Asn Asp Cys Gly	920	Ala Leu Lys Arg Ala	925	His Gly Gly Ile Ser	930
	935		940		945
Ser Glu Leu Glu	950	Ala Ser Val Ala Ser	955	Pro Phe Thr Ser Lys	960
	965		970		975
Pro Ser Ile Ser	980	Cys Val Pro Asn Leu	985	Arg Glu Gly Arg	990
	995		1000		1005
Ala Leu Ile Thr	1010	Ser Phe Cys Val Phe	1015	Lys Phe Met Ala Leu	1020
	1025		1030		1035
Ser Ile Ile Gln	1040	Tyr Phe Ser Val Thr	1045	Leu Tyr Ser Ile	1050
	1055		1060		1065
Ser Asn Leu Gly	1070	Asp Phe Gln Phe Leu	1075	Phe Ile Asp Leu Ala	1080
	1085		1090		1095
Ile Leu Val Val	1100	Val Phe Thr Met Ser	1105	Leu Asn Pro Ala Trp	1110
	1115		1120		1125
Glu Leu Val Ala	1130	Gln Arg Pro Pro Ser	1135	Gly Leu Ile Ser Gly	1140
	1145		1150		1155
Leu Leu Phe Ser	1160	Val Leu Ser Gln Ile	1165	Ile Ile Cys Ile Gly	1170
	1175		1180		1185
Gln Ser Leu Gly	1190	Phe Trp Val Lys Gln	1195	Phe Gln Pro Trp Tyr	1200
	1205		1210		1215
Val Trp His Pro	1220	Lys Ser Asp Ala Cys	1225	Asn Thr Thr Gly Ser	1230
	1235		1240		1245
Phe Trp Asn Ser	1250	Ser His Val Asp Asn		Glu Thr Glu Leu Asp	
His Asn Ile Gln		Asn Tyr Glu Asn Thr		Thr Val Phe Phe Ile	
Ser Phe Gln Tyr		Leu Ile Val Ala Ile		Ala Phe Ser Lys Gly	
Pro Phe Arg Gln		Pro Cys Tyr Lys Asn		Tyr Phe Phe Val Phe	
Val Ile Phe Leu		Tyr Ile Phe Ile Leu		Phe Ile Met Leu Tyr	
Val Ala Ser Val		Asp Gln Val Leu Gln		Ile Val Cys Val Pro	
Gln Trp Arg Val		Thr Met Leu Ile Ile		Val Leu Val Asn Ala	
Val Ser Ile Thr		Val Glu Asn Phe Phe		Leu Asp Met Val Leu	
Lys Val Val Phe		Asn Arg Asp Lys Gln		Gly Glu Tyr Arg Phe	
Thr Thr Gln Pro		Pro Gln Glu Ser Val		Asp Arg Trp Gly Lys	
Cys Leu Pro Trp		Ala Leu Gly Cys Arg		Lys Lys Thr Pro Lys	
Lys Tyr Met Tyr		Leu Ala Gln Glu Leu		Leu Val Asp Pro Glu	
Pro Pro Lys Pro		Gln Thr Thr Thr Glu		Ala Lys Ala Leu Val	
Glu Asn Gly Ser		Cys Gln Ile Ile Thr		Ile Thr	

<210> 15
 <211> 499
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3046849CD1

<400> 15

Met	Leu	His	Ala	Leu	Leu	Arg	Ser	Arg	Thr	Ile	Gln	Gly	Arg	Ile
1				5					10					15
Leu	Leu	Leu	Thr	Ile	Cys	Ala	Ala	Gly	Ile	Gly	Gly	Thr	Phe	Gln
				20					25					30
Phe	Gly	Tyr	Asn	Leu	Ser	Ile	Ile	Asn	Ala	Pro	Thr	Leu	His	Ile
				35					40					45
Gln	Glu	Phe	Thr	Asn	Glu	Thr	Trp	Gln	Ala	Arg	Thr	Gly	Glu	Pro
				50					55					60
Leu	Pro	Asp	His	Leu	Val	Leu	Leu	Met	Trp	Ser	Leu	Ile	Val	Ser
				65					70					75
Leu	Tyr	Pro	Leu	Gly	Gly	Leu	Phe	Gly	Ala	Leu	Leu	Ala	Gly	Pro
				80					85					90
Leu	Ala	Ile	Thr	Leu	Gly	Arg	Lys	Lys	Ser	Leu	Leu	Val	Asn	Asn
				95					100					105
Ile	Phe	Val	Val	Ser	Ala	Ala	Ile	Leu	Phe	Gly	Phe	Ser	Arg	Lys
				110					115					120
Ala	Gly	Ser	Phe	Glu	Met	Ile	Met	Leu	Gly	Arg	Leu	Leu	Val	Gly
				125					130					135
Val	Asn	Ala	Gly	Val	Ser	Met	Asn	Ile	Gln	Pro	Met	Tyr	Leu	Gly
				140					145					150
Glu	Ser	Ala	Pro	Lys	Glu	Leu	Arg	Gly	Ala	Val	Ala	Met	Ser	Ser
				155					160					165
Ala	Ile	Phe	Thr	Ala	Leu	Gly	Ile	Val	Met	Gly	Gln	Val	Val	Gly
				170					175					180
Leu	Arg	Glu	Leu	Leu	Gly	Gly	Pro	Gln	Ala	Trp	Pro	Leu	Leu	Leu
				185					190					195
Ala	Ser	Cys	Leu	Val	Pro	Gly	Ala	Leu	Gln	Leu	Ala	Ser	Leu	Pro
				200					205					210
Leu	Leu	Pro	Glu	Ser	Pro	Arg	Tyr	Leu	Leu	Ile	Asp	Cys	Gly	Asp
				215					220					225
Thr	Glu	Ala	Cys	Leu	Ala	Ala	Leu	Arg	Gln	Leu	Arg	Gly	Ser	Gly
				230					235					240
Asp	Leu	Ala	Gly	Glu	Leu	Glu	Glu	Leu	Glu	Glu	Glu	Arg	Ala	Ala
				245					250					255
Cys	Gln	Gly	Cys	Arg	Ala	Arg	Arg	Pro	Trp	Glu	Leu	Phe	Gln	His
				260					265					270
Arg	Ala	Leu	Arg	Arg	Gln	Val	Thr	Ser	Leu	Val	Val	Leu	Gly	Ser
				275					280					285
Ala	Met	Glu	Leu	Cys	Gly	Asn	Asp	Ser	Val	Tyr	Ala	Tyr	Ala	Ser
				290					295					300
Ser	Val	Phe	Arg	Lys	Ala	Gly	Val	Pro	Glu	Ala	Lys	Ile	Gln	Tyr
				305					310					315
Ala	Ile	Ile	Gly	Thr	Gly	Ser	Cys	Glu	Leu	Leu	Thr	Ala	Val	Val
				320					325					330
Ser	Cys	Val	Val	Ile	Glu	Arg	Val	Gly	Arg	Arg	Val	Leu	Leu	Ile
				335					340					345
Gly	Gly	Tyr	Ser	Leu	Met	Thr	Cys	Trp	Gly	Ser	Ile	Phe	Thr	Val
				350					355					360
Ala	Leu	Cys	Leu	Gln	Ser	Ser	Phe	Pro	Trp	Thr	Leu	Tyr	Leu	Ala
				365					370					375
Met	Ala	Cys	Ile	Phe	Ala	Phe	Ile	Leu	Ser	Phe	Gly	Ile	Gly	Pro
				380					385					390
Ala	Gly	Val	Thr	Gly	Ile	Leu	Ala	Thr	Glu	Leu	Phe	Asp	Gln	Met
				395					400					405
Ala	Arg	Pro	Ala	Ala	Cys	Met	Val	Cys	Gly	Ala	Leu	Met	Trp	Ile
				410					415					420
Met	Leu	Ile	Leu	Val	Gly	Leu	Gly	Phe	Pro	Phe	Ile	Met	Glu	Ala
				425					430					435
Leu	Ser	His	Phe	Leu	Tyr	Val	Pro	Phe	Leu	Gly	Val	Cys	Val	Cys
				440					445					450
Gly	Ala	Ile	Tyr	Thr	Gly	Leu	Phe	Leu	Pro	Glu	Thr	Lys	Gly	Lys
				455					460					465
Thr	Phe	Gln	Glu	Ile	Ser	Lys	Glu	Leu	His	Arg	Leu	Asn	Phe	Pro
				470					475					480
Arg	Arg	Ala	Gln	Gly	Pro	Thr	Trp	Arg	Ser	Leu	Glu	Val	Ile	Gln
				485					490					495

Ser Thr Glu Leu

<210> 16
 <211> 596
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4538363CD1

<400> 16
 Met Ala Ala Asn Ser Thr Ser Asp Leu His Thr Pro Gly Thr Gln
 1 5 10 15
 Leu Ser Val Ala Asp Ile Ile Val Ile Thr Val Tyr Phe Ala Leu
 20 25 30
 Asn Val Ala Val Gly Ile Trp Ser Ser Cys Arg Ala Ser Arg Asn
 35 40 45
 Thr Val Asn Gly Tyr Phe Leu Ala Gly Arg Asp Met Thr Trp Trp
 50 55 60
 Pro Ile Gly Ala Ser Leu Phe Ala Ser Ser Glu Gly Ser Gly Leu
 65 70 75
 Phe Ile Gly Leu Ala Gly Ser Gly Ala Ala Gly Gly Leu Ala Val
 80 85 90
 Ala Gly Phe Glu Trp Asn Ala Thr Tyr Val Leu Leu Ala Leu Ala
 95 100 105
 Trp Val Phe Val Pro Ile Tyr Ile Ser Ser Glu Ile Val Thr Leu
 110 115 120
 Pro Glu Tyr Ile Gln Lys Arg Tyr Gly Gly Gln Arg Ile Arg Met
 125 130 135
 Tyr Leu Ser Val Leu Ser Leu Leu Leu Ser Val Phe Thr Lys Ile
 140 145 150
 Ser Leu Asp Leu Tyr Ala Gly Ala Leu Phe Val His Ile Cys Leu
 155 160 165
 Gly Trp Asn Phe Tyr Leu Ser Thr Ile Leu Thr Leu Gly Ile Thr
 170 175 180
 Ala Leu Tyr Thr Ile Ala Gly Gly Leu Ala Ala Val Ile Tyr Thr
 185 190 195
 Asp Ala Leu Gln Thr Leu Ile Met Val Val Gly Ala Val Ile Leu
 200 205 210
 Thr Ile Lys Ala Phe Asp Gln Ile Gly Gly Tyr Gly Gln Leu Glu
 215 220 225
 Ala Ala Tyr Ala Gln Ala Ile Pro Ser Arg Thr Ile Ala Asn Thr
 230 235 240
 Thr Cys His Leu Pro Arg Thr Asp Ala Met His Met Phe Arg Asp
 245 250 255
 Pro His Thr Gly Asp Leu Pro Trp Thr Gly Met Thr Phe Gly Leu
 260 265 270
 Thr Ile Met Ala Thr Trp Tyr Trp Cys Thr Asp Gln Val Ile Val
 275 280 285
 Gln Arg Ser Leu Ser Ala Arg Asp Leu Asn His Ala Lys Ala Gly
 290 295 300
 Ser Ile Leu Ala Ser Tyr Leu Lys Met Leu Pro Met Gly Leu Ile
 305 310 315
 Ile Met Pro Gly Met Ile Ser Arg Ala Leu Phe Pro Asp Asp Val
 320 325 330
 Gly Cys Val Val Pro Ser Glu Cys Leu Arg Ala Cys Gly Ala Glu
 335 340 345
 Val Gly Cys Ser Asn Ile Ala Tyr Pro Lys Leu Val Met Glu Leu
 350 355 360
 Met Pro Ile Gly Leu Arg Gly Leu Met Ile Ala Val Met Leu Ala
 365 370 375
 Ala Leu Met Ser Ser Leu Thr Ser Ile Phe Asn Ser Ser Ser Thr
 380 385 390
 Leu Phe Thr Met Asp Ile Trp Arg Arg Leu Arg Pro Arg Ser Gly
 395 400 405

Glu	Arg	Glu	Leu	Leu	Leu	Val	Gly	Arg	Leu	Val	Ile	Val	Ala	Leu
				410					415					420
Ile	Gly	Val	Ser	Val	Ala	Trp	Ile	Pro	Val	Leu	Gln	Asp	Ser	Asn
				425					430					435
Ser	Gly	Gln	Leu	Phe	Ile	Tyr	Met	Gln	Ser	Val	Thr	Ser	Ser	Leu
				440					445					450
Ala	Pro	Pro	Val	Thr	Ala	Val	Phe	Val	Leu	Gly	Val	Phe	Trp	Arg
				455					460					465
Arg	Ala	Asn	Glu	Gln	Gly	Ala	Phe	Trp	Gly	Leu	Ile	Ala	Gly	Leu
				470					475					480
Val	Val	Gly	Ala	Thr	Arg	Leu	Val	Leu	Glu	Phe	Leu	Asn	Pro	Ala
				485					490					495
Pro	Pro	Cys	Gly	Glu	Pro	Asp	Thr	Arg	Pro	Ala	Val	Leu	Gly	Ser
				500					505					510
Ile	His	Tyr	Leu	His	Phe	Ala	Val	Ala	Leu	Phe	Ala	Leu	Ser	Gly
				515					520					525
Ala	Val	Val	Val	Ala	Gly	Ser	Leu	Leu	Thr	Pro	Pro	Pro	Gln	Ser
				530					535					540
Val	Gln	Ile	Glu	Asn	Leu	Thr	Trp	Trp	Thr	Leu	Ala	Gln	Asp	Val
				545					550					555
Pro	Leu	Gly	Thr	Lys	Ala	Gly	Asp	Gly	Gln	Thr	Pro	Gln	Lys	His
				560					565					570
Ala	Phe	Trp	Ala	Arg	Val	Cys	Gly	Phe	Asn	Ala	Ile	Leu	Leu	Met
				575					580					585
Cys	Val	Asn	Ile	Phe	Phe	Tyr	Ala	Tyr	Phe	Ala				
				590					595					

<210> 17

<211> 1192

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6427460CD1

<400> 17

Met	Asp	Cys	Ser	Leu	Val	Arg	Thr	Leu	Val	His	Arg	Tyr	Cys	Ala
1				5					10					15
Gly	Glu	Glu	Asn	Trp	Val	Asp	Ser	Arg	Thr	Ile	Tyr	Val	Gly	His
				20					25					30
Arg	Glu	Pro	Pro	Pro	Gly	Ala	Glu	Ala	Tyr	Ile	Pro	Gln	Arg	Tyr
				35					40					45
Pro	Asp	Asn	Arg	Ile	Val	Ser	Ser	Lys	Tyr	Thr	Phe	Trp	Asn	Phe
				50					55					60
Ile	Pro	Lys	Asn	Leu	Phe	Glu	Gln	Phe	Arg	Arg	Val	Ala	Asn	Phe
				65					70					75
Tyr	Phe	Leu	Ile	Ile	Phe	Leu	Val	Gln	Leu	Ile	Ile	Asp	Thr	Pro
				80					85					90
Thr	Ser	Pro	Val	Thr	Ser	Gly	Leu	Pro	Leu	Phe	Phe	Val	Ile	Thr
				95					100					105
Val	Thr	Ala	Ile	Lys	Gln	Gly	Tyr	Glu	Asp	Trp	Leu	Arg	His	Lys
				110					115					120
Ala	Asp	Asn	Ala	Met	Asn	Gln	Cys	Pro	Val	His	Phe	Ile	Gln	His
				125					130					135
Gly	Lys	Leu	Val	Arg	Lys	Gln	Ser	Arg	Lys	Leu	Arg	Val	Gly	Asp
				140					145					150
Ile	Val	Met	Val	Lys	Glu	Asp	Glu	Thr	Phe	Pro	Cys	Asp	Leu	Ile
				155					160					165
Phe	Leu	Ser	Ser	Asn	Arg	Gly	Asp	Gly	Thr	Cys	His	Val	Thr	Thr
				170					175					180
Ala	Ser	Leu	Asp	Gly	Glu	Ser	Ser	His	Lys	Thr	His	Tyr	Ala	Val
				185					190					195
Gln	Asp	Thr	Lys	Gly	Phe	His	Thr	Glu	Glu	Asp	Ile	Gly	Gly	Leu
				200					205					210
His	Ala	Thr	Ile	Glu	Cys	Glu	Gln	Pro	Gln	Pro	Asp	Leu	Tyr	Lys
				215					220					225

Phe	Val	Gly	Arg	Ile	Asn	Val	Tyr	Ser	Asp	Leu	Asn	Asp	Pro	Val
				230					235					240
Val	Arg	Pro	Leu	Gly	Ser	Glu	Asn	Leu	Leu	Leu	Arg	Gly	Ala	Thr
				245					250					255
Leu	Lys	Asn	Thr	Glu	Lys	Ile	Phe	Gly	Val	Ala	Ile	Tyr	Thr	Gly
				260					265					270
Met	Glu	Thr	Lys	Met	Ala	Leu	Asn	Tyr	Gln	Ser	Lys	Ser	Gln	Lys
				275					280					285
Arg	Ser	Ala	Val	Glu	Lys	Ser	Met	Asn	Ala	Phe	Leu	Ile	Val	Tyr
				290					295					300
Leu	Cys	Ile	Leu	Ile	Ser	Lys	Ala	Leu	Ile	Asn	Thr	Val	Leu	Lys
				305					310					315
Tyr	Val	Trp	Gln	Ser	Glu	Pro	Phe	Arg	Asp	Glu	Pro	Trp	Tyr	Asn
				320					325					330
Gln	Lys	Thr	Glu	Ser	Glu	Arg	Gln	Arg	Asn	Leu	Phe	Leu	Lys	Ala
				335					340					345
Phe	Thr	Asp	Phe	Leu	Ala	Phe	Met	Val	Leu	Phe	Asn	Tyr	Ile	Ile
				350					355					360
Pro	Val	Ser	Met	Tyr	Val	Thr	Val	Glu	Met	Gln	Lys	Phe	Leu	Gly
				365					370					375
Ser	Tyr	Phe	Ile	Thr	Trp	Asp	Glu	Asp	Met	Phe	Asp	Glu	Glu	Thr
				380					385					390
Gly	Glu	Gly	Pro	Leu	Val	Asn	Thr	Ser	Asp	Leu	Asn	Glu	Glu	Leu
				395					400					405
Gly	Gln	Val	Glu	Tyr	Ile	Phe	Thr	Asp	Lys	Thr	Gly	Thr	Leu	Thr
				410					415					420
Glu	Asn	Asn	Met	Glu	Phe	Lys	Glu	Cys	Cys	Ile	Glu	Gly	His	Val
				425					430					435
Tyr	Val	Pro	His	Val	Ile	Cys	Asn	Gly	Gln	Val	Leu	Pro	Glu	Ser
				440					445					450
Ser	Gly	Ile	Asp	Met	Ile	Asp	Ser	Ser	Pro	Ser	Val	Asn	Gly	Arg
				455					460					465
Glu	Arg	Glu	Glu	Leu	Phe	Phe	Arg	Ala	Leu	Cys	Leu	Cys	His	Thr
				470					475					480
Val	Gln	Val	Lys	Asp	Asp	Asp	Ser	Val	Asp	Gly	Pro	Arg	Lys	Ser
				485					490					495
Pro	Asp	Gly	Gly	Lys	Ser	Cys	Val	Tyr	Ile	Ser	Ser	Ser	Pro	Asp
				500					505					510
Glu	Val	Ala	Leu	Val	Glu	Gly	Val	Gln	Arg	Leu	Gly	Phe	Thr	Tyr
				515					520					525
Leu	Arg	Leu	Lys	Asp	Asn	Tyr	Met	Glu	Ile	Leu	Asn	Arg	Glu	Asn
				530					535					540
His	Ile	Glu	Arg	Phe	Glu	Leu	Leu	Glu	Ile	Leu	Ser	Phe	Asp	Ser
				545					550					555
Val	Arg	Arg	Arg	Met	Ser	Val	Ile	Val	Lys	Ser	Ala	Thr	Gly	Glu
				560					565					570
Ile	Tyr	Leu	Phe	Cys	Lys	Gly	Ala	Asp	Ser	Ser	Ile	Phe	Pro	Arg
				575					580					585
Val	Ile	Glu	Gly	Lys	Val	Asp	Gln	Ile	Arg	Ala	Arg	Val	Glu	Arg
				590					595					600
Asn	Ala	Val	Glu	Gly	Leu	Arg	Thr	Leu	Cys	Val	Ala	Tyr	Lys	Arg
				605					610					615
Leu	Ile	Gln	Glu	Glu	Tyr	Glu	Gly	Ile	Cys	Lys	Leu	Leu	Gln	Ala
				620					625					630
Ala	Lys	Val	Ala	Leu	Gln	Asp	Arg	Glu	Lys	Lys	Leu	Ala	Glu	Ala
				635					640					645
Tyr	Glu	Gln	Ile	Glu	Lys	Asp	Leu	Thr	Leu	Leu	Gly	Ala	Thr	Ala
				650					655					660
Val	Glu	Asp	Arg	Leu	Gln	Glu	Lys	Ala	Ala	Asp	Thr	Ile	Glu	Ala
				665					670					675
Leu	Gln	Lys	Ala	Gly	Ile	Lys	Val	Trp	Val	Leu	Thr	Gly	Asp	Lys
				680					685					690
Met	Glu	Thr	Ala	Ala	Ala	Thr	Cys	Tyr	Ala	Cys	Lys	Leu	Phe	Arg
				695					700					705
Arg	Asn	Thr	Gln	Leu	Leu	Glu	Leu	Thr	Thr	Lys	Arg	Ile	Glu	Glu
				710					715					720
Gln	Ser	Leu	His	Asp	Val	Leu	Phe	Glu	Leu	Ser	Lys	Thr	Val	Leu

Arg His Ser Gly	725	Leu Thr Arg Asp	730	Leu Ser Gly Leu	735
Ala Asp Met Gln	740	Asn Ile Ile	745	Asp Gly Ala Ala	750
Ser Leu Ile Met	755	Pro Arg Glu Asp	760	Ser Ser Gly Asn	765
Arg Glu Leu Phe	770	Lys Cys Arg Ser	775	Ser Ser Gly Asn	780
Cys Cys Arg Met	785	Ala Pro Leu Gln Lys	790	Ala Gln Ile Val Lys	795
Ile Lys Phe Ser	800	Lys Glu His Pro Ile	805	Leu Ala Ile Gly	810
Gly Ala Asn Asp	815	Val Ser Met Ile Leu	820	Glu Ala His Val Gly	825
Gly Val Ile Gly	830	Lys Glu Gly Arg Gln	835	Ala Arg Asn Ser	840
Tyr Ala Ile Pro	845	Lys Phe Lys His Leu	850	Lys Lys Met Leu Leu	855
His Gly His Phe	860	Tyr Tyr Ile Arg Ile	865	Ser Glu Leu Val Gln	870
Phe Phe Tyr Lys	875	Asn Val Cys Phe Ile	880	Phe Pro Gln Phe Leu	885
Gln Phe Phe Cys	890	Gly Phe Ser Gln Gln	895	Thr Leu Tyr Asp Thr	900
Tyr Leu Thr Leu	905	Asn Ile Ser Phe Thr	910	Ser Leu Pro Ile	915
Leu Tyr Ser Leu	920	Met Glu Gln His Val	925	Gly Ile Asp Val Leu	930
Arg Asp Pro Thr	935	Leu Tyr Arg Asp Val	940	Lys Asn Ala Leu Leu	945
Arg Trp Arg Val	950	Phe Ile Tyr Trp Thr	955	Leu Leu Gly Leu Phe	960
Ala Leu Val Phe	965	Phe Phe Gly Ala Tyr	970	Phe Val Phe Glu Asn	975
Thr Val Thr Ser	980	Asn Gly Gln Ile Phe	985	Val Phe Glu Asn Thr	990
Thr Leu Val Phe	995	Thr Val Met Val Phe	1000	Val Thr Leu Lys Leu	1005
Ala Leu Asp Thr	1010	His Tyr Trp Thr Trp	1015	Val Thr Leu Lys Leu	1020
Trp Gly Ser Leu	1025	Leu Phe Tyr Val Val	1030	Asn His Phe Val Ile	1035
Gly Val Ile Trp	1040	Pro Phe Leu Asn Tyr	1045	Ser Leu Leu Trp Gly	1050
Phe Ile Gln Met	1055	Leu Ser Ser Gly Pro	1060	Arg Met Tyr Tyr Val	1065
Leu Leu Val Thr	1070	Ile Ser Leu Leu Pro	1075	Trp Leu Ala Ile Val	1080
Leu Cys Arg Gln	1085	Leu Trp Pro Thr Ala	1090	Val Leu Lys Lys Val	1095
Asn Gly Cys Ala	1100	Gln Pro Arg Asp Arg	1105	Glu Arg Val Gln Gln	1110
Leu Ala Ser Leu	1115	Gln Ser Pro Gly Tyr	1120	Ser Glu Phe Thr Pro	1125
Ala Ala Trp Tyr	1130	Ser Ser His Ser Gln	1135	Ser Thr Cys Pro Ser	1140
Trp Lys Glu Lys	1145	Val Ser Thr Glu Pro	1150	Val Thr Leu Ala Ala	1155
Ser His His His	1160	Cys Ser Ser Ile Pro	1165	Pro Ile Leu Gly Gly	1170
Ser Arg Val Gly	1175	Met Leu Val	1180	His Ser Cys Pro Arg	1185
	1190				

<210> 18
 <211> 638
 <212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474127CD1

<400> 18

Met	Gly	Lys	Ile	Glu	Asn	Asn	Glu	Arg	Val	Ile	Leu	Asn	Val	Gly
1				5					10					15
Gly	Thr	Arg	His	Glu	Thr	Tyr	Arg	Ser	Thr	Leu	Lys	Thr	Leu	Pro
				20					25					30
Gly	Thr	Arg	Leu	Ala	Leu	Leu	Ala	Ser	Ser	Glu	Pro	Pro	Gly	Asp
				35					40					45
Cys	Leu	Thr	Thr	Ala	Gly	Asp	Lys	Leu	Gln	Pro	Ser	Pro	Pro	Pro
				50					55					60
Leu	Ser	Pro	Pro	Pro	Arg	Ala	Pro	Pro	Leu	Ser	Pro	Gly	Pro	Gly
				65					70					75
Gly	Cys	Phe	Glu	Gly	Gly	Ala	Gly	Asn	Cys	Ser	Ser	Arg	Gly	Gly
				80					85					90
Arg	Ala	Ser	Asp	His	Pro	Gly	Gly	Gly	Arg	Glu	Phe	Phe	Phe	Asp
				95					100					105
Arg	His	Pro	Gly	Val	Phe	Ala	Tyr	Val	Leu	Asn	Tyr	Tyr	Arg	Thr
				110					115					120
Gly	Lys	Leu	His	Cys	Pro	Ala	Asp	Val	Cys	Gly	Pro	Leu	Phe	Glu
				125					130					135
Glu	Glu	Leu	Ala	Phe	Trp	Gly	Ile	Asp	Glu	Thr	Asp	Val	Glu	Pro
				140					145					150
Cys	Cys	Trp	Met	Thr	Tyr	Arg	Gln	His	Arg	Asp	Ala	Glu	Glu	Ala
				155					160					165
Leu	Asp	Ile	Phe	Glu	Thr	Pro	Asp	Leu	Ile	Gly	Gly	Asp	Pro	Gly
				170					175					180
Asp	Asp	Glu	Asp	Leu	Ala	Ala	Lys	Arg	Leu	Gly	Ile	Glu	Asp	Ala
				185					190					195
Ala	Gly	Leu	Gly	Gly	Pro	Asp	Gly	Lys	Ser	Gly	Arg	Trp	Arg	Arg
				200					205					210
Leu	Gln	Pro	Arg	Met	Trp	Ala	Leu	Phe	Glu	Asp	Pro	Tyr	Ser	Ser
				215					220					225
Arg	Ala	Ala	Arg	Phe	Ile	Ala	Phe	Ala	Ser	Leu	Phe	Phe	Ile	Leu
				230					235					240
Val	Ser	Ile	Thr	Thr	Phe	Cys	Leu	Glu	Thr	His	Glu	Ala	Phe	Asn
				245					250					255
Ile	Val	Lys	Asn	Lys	Thr	Glu	Pro	Val	Ile	Asn	Gly	Thr	Ser	Val
				260					265					270
Val	Leu	Gln	Tyr	Glu	Ile	Glu	Thr	Asp	Pro	Ala	Leu	Thr	Tyr	Val
				275					280					285
Glu	Gly	Val	Cys	Val	Val	Trp	Phe	Thr	Phe	Glu	Phe	Leu	Val	Arg
				290					295					300
Ile	Val	Phe	Ser	Pro	Asn	Lys	Leu	Glu	Phe	Ile	Lys	Asn	Leu	Leu
				305					310					315
Asn	Ile	Ile	Asp	Phe	Val	Ala	Ile	Leu	Pro	Phe	Tyr	Leu	Glu	Val
				320					325					330
Gly	Leu	Ser	Gly	Leu	Ser	Ser	Lys	Ala	Ala	Lys	Asp	Val	Leu	Gly
				335					340					345
Phe	Leu	Arg	Val	Val	Arg	Phe	Val	Arg	Ile	Leu	Arg	Ile	Phe	Lys
				350					355					360
Leu	Thr	Arg	His	Phe	Val	Gly	Leu	Arg	Val	Leu	Gly	His	Thr	Leu
				365					370					375
Arg	Ala	Ser	Thr	Asn	Glu	Phe	Leu	Leu	Leu	Ile	Ile	Phe	Leu	Ala
				380					385					390
Leu	Gly	Val	Leu	Ile	Phe	Ala	Thr	Met	Ile	Tyr	Tyr	Ala	Glu	Arg
				395					400					405
Val	Gly	Ala	Gln	Pro	Asn	Asp	Pro	Ser	Ala	Ser	Glu	His	Thr	Gln
				410					415					420
Phe	Lys	Asn	Ile	Pro	Ile	Gly	Phe	Trp	Trp	Ala	Val	Val	Thr	Met
				425					430					435
Thr	Thr	Leu	Gly	Tyr	Gly	Asp	Met	Tyr	Pro	Gln	Thr	Trp	Ser	Gly
				440					445					450

Met	Leu	Val	Gly	Ala	Leu	Cys	Ala	Leu	Ala	Gly	Val	Leu	Thr	Ile
				455					460					465
Ala	Met	Pro	Val	Pro	Val	Ile	Val	Asn	Asn	Phe	Gly	Met	Tyr	Tyr
				470					475					480
Ser	Leu	Ala	Met	Ala	Lys	Gln	Lys	Leu	Pro	Arg	Lys	Arg	Lys	Lys
				485					490					495
His	Ile	Pro	Pro	Ala	Pro	Gln	Ala	Ser	Ser	Pro	Thr	Phe	Cys	Lys
				500					505					510
Thr	Glu	Leu	Asn	Met	Ala	Cys	Asn	Ser	Thr	Gln	Ser	Asp	Thr	Cys
				515					520					525
Leu	Gly	Lys	Asp	Asn	Arg	Leu	Leu	Glu	His	Asn	Arg	Ser	Val	Leu
				530					535					540
Ser	Gly	Asp	Asp	Ser	Thr	Gly	Ser	Glu	Pro	Pro	Leu	Ser	Pro	Pro
				545					550					555
Glu	Arg	Leu	Pro	Ile	Arg	Arg	Ser	Ser	Thr	Arg	Asp	Lys	Asn	Arg
				560					565					570
Arg	Gly	Glu	Thr	Cys	Phe	Leu	Leu	Thr	Thr	Gly	Asp	Tyr	Thr	Cys
				575					580					585
Ala	Ser	Asp	Gly	Gly	Ile	Arg	Lys	Gly	Tyr	Glu	Lys	Ser	Arg	Ser
				590					595					600
Leu	Asn	Asn	Ile	Ala	Gly	Leu	Ala	Gly	Asn	Ala	Leu	Arg	Leu	Ser
				605					610					615
Pro	Val	Thr	Ser	Pro	Tyr	Asn	Ser	Pro	Cys	Pro	Leu	Arg	Arg	Ser
				620					625					630
Arg	Ser	Pro	Ile	Pro	Ser	Ile	Leu							
				635										

<210> 19

<211> 681

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7476949CD1

<400> 19

Met	Ser	Lys	Asp	Leu	Ala	Ala	Met	Gly	Pro	Gly	Ala	Ser	Gly	Asp
1				5					10					15
Gly	Val	Arg	Thr	Glu	Thr	Ala	Pro	His	Ile	Ala	Leu	Asp	Ser	Arg
				20					25					30
Val	Gly	Leu	His	Ala	Tyr	Asp	Ile	Ser	Val	Val	Val	Ile	Tyr	Phe
				35					40					45
Val	Phe	Val	Ile	Ala	Val	Gly	Ile	Trp	Ser	Ser	Ile	Arg	Ala	Ser
				50					55					60
Arg	Gly	Thr	Ile	Gly	Gly	Tyr	Phe	Leu	Ala	Gly	Arg	Ser	Met	Ser
				65					70					75
Trp	Trp	Pro	Ile	Gly	Ala	Ser	Leu	Met	Ser	Ser	Asn	Val	Gly	Ser
				80					85					90
Gly	Leu	Phe	Ile	Gly	Leu	Ala	Gly	Thr	Gly	Ala	Ala	Gly	Gly	Leu
				95					100					105
Ala	Val	Gly	Gly	Phe	Glu	Trp	Asn	Ala	Thr	Trp	Leu	Leu	Leu	Ala
				110					115					120
Leu	Gly	Trp	Val	Phe	Val	Pro	Val	Tyr	Ile	Ala	Ala	Gly	Val	Val
				125					130					135
Thr	Met	Pro	Gln	Tyr	Leu	Lys	Lys	Arg	Phe	Gly	Gly	Gln	Arg	Ile
				140					145					150
Gln	Val	Tyr	Met	Ser	Val	Leu	Ser	Leu	Ile	Leu	Tyr	Ile	Phe	Thr
				155					160					165
Lys	Ile	Ser	Thr	Asp	Ile	Phe	Ser	Gly	Ala	Leu	Phe	Ile	Gln	Met
				170					175					180
Ala	Leu	Gly	Trp	Asn	Leu	Tyr	Leu	Ser	Thr	Gly	Ile	Leu	Leu	Val
				185					190					195
Val	Thr	Ala	Val	Tyr	Thr	Ile	Ala	Gly	Gly	Leu	Met	Ala	Val	Ile
				200					205					210
Tyr	Thr	Asp	Ala	Leu	Gln	Thr	Val	Ile	Met	Val	Gly	Gly	Ala	Leu
				215					220					225

Val	Leu	Met	Phe	Leu	Gly	Phe	Gln	Asp	Val	Gly	Trp	Tyr	Pro	Gly
				230					235					240
Leu	Glu	Gln	Arg	Tyr	Arg	Gln	Ala	Ile	Pro	Asn	Val	Thr	Val	Pro
				245					250					255
Asn	Thr	Thr	Cys	His	Leu	Pro	Arg	Pro	Asp	Ala	Phe	His	Ile	Leu
				260					265					270
Arg	Asp	Pro	Val	Ser	Gly	Asp	Ile	Pro	Trp	Pro	Gly	Leu	Ile	Phe
				275					280					285
Gly	Leu	Thr	Val	Leu	Ala	Thr	Trp	Cys	Trp	Cys	Thr	Asp	Gln	Val
				290					295					300
Ile	Val	Gln	Arg	Ser	Leu	Ser	Ala	Lys	Ser	Leu	Ser	His	Ala	Lys
				305					310					315
Gly	Gly	Ser	Val	Leu	Gly	Gly	Tyr	Leu	Lys	Ile	Leu	Pro	Met	Phe
				320					325					330
Phe	Ile	Val	Met	Pro	Gly	Met	Ile	Ser	Arg	Ala	Leu	Phe	Pro	Asp
				335					340					345
Glu	Val	Gly	Cys	Val	Asp	Pro	Asp	Val	Cys	Gln	Arg	Ile	Cys	Gly
				350					355					360
Ala	Arg	Val	Gly	Cys	Ser	Asn	Ile	Ala	Tyr	Pro	Lys	Leu	Val	Met
				365					370					375
Ala	Leu	Met	Pro	Val	Gly	Leu	Arg	Gly	Leu	Met	Ile	Ala	Val	Ile
				380					385					390
Met	Ala	Ala	Leu	Met	Ser	Ser	Leu	Thr	Ser	Ile	Phe	Asn	Ser	Ser
				395					400					405
Ser	Thr	Leu	Phe	Thr	Ile	Asp	Val	Trp	Gln	Arg	Phe	Arg	Arg	Lys
				410					415					420
Ser	Thr	Glu	Gln	Glu	Leu	Met	Val	Val	Gly	Arg	Val	Phe	Val	Val
				425					430					435
Phe	Leu	Val	Val	Ile	Ser	Ile	Leu	Trp	Ile	Pro	Ile	Ile	Gln	Ser
				440					445					450
Ser	Asn	Ser	Gly	Gln	Leu	Phe	Asp	Tyr	Ile	Gln	Ala	Val	Thr	Ser
				455					460					465
Tyr	Leu	Ala	Pro	Pro	Ile	Thr	Ala	Leu	Phe	Leu	Leu	Ala	Ile	Phe
				470					475					480
Cys	Lys	Arg	Val	Thr	Glu	Pro	Gly	Ala	Phe	Trp	Gly	Leu	Val	Phe
				485					490					495
Gly	Leu	Gly	Val	Gly	Leu	Leu	Arg	Met	Ile	Leu	Glu	Phe	Ser	Tyr
				500					505					510
Pro	Ala	Pro	Ala	Cys	Gly	Glu	Val	Asp	Arg	Arg	Pro	Ala	Val	Leu
				515					520					525
Lys	Asp	Phe	His	Tyr	Leu	Tyr	Phe	Ala	Ile	Leu	Leu	Cys	Gly	Leu
				530					535					540
Thr	Ala	Ile	Val	Ile	Val	Ile	Val	Ser	Leu	Cys	Thr	Thr	Pro	Ile
				545					550					555
Pro	Glu	Glu	Gln	Leu	Thr	Arg	Leu	Thr	Trp	Trp	Thr	Arg	Asn	Cys
				560					565					570
Pro	Leu	Ser	Glu	Leu	Glu	Lys	Glu	Ala	His	Glu	Ser	Thr	Pro	Glu
				575					580					585
Ile	Ser	Glu	Arg	Pro	Ala	Gly	Glu	Cys	Pro	Ala	Gly	Gly	Gly	Ala
				590					595					600
Ala	Glu	Asn	Ser	Ser	Leu	Gly	Gln	Glu	Gln	Pro	Glu	Ala	Pro	Ser
				605					610					615
Arg	Ser	Trp	Gly	Lys	Leu	Leu	Trp	Ser	Trp	Phe	Cys	Gly	Leu	Ser
				620					625					630
Gly	Thr	Pro	Glu	Gln	Ala	Leu	Ser	Pro	Ala	Glu	Lys	Ala	Ala	Leu
				635					640					645
Glu	Gln	Lys	Leu	Thr	Ser	Ile	Glu	Glu	Glu	Pro	Leu	Trp	Arg	His
				650					655					660
Val	Cys	Asn	Ile	Asn	Ala	Val	Leu	Leu	Leu	Ala	Ile	Asn	Ile	Phe
				665					670					675
Leu	Trp	Gly	Tyr	Phe	Ala									
				680										

<210> 20
 <211> 1096
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477249CD1

<400> 20

Met	Trp	Arg	Trp	Ile	Arg	Gln	Gln	Leu	Gly	Phe	Asp	Pro	Pro	His
1				5					10					15
Gln	Ser	Asp	Thr	Arg	Thr	Ile	Tyr	Val	Ala	Asn	Arg	Phe	Pro	Gln
				20					25					30
Asn	Gly	Leu	Tyr	Thr	Pro	Gln	Lys	Phe	Ile	Asp	Asn	Arg	Ile	Ile
				35					40					45
Ser	Ser	Lys	Tyr	Thr	Val	Trp	Asn	Phe	Val	Pro	Lys	Asn	Leu	Phe
				50					55					60
Glu	Gln	Phe	Arg	Arg	Val	Ala	Asn	Phe	Tyr	Phe	Leu	Ile	Ile	Phe
				65					70					75
Leu	Val	Gln	Leu	Met	Ile	Asp	Thr	Pro	Thr	Ser	Pro	Val	Thr	Ser
				80					85					90
Gly	Leu	Pro	Leu	Phe	Phe	Val	Ile	Thr	Val	Thr	Ala	Ile	Lys	Gln
				95					100					105
Gly	Tyr	Glu	Asp	Trp	Leu	Arg	His	Asn	Ser	Asp	Asn	Glu	Val	Asn
				110					115					120
Gly	Ala	Pro	Val	Tyr	Val	Val	Arg	Ser	Gly	Gly	Leu	Val	Lys	Thr
				125					130					135
Arg	Ser	Lys	Asn	Ile	Arg	Val	Gly	Asp	Ile	Val	Arg	Ile	Ala	Lys
				140					145					150
Asp	Glu	Ile	Phe	Pro	Ala	Asp	Leu	Val	Leu	Leu	Ser	Ser	Asp	Arg
				155					160					165
Leu	Asp	Gly	Ser	Cys	His	Val	Thr	Thr	Ala	Ser	Leu	Asp	Gly	Glu
				170					175					180
Thr	Asn	Leu	Lys	Thr	His	Val	Ala	Val	Pro	Glu	Thr	Ala	Leu	Leu
				185					190					195
Gln	Thr	Val	Ala	Asn	Leu	Asp	Thr	Leu	Val	Ala	Val	Ile	Glu	Cys
				200					205					210
Gln	Gln	Pro	Glu	Ala	Asp	Leu	Tyr	Arg	Phe	Met	Gly	Arg	Met	Ile
				215					220					225
Ile	Thr	Gln	Gln	Met	Glu	Glu	Ile	Val	Arg	Pro	Leu	Gly	Pro	Glu
				230					235					240
Ser	Leu	Leu	Leu	Arg	Gly	Ala	Arg	Leu	Lys	Asn	Thr	Lys	Glu	Ile
				245					250					255
Phe	Gly	Val	Ala	Val	Tyr	Thr	Gly	Met	Glu	Thr	Lys	Met	Ala	Leu
				260					265					270
Asn	Tyr	Lys	Ser	Lys	Ser	Gln	Lys	Arg	Ser	Ala	Val	Glu	Lys	Ser
				275					280					285
Met	Asn	Thr	Phe	Leu	Ile	Ile	Tyr	Leu	Val	Ile	Leu	Ile	Ser	Glu
				290					295					300
Ala	Val	Ile	Ser	Thr	Ile	Leu	Lys	Tyr	Thr	Trp	Gln	Ala	Glu	Glu
				305					310					315
Lys	Trp	Asp	Glu	Pro	Trp	Tyr	Asn	Gln	Lys	Thr	Glu	His	Gln	Arg
				320					325					330
Asn	Ser	Ser	Lys	Val	Glu	Tyr	Val	Phe	Thr	Asp	Lys	Thr	Gly	Thr
				335					340					345
Leu	Thr	Glu	Asn	Glu	Met	Gln	Phe	Arg	Glu	Cys	Ser	Ile	Asn	Gly
				350					355					360
Met	Lys	Tyr	Gln	Glu	Ile	Asn	Gly	Arg	Leu	Val	Pro	Glu	Gly	Pro
				365					370					375
Thr	Pro	Asp	Ser	Ser	Glu	Gly	Asn	Leu	Ser	Tyr	Leu	Ser	Ser	Leu
				380					385					390
Ser	His	Leu	Asn	Asn	Leu	Ser	His	Leu	Thr	Thr	Ser	Ser	Ser	Phe
				395					400					405
Arg	Thr	Ser	Pro	Glu	Asn	Glu	Thr	Glu	Leu	Ile	Lys	Glu	His	Asp
				410					415					420
Leu	Phe	Phe	Lys	Ala	Val	Ser	Leu	Cys	His	Thr	Val	Gln	Ile	Ser
				425					430					435
Asn	Val	Gln	Thr	Asp	Cys	Thr	Gly	Asp	Gly	Pro	Trp	Gln	Ser	Asn
				440					445					450
Leu	Ala	Pro	Ser	Gln	Leu	Glu	Tyr	Tyr	Ala	Ser	Ser	Pro	Asp	Glu
				455					460					465

Lys	Ala	Leu	Val	Glu	Ala	Ala	Ala	Arg	Ile	Gly	Ile	Val	Phe	Ile
				470					475					480
Gly	Asn	Ser	Glu	Glu	Thr	Met	Glu	Val	Lys	Thr	Leu	Gly	Lys	Leu
				485					490					495
Glu	Arg	Tyr	Lys	Leu	Leu	His	Ile	Leu	Glu	Phe	Asp	Ser	Asp	Arg
				500					505					510
Arg	Arg	Met	Ser	Val	Ile	Val	Gln	Ala	Pro	Ser	Gly	Glu	Lys	Leu
				515					520					525
Leu	Phe	Ala	Lys	Gly	Ala	Glu	Ser	Ser	Ile	Leu	Pro	Lys	Cys	Ile
				530					535					540
Gly	Gly	Glu	Ile	Glu	Lys	Thr	Arg	Ile	His	Val	Asp	Glu	Phe	Ala
				545					550					555
Leu	Lys	Gly	Leu	Arg	Thr	Leu	Cys	Ile	Ala	Tyr	Arg	Lys	Phe	Thr
				560					565					570
Ser	Lys	Glu	Tyr	Glu	Glu	Ile	Asp	Lys	Arg	Ile	Phe	Glu	Ala	Arg
				575					580					585
Thr	Ala	Leu	Gln	Gln	Arg	Glu	Glu	Lys	Leu	Ala	Ala	Val	Phe	Gln
				590					595					600
Phe	Ile	Glu	Lys	Asp	Leu	Ile	Leu	Leu	Gly	Ala	Thr	Ala	Val	Glu
				605					610					615
Asp	Arg	Leu	Gln	Asp	Lys	Val	Arg	Glu	Thr	Ile	Glu	Ala	Leu	Arg
				620					625					630
Met	Ala	Gly	Ile	Lys	Val	Trp	Val	Leu	Thr	Gly	Asp	Lys	His	Glu
				635					640					645
Thr	Ala	Val	Ser	Val	Ser	Leu	Ser	Cys	Gly	His	Phe	His	Arg	Thr
				650					655					660
Met	Asn	Ile	Leu	Glu	Leu	Ile	Asn	Gln	Lys	Ser	Asp	Ser	Glu	Cys
				665					670					675
Ala	Glu	Gln	Leu	Arg	Gln	Leu	Ala	Arg	Arg	Ile	Thr	Glu	Asp	His
				680					685					690
Val	Ile	Gln	His	Gly	Leu	Val	Val	Asp	Gly	Thr	Ser	Leu	Ser	Leu
				695					700					705
Ala	Leu	Arg	Glu	His	Glu	Lys	Leu	Phe	Met	Glu	Val	Cys	Arg	Asn
				710					715					720
Cys	Ser	Ala	Val	Leu	Cys	Cys	Arg	Met	Ala	Pro	Leu	Gln	Lys	Ala
				725					730					735
Lys	Val	Ile	Arg	Leu	Ile	Lys	Ile	Ser	Pro	Glu	Lys	Pro	Ile	Thr
				740					745					750
Leu	Ala	Val	Gly	Asp	Gly	Ala	Asn	Asp	Val	Ser	Met	Ile	Gln	Glu
				755					760					765
Ala	His	Val	Gly	Ile	Gly	Ile	Met	Gly	Lys	Glu	Gly	Arg	Gln	Ala
				770					775					780
Ala	Arg	Asn	Ser	Asp	Tyr	Ala	Ile	Ala	Arg	Phe	Lys	Phe	Leu	Ser
				785					790					795
Lys	Leu	Leu	Phe	Val	His	Gly	His	Phe	Tyr	Tyr	Ile	Arg	Ile	Ala
				800					805					810
Thr	Leu	Val	Gln	Tyr	Phe	Phe	Tyr	Lys	Asn	Val	Cys	Phe	Ile	Thr
				815					820					825
Pro	Gln	Phe	Leu	Tyr	Gln	Phe	Tyr	Cys	Leu	Phe	Ser	Gln	Gln	Thr
				830					835					840
Leu	Tyr	Asp	Ser	Val	Tyr	Leu	Thr	Leu	Tyr	Asn	Ile	Cys	Phe	Thr
				845					850					855
Ser	Leu	Pro	Ile	Leu	Ile	Tyr	Ser	Leu	Leu	Glu	Gln	His	Val	Asp
				860					865					870
Pro	His	Val	Leu	Gln	Asn	Lys	Pro	Thr	Leu	Tyr	Arg	Asp	Ile	Ser
				875					880					885
Lys	Asn	Arg	Leu	Leu	Ser	Ile	Lys	Thr	Phe	Leu	Tyr	Trp	Thr	Ile
				890					895					900
Leu	Gly	Phe	Ser	His	Ala	Phe	Ile	Phe	Phe	Phe	Gly	Ser	Tyr	Leu
				905					910					915
Leu	Ile	Gly	Lys	Asp	Thr	Ser	Leu	Leu	Gly	Asn	Gly	Gln	Met	Phe
				920					925					930
Gly	Asn	Trp	Thr	Phe	Gly	Thr	Leu	Val	Phe	Thr	Val	Met	Val	Ile
				935					940					945
Thr	Val	Thr	Val	Lys	Met	Ala	Leu	Glu	Thr	His	Phe	Trp	Thr	Trp
				950					955					960
Ile	Asn	His	Leu	Val	Thr	Trp	Gly	Ser	Ile	Ile	Phe	Tyr	Phe	Val

Phe	Ser	Leu	Phe	965	Gly	Gly	Ile	Leu	970	Pro	Phe	Leu	Gly	Ser	975
			Tyr	980	Val	Phe	Ile	Gln	985	Leu	Ser	Ser	Gly	Ser	990
Gln	Asn	Met	Tyr	995					1000						1005
Ala	Trp	Phe	Ala	1010	Ile	Leu	Met	Val	1015	Thr	Cys	Leu	Phe	Leu	1020
Asp	Ile	Ile	Lys	1025	Val	Phe	Asp	Arg	1030	Leu	His	Pro	Thr	Ser	1035
Thr	Glu	Lys	Ala	1040	Leu	Thr	Glu	Thr	1045	Ala	Gly	Ile	Lys	Cys	1050
Leu	Asp	Ser	Met	1055	Cys	Phe	Pro	Glu	1060	Glu	Ala	Ala	Cys	Ala	1065
Ser	Val	Gly	Arg	1070	Leu	Glu	Arg	Val	1075	Gly	Arg	Cys	Ser	Pro	1080
Thr	His	Ile	Ser	1085	Cys	Glu	Ile	Ser	1090	Ser	Ser	Leu	Cys	Cys	1095

Arg

<210> 21
 <211> 707
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7477720CD1

<400> 21

Met	Ala	Leu	Gln	Met	Phe	Val	Thr	Tyr	Ser	Pro	Trp	Asn	Cys	Leu	
1				5					10					15	
Leu	Leu	Leu	Val	Ala	Leu	Glu	Cys	Ser	Glu	Ala	Ser	Ser	Asp	Leu	
				20					25					30	
Asn	Glu	Ser	Ala	Asn	Ser	Thr	Ala	Gln	Tyr	Ala	Ser	Asn	Ala	Trp	
				35					40					45	
Phe	Ala	Ala	Ala	Ser	Ser	Glu	Pro	Glu	Glu	Gly	Ile	Ser	Val	Phe	
				50					55					60	
Glu	Leu	Asp	Tyr	Asp	Tyr	Val	Gln	Ile	Pro	Tyr	Glu	Val	Thr	Leu	
				65					70					75	
Trp	Ile	Leu	Leu	Ala	Ser	Leu	Ala	Lys	Ile	Gly	Phe	His	Leu	Tyr	
				80					85					90	
His	Arg	Leu	Pro	Gly	Leu	Met	Pro	Glu	Ser	Cys	Leu	Leu	Ile	Leu	
				95					100					105	
Val	Gly	Ala	Leu	Val	Gly	Gly	Ile	Ile	Phe	Gly	Thr	Asp	His	Lys	
				110					115					120	
Ser	Pro	Pro	Val	Met	Asp	Ser	Ser	Ile	Tyr	Phe	Leu	Tyr	Leu	Leu	
				125					130					135	
Pro	Pro	Ile	Val	Leu	Glu	Gly	Gly	Tyr	Phe	Met	Pro	Thr	Arg	Pro	
				140					145					150	
Phe	Phe	Glu	Asn	Ile	Gly	Ser	Ile	Leu	Trp	Trp	Ala	Val	Leu	Gly	
				155					160					165	
Ala	Leu	Ile	Asn	Ala	Leu	Gly	Ile	Gly	Leu	Ser	Leu	Tyr	Leu	Ile	
				170					175					180	
Cys	Gln	Val	Lys	Ala	Phe	Gly	Leu	Gly	Asp	Val	Asn	Leu	Leu	Gln	
				185					190					195	
Asn	Leu	Leu	Phe	Gly	Ser	Leu	Ile	Ser	Ala	Val	Asp	Pro	Val	Ala	
				200					205					210	
Val	Leu	Ala	Val	Phe	Glu	Glu	Ala	Arg	Val	Asn	Glu	Gln	Leu	Tyr	
				215					220					225	
Met	Met	Ile	Phe	Gly	Glu	Ala	Leu	Leu	Asn	Asp	Gly	Ile	Thr	Val	
				230					235					240	
Val	Leu	Tyr	Asn	Met	Leu	Ile	Ala	Phe	Thr	Lys	Met	His	Lys	Phe	
				245					250					255	
Glu	Asp	Ile	Glu	Thr	Val	Asp	Ile	Leu	Ala	Gly	Cys	Ala	Arg	Phe	
				260					265					270	
Ile	Val	Val	Gly	Leu	Gly	Gly	Val	Leu	Phe	Gly	Ile	Val	Phe	Gly	

Phe	Ile	Ser	Ala	275	Phe	Ile	Thr	Arg	Phe	280	Thr	Gln	Asn	Ile	Ser	Ala	285
Ile	Glu	Pro	Leu	290	Ile	Val	Phe	Met	Phe	295	Ser	Tyr	Leu	Ser	Tyr	Leu	300
Ala	Ala	Glu	Thr	305	Leu	Tyr	Leu	Ser	Gly	310	Ile	Leu	Ala	Ile	Thr	Ala	315
Cys	Ala	Val	Thr	320	Met	Lys	Lys	Tyr	Val	325	Glu	Glu	Asn	Val	Ser	Gln	330
Thr	Ser	Tyr	Thr	335	Thr	Ile	Lys	Tyr	Phe	340	Met	Lys	Met	Leu	Ser	Ser	345
Val	Ser	Glu	Thr	350	Leu	Ile	Phe	Ile	Phe	355	Met	Gly	Val	Ser	Thr	Val	360
Gly	Lys	Asn	His	365	Glu	Trp	Asn	Trp	Ala	370	Phe	Ile	Cys	Phe	Thr	Leu	375
Ala	Phe	Cys	Gln	380	Ile	Trp	Arg	Ala	Ile	385	Ser	Val	Phe	Ala	Leu	Phe	390
Tyr	Ile	Ser	Asn	395	Gln	Phe	Arg	Thr	Phe	400	Pro	Phe	Ser	Ile	Lys	Asp	405
Gln	Cys	Ile	Ile	410	Phe	Tyr	Ser	Gly	Val	415	Arg	Gly	Ala	Gly	Ser	Phe	420
Ser	Leu	Ala	Phe	425	Leu	Leu	Pro	Leu	Ser	430	Leu	Phe	Pro	Arg	Lys	Lys	435
Met	Phe	Val	Thr	440	Ala	Thr	Leu	Val	Val	445	Ile	Tyr	Phe	Thr	Val	Phe	450
Ile	Gln	Gly	Ile	455	Thr	Val	Gly	Pro	Leu	460	Val	Arg	Tyr	Leu	Asp	Val	465
Lys	Lys	Thr	Asn	470	Lys	Lys	Glu	Ser	Ile	475	Asn	Glu	Glu	Leu	His	Ile	480
Arg	Leu	Met	Asp	485	His	Leu	Lys	Ala	Gly	490	Ile	Glu	Asp	Val	Cys	Gly	495
His	Trp	Ser	His	500	Tyr	Gln	Val	Arg	Asp	505	Lys	Phe	Lys	Lys	Phe	Asp	510
His	Arg	Tyr	Leu	515	Arg	Lys	Ile	Leu	Ile	520	Arg	Lys	Asn	Leu	Pro	Lys	525
Ser	Ser	Ile	Val	530	Ser	Leu	Tyr	Lys	Lys	535	Leu	Glu	Met	Lys	Gln	Ala	540
Ile	Glu	Met	Val	545	Glu	Thr	Gly	Ile	Leu	550	Ser	Ser	Thr	Ala	Phe	Ser	555
Ile	Pro	His	Gln	560	Ala	Gln	Arg	Ile	Gln	565	Gly	Ile	Lys	Arg	Leu	Ser	570
Pro	Glu	Asp	Val	575	Glu	Ser	Ile	Arg	Asp	580	Ile	Leu	Thr	Ser	Asn	Met	585
Tyr	Gln	Val	Arg	590	Gln	Arg	Thr	Leu	Ser	595	Tyr	Asn	Lys	Tyr	Asn	Leu	600
Lys	Pro	Gln	Thr	605	Ser	Glu	Lys	Gln	Ala	610	Lys	Glu	Ile	Leu	Ile	Arg	615
Arg	Gln	Asn	Thr	620	Leu	Arg	Glu	Ser	Met	625	Arg	Lys	Gly	His	Ser	Leu	630
Pro	Trp	Gly	Lys	635	Pro	Ala	Gly	Thr	Lys	640	Asn	Ile	Arg	Tyr	Leu	Ser	645
Tyr	Pro	Tyr	Gly	650	Asn	Pro	Gln	Ser	Ala	655	Gly	Arg	Asp	Thr	Arg	Ala	660
Ala	Gly	Phe	Ser	665	Gly	Lys	Leu	Pro	Thr	670	Trp	Leu	Leu	Cys	Cys	Phe	675
Ser	Val	Glu	Ser	680	Gly	Gly	Lys	Tyr	Leu	685	Gly	Val	Trp	Ala	Lys	Arg	690
Gln	His			695						700							705

<210> 22
 <211> 729
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID No: 7477852CD1

<400> 22

Met	Gly	Gly	Phe	Leu	Pro	Lys	Ala	Glu	Gly	Pro	Gly	Ser	Gln	Leu
1				5					10					15
Gln	Lys	Leu	Leu	Pro	Ser	Phe	Leu	Val	Arg	Glu	Gln	Asp	Trp	Asp
				20					25					30
Gln	His	Leu	Asp	Lys	Leu	His	Met	Leu	Gln	Gln	Lys	Arg	Ile	Leu
				35					40					45
Glu	Ser	Pro	Leu	Leu	Arg	Ala	Ser	Lys	Glu	Asn	Asp	Leu	Ser	Val
				50					55					60
Leu	Arg	Gln	Leu	Leu	Leu	Asp	Cys	Thr	Cys	Asp	Val	Arg	Gln	Arg
				65					70					75
Gly	Ala	Leu	Gly	Glu	Thr	Ala	Leu	His	Ile	Ala	Ala	Leu	Tyr	Asp
				80					85					90
Asn	Leu	Glu	Ala	Ala	Leu	Val	Leu	Met	Glu	Ala	Ala	Pro	Glu	Leu
				95					100					105
Val	Phe	Glu	Pro	Thr	Thr	Cys	Glu	Ala	Phe	Ala	Gly	Gln	Thr	Ala
				110					115					120
Leu	His	Ile	Ala	Val	Val	Asn	Gln	Asn	Val	Asn	Leu	Val	Arg	Ala
				125					130					135
Leu	Leu	Thr	Arg	Arg	Ala	Ser	Val	Ser	Ala	Arg	Ala	Thr	Gly	Thr
				140					145					150
Ala	Phe	Arg	Arg	Ser	Pro	Arg	Asn	Leu	Ile	Tyr	Phe	Gly	Glu	His
				155					160					165
Pro	Leu	Ser	Phe	Ala	Ala	Cys	Val	Asn	Ser	Glu	Glu	Ile	Val	Arg
				170					175					180
Leu	Leu	Ile	Glu	His	Gly	Ala	Asp	Ile	Arg	Ala	Gln	Asp	Ser	Leu
				185					190					195
Gly	Asn	Thr	Val	Leu	His	Ile	Leu	Ile	Leu	Gln	Pro	Asn	Lys	Thr
				200					205					210
Phe	Ala	Cys	Gln	Met	Tyr	Asn	Leu	Leu	Leu	Ser	Tyr	Asp	Gly	His
				215					220					225
Gly	Asp	His	Leu	Gln	Pro	Leu	Asp	Leu	Val	Pro	Asn	His	Gln	Gly
				230					235					240
Leu	Thr	Pro	Phe	Lys	Leu	Ala	Gly	Val	Glu	Gly	Asn	Thr	Val	Met
				245					250					255
Phe	Gln	His	Leu	Met	Gln	Lys	Arg	Arg	His	Ile	Gln	Trp	Thr	Tyr
				260					265					270
Gly	Pro	Leu	Thr	Ser	Ile	Leu	Tyr	Asp	Leu	Thr	Glu	Ile	Asp	Ser
				275					280					285
Trp	Gly	Glu	Glu	Leu	Ser	Phe	Leu	Glu	Leu	Val	Val	Ser	Ser	Asp
				290					295					300
Lys	Arg	Glu	Ala	Arg	Gln	Ile	Leu	Glu	Gln	Thr	Pro	Val	Lys	Glu
				305					310					315
Leu	Val	Ser	Phe	Lys	Trp	Asn	Lys	Tyr	Gly	Arg	Pro	Tyr	Phe	Cys
				320					325					330
Ile	Leu	Ala	Ala	Leu	Tyr	Leu	Leu	Tyr	Met	Ile	Cys	Phe	Thr	Thr
				335					340					345
Cys	Cys	Val	Tyr	Arg	Pro	Leu	Lys	Phe	Arg	Gly	Gly	Asn	Arg	Thr
				350					355					360
His	Ser	Arg	Asp	Ile	Thr	Ile	Leu	Gln	Gln	Lys	Leu	Leu	Gln	Glu
				365					370					375
Ala	Tyr	Glu	Thr	Arg	Glu	Asp	Ile	Ile	Arg	Leu	Val	Gly	Glu	Leu
				380					385					390
Val	Ser	Ile	Val	Gly	Ala	Val	Ile	Ile	Leu	Leu	Leu	Glu	Ile	Pro
				395					400					405
Asp	Ile	Phe	Arg	Val	Gly	Ala	Ser	Arg	Tyr	Phe	Gly	Lys	Thr	Ile
				410					415					420
Leu	Gly	Gly	Pro	Phe	His	Val	Ile	Met	Ile	Thr	Tyr	Ala	Ser	Leu
				425					430					435
Val	Leu	Val	Thr	Met	Val	Met	Arg	Leu	Thr	Asn	Thr	Asn	Gly	Glu
				440					445					450
Val	Val	Pro	Met	Ser	Phe	Ala	Leu	Val	Leu	Gly	Trp	Cys	Ser	Val
				455					460					465
Met	Tyr	Phe	Thr	Arg	Gly	Phe	Gln	Met	Leu	Gly	Pro	Phe	Thr	Ile
				470					475					480

Met	Ile	Gln	Lys	Met	Ile	Phe	Gly	Asp	Leu	Met	Arg	Phe	Cys	Trp
				485					490					495
Leu	Met	Ala	Val	Val	Ile	Leu	Gly	Phe	Ala	Ser	Ala	Phe	Tyr	Ile
				500					505					510
Ile	Phe	Gln	Thr	Glu	Asp	Pro	Thr	Ser	Leu	Gly	Gln	Phe	Tyr	Asp
				515					520					525
Tyr	Pro	Met	Ala	Leu	Phe	Thr	Thr	Phe	Glu	Leu	Phe	Leu	Thr	Val
				530					535					540
Ile	Asp	Ala	Pro	Ala	Asn	Tyr	Asp	Val	Asp	Leu	Pro	Phe	Met	Phe
				545					550					555
Ser	Ile	Val	Asn	Phe	Ala	Phe	Ala	Ile	Ile	Ala	Thr	Leu	Leu	Met
				560					565					570
Leu	Asn	Leu	Phe	Ile	Ala	Met	Met	Gly	Asp	Thr	His	Trp	Arg	Val
				575					580					585
Ala	Gln	Glu	Arg	Asp	Glu	Leu	Trp	Arg	Ala	Gln	Val	Val	Ala	Thr
				590					595					600
Thr	Val	Met	Leu	Glu	Arg	Lys	Leu	Pro	Arg	Cys	Leu	Trp	Pro	Arg
				605					610					615
Ser	Gly	Ile	Cys	Gly	Cys	Glu	Phe	Gly	Leu	Gly	Asp	Arg	Trp	Phe
				620					625					630
Leu	Arg	Val	Glu	Asn	His	Asn	Asp	Gln	Asn	Pro	Leu	Arg	Val	Leu
				635					640					645
Arg	Tyr	Val	Glu	Val	Phe	Lys	Asn	Ser	Asp	Lys	Glu	Asp	Asp	Gln
				650					655					660
Glu	His	Pro	Ser	Glu	Lys	Gln	Pro	Ser	Gly	Ala	Glu	Ser	Gly	Thr
				665					670					675
Leu	Ala	Arg	Ala	Ser	Leu	Ala	Leu	Pro	Thr	Ser	Ser	Leu	Ser	Arg
				680					685					690
Thr	Ala	Ser	Gln	Ser	Ser	Ser	His	Arg	Gly	Trp	Glu	Ile	Leu	Arg
				695					700					705
Gln	Asn	Thr	Leu	Gly	His	Leu	Asn	Leu	Gly	Leu	Asn	Leu	Ser	Glu
				710					715					720
Gly	Asp	Gly	Glu	Glu	Val	Tyr	His	Phe						
				725										

<210> 23
 <211> 492
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1471717CD1

<400> 23														
Met	Ala	Thr	Lys	Pro	Thr	Glu	Pro	Val	Thr	Ile	Leu	Ser	Leu	Arg
1				5					10					15
Lys	Leu	Ser	Leu	Gly	Thr	Ala	Glu	Pro	Gln	Val	Lys	Glu	Pro	Lys
				20					25					30
Thr	Phe	Thr	Val	Glu	Asp	Ala	Val	Glu	Thr	Ile	Gly	Phe	Gly	Arg
				35					40					45
Phe	His	Ile	Ala	Leu	Phe	Leu	Ile	Met	Gly	Ser	Thr	Gly	Val	Val
				50					55					60
Glu	Ala	Met	Glu	Ile	Met	Leu	Ile	Ala	Val	Val	Ser	Pro	Val	Ile
				65					70					75
Arg	Cys	Glu	Trp	Gln	Leu	Glu	Asn	Trp	Gln	Val	Ala	Leu	Val	Thr
				80					85					90
Thr	Met	Val	Phe	Phe	Gly	Tyr	Met	Val	Phe	Ser	Ile	Leu	Phe	Gly
				95					100					105
Leu	Leu	Ala	Asp	Arg	Tyr	Gly	Arg	Trp	Lys	Ile	Leu	Leu	Ile	Ser
				110					115					120
Phe	Leu	Trp	Gly	Ala	Tyr	Phe	Ser	Leu	Leu	Thr	Ser	Phe	Ala	Pro
				125					130					135
Ser	Tyr	Ile	Trp	Phe	Val	Phe	Leu	Arg	Thr	Met	Val	Gly	Cys	Gly
				140					145					150
Val	Ser	Gly	His	Ser	Gln	Gly	Leu	Ile	Ile	Lys	Thr	Glu	Phe	Leu
				155					160					165

Pro	Thr	Lys	Tyr	Arg	Gly	Tyr	Met	Leu	Pro	Leu	Ser	Gln	Val	Phe	
				170					175					180	
Trp	Leu	Ala	Gly	Ser	Leu	Leu	Ile	Ile	Gly	Leu	Ala	Ser	Val	Ile	
				185					190					195	
Ile	Pro	Thr	Ile	Gly	Trp	Arg	Trp	Leu	Ile	Arg	Val	Ala	Ser	Ile	
				200					205					210	
Pro	Gly	Ile	Ile	Leu	Ile	Val	Ala	Phe	Lys	Phe	Ile	Pro	Glu	Ser	
				215					220					225	
Ala	Arg	Phe	Asn	Val	Ser	Thr	Gly	Asn	Thr	Arg	Ala	Ala	Leu	Ala	
				230					235					240	
Thr	Leu	Glu	Arg	Val	Ala	Lys	Met	Asn	Arg	Ser	Val	Met	Pro	Glu	
				245					250					255	
Gly	Lys	Leu	Val	Glu	Pro	Val	Leu	Glu	Lys	Arg	Gly	Arg	Phe	Ala	
				260					265					270	
Asp	Leu	Leu	Asp	Ala	Lys	Tyr	Leu	Arg	Thr	Thr	Leu	Gln	Ile	Trp	
				275					280					285	
Val	Ile	Trp	Leu	Gly	Ile	Ser	Phe	Ala	Tyr	Tyr	Gly	Val	Ile	Leu	
				290					295					300	
Ala	Ser	Ala	Glu	Leu	Leu	Glu	Arg	Asp	Leu	Val	Cys	Gly	Ser	Lys	
				305					310					315	
Ser	Asp	Ser	Ala	Val	Val	Val	Thr	Gly	Gly	Asp	Ser	Gly	Glu	Ser	
				320					325					330	
Gln	Ser	Pro	Cys	Tyr	Cys	His	Met	Phe	Ala	Pro	Ser	Asp	Tyr	Arg	
				335					340					345	
Thr	Met	Ile	Ile	Ser	Thr	Ile	Gly	Glu	Ile	Ala	Leu	Asn	Pro	Leu	
				350					355					360	
Asn	Ile	Leu	Gly	Ile	Asn	Phe	Leu	Gly	Arg	Arg	Leu	Ser	Leu	Ser	
				365					370					375	
Ile	Thr	Met	Gly	Cys	Thr	Ala	Leu	Phe	Cys	Leu	Leu	Leu	Asn	Ile	
				380					385					390	
Cys	Thr	Ser	Ser	Ala	Gly	Leu	Ile	Gly	Phe	Leu	Phe	Met	Leu	Arg	
				395					400					405	
Ala	Leu	Val	Ala	Ala	Asn	Phe	Asn	Thr	Val	Tyr	Ile	Tyr	Thr	Ala	
				410					415					420	
Glu	Val	Tyr	Pro	Thr	Thr	Met	Arg	Ala	Leu	Gly	Met	Gly	Thr	Ser	
				425					430					435	
Gly	Ser	Leu	Cys	Arg	Ile	Gly	Ala	Met	Val	Ala	Pro	Phe	Ile	Ser	
				440					445					450	
Gln	Val	Leu	Met	Ser	Ala	Ser	Ile	Leu	Gly	Ala	Leu	Cys	Leu	Phe	
				455					460					465	
Ser	Ser	Val	Cys	Val	Val	Cys	Ala	Ile	Ser	Ala	Phe	Thr	Leu	Pro	
				470					475					480	
Ile	Glu	Thr	Lys	Gly	Arg	Ala	Leu	Gln	Gln	Ile	Lys				
				485					490						

<210> 24

<211> 1494

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3874406CD1

<400> 24

Met	Asn	Met	Lys	Gln	Lys	Ser	Val	Tyr	Gln	Gln	Thr	Lys	Ala	Leu	
1				5					10					15	
Leu	Cys	Lys	Asn	Phe	Leu	Lys	Lys	Trp	Arg	Met	Lys	Arg	Glu	Ser	
				20					25					30	
Leu	Leu	Glu	Trp	Gly	Leu	Ser	Ile	Leu	Leu	Gly	Leu	Cys	Ile	Ala	
				35					40					45	
Leu	Phe	Ser	Ser	Ser	Met	Arg	Asn	Val	Gln	Phe	Pro	Gly	Met	Ala	
				50					55					60	
Pro	Gln	Asn	Leu	Gly	Arg	Val	Asp	Lys	Phe	Asn	Ser	Ser	Ser	Leu	
				65					70					75	
Met	Val	Val	Tyr	Thr	Pro	Ile	Ser	Asn	Leu	Thr	Gln	Gln	Ile	Met	
				80					85					90	

Asn	Lys	Thr	Ala	Leu	Ala	Pro	Leu	Leu	Lys	Gly	Thr	Ser	Val	Ile
				95					100					105
Gly	Ala	Pro	Asn	Lys	Thr	His	Met	Asp	Glu	Ile	Leu	Leu	Glu	Asn
				110					115					120
Leu	Pro	Tyr	Ala	Met	Gly	Ile	Ile	Phe	Asn	Glu	Thr	Phe	Ser	Tyr
				125					130					135
Lys	Leu	Ile	Phe	Phe	Gln	Gly	Tyr	Asn	Ser	Pro	Leu	Trp	Lys	Glu
				140					145					150
Asp	Phe	Ser	Ala	His	Cys	Trp	Asp	Gly	Tyr	Gly	Glu	Phe	Ser	Cys
				155					160					165
Thr	Leu	Thr	Lys	Tyr	Trp	Asn	Arg	Gly	Phe	Val	Ala	Leu	Gln	Thr
				170					175					180
Ala	Ile	Asn	Thr	Ala	Ile	Ile	Glu	Val	Ala	Leu	Val	Phe	Leu	Met
				185					190					195
Ser	Val	Leu	Leu	Lys	Lys	Ala	Val	Leu	Thr	Asn	Leu	Val	Val	Phe
				200					205					210
Leu	Leu	Thr	Leu	Phe	Trp	Gly	Cys	Leu	Gly	Phe	Thr	Val	Phe	Tyr
				215					220					225
Glu	Gln	Leu	Pro	Ser	Ser	Leu	Glu	Trp	Ile	Leu	Asn	Ile	Cys	Ser
				230					235					240
Pro	Phe	Ala	Phe	Thr	Thr	Gly	Met	Ile	Gln	Ile	Ile	Lys	Leu	Asp
				245					250					255
Tyr	Asn	Leu	Asn	Gly	Val	Ile	Phe	Pro	Asp	Pro	Ser	Gly	Asp	Ser
				260					265					270
Tyr	Thr	Met	Ile	Ala	Thr	Phe	Ser	Met	Leu	Leu	Leu	Asp	Gly	Leu
				275					280					285
Ile	Tyr	Leu	Leu	Leu	Ala	Leu	Tyr	Phe	Asp	Lys	Ile	Leu	Pro	Tyr
				290					295					300
Gly	Asp	Glu	Arg	His	Tyr	Ser	Pro	Leu	Phe	Phe	Leu	Asn	Ser	Ser
				305					310					315
Ser	Cys	Phe	Gln	His	Gln	Arg	Thr	Asn	Ala	Lys	Val	Ile	Glu	Lys
				320					325					330
Glu	Ile	Asp	Ala	Glu	His	Pro	Ser	Asp	Asp	Tyr	Phe	Glu	Pro	Val
				335					340					345
Ala	Pro	Glu	Phe	Gln	Gly	Lys	Glu	Ala	Ile	Arg	Ile	Arg	Asn	Val
				350					355					360
Lys	Lys	Glu	Tyr	Lys	Gly	Lys	Ser	Gly	Lys	Val	Glu	Ala	Leu	Lys
				365					370					375
Gly	Leu	Leu	Phe	Asp	Ile	Tyr	Glu	Gly	Gln	Ile	Thr	Ala	Ile	Leu
				380					385					390
Gly	His	Ser	Gly	Ala	Gly	Lys	Ser	Ser	Leu	Leu	Asn	Ile	Leu	Asn
				395					400					405
Gly	Leu	Ser	Val	Pro	Thr	Glu	Gly	Ser	Val	Thr	Ile	Tyr	Asn	Lys
				410					415					420
Asn	Leu	Ser	Glu	Met	Gln	Asp	Leu	Glu	Glu	Ile	Arg	Lys	Ile	Thr
				425					430					435
Gly	Val	Cys	Pro	Gln	Phe	Asn	Val	Gln	Phe	Asp	Ile	Leu	Thr	Val
				440					445					450
Lys	Glu	Asn	Leu	Ser	Leu	Phe	Ala	Lys	Ile	Lys	Gly	Ile	His	Leu
				455					460					465
Lys	Glu	Val	Glu	Gln	Glu	Val	Gln	Arg	Ile	Leu	Leu	Glu	Leu	Asp
				470					475					480
Met	Gln	Asn	Ile	Gln	Asp	Asn	Leu	Ala	Lys	His	Leu	Ser	Glu	Gly
				485					490					495
Gln	Lys	Arg	Lys	Leu	Thr	Phe	Gly	Ile	Thr	Ile	Leu	Gly	Asp	Pro
				500					505					510
Gln	Ile	Leu	Leu	Leu	Asp	Glu	Pro	Thr	Thr	Gly	Leu	Asp	Pro	Phe
				515					520					525
Ser	Arg	Asp	Gln	Val	Trp	Ser	Leu	Leu	Arg	Glu	Arg	Arg	Ala	Asp
				530					535					540
His	Val	Ile	Leu	Phe	Ser	Thr	Gln	Ser	Met	Asp	Glu	Ala	Asp	Ile
				545					550					555
Leu	Ala	Asp	Arg	Lys	Val	Ile	Met	Ser	Asn	Gly	Arg	Leu	Lys	Cys
				560					565					570
Ala	Gly	Ser	Ser	Ile	Phe	Leu	Lys	Arg	Arg	Trp	Gly	Leu	Gly	Tyr
				575					580					585
His	Leu	Ser	Leu	His	Arg	Asn	Glu	Ile	Cys	Asn	Pro	Glu	Gln	Ile

Thr Ser Phe Ile	590	Thr His His Ile Pro	595	Ala Lys Leu Lys	600
605	605	610	610	615	615
Glu Asn Lys Glu	620	Leu Val Tyr Thr	625	Leu Pro Leu Glu Arg	630
635	635	640	640	645	645
Asn Thr Phe Pro	650	Leu Phe Ser Asp	655	Leu Asp Lys Cys Ser	660
665	665	670	670	675	675
Gln Gly Val Thr	680	Gly Tyr Asp Ile Ser	685	Met Ser Thr Leu Asn	690
695	695	700	700	705	705
Val Phe Met Lys	710	Leu Glu Gly Gln Ser	715	Thr Ile Glu Gln Asp	720
725	725	730	730	735	735
Glu Gln Val Glu	740	Met Ile Arg Asp Ser	745	Glu Ser Leu Asn Glu	750
755	755	760	760	765	765
Glu Leu Ala His	770	Ser Ser Phe Ser Glu	775	Met Gln Thr Ala Val	780
785	785	790	790	795	795
Asp Met Gly Leu	800	Trp Arg Met Gln Val	805	Phe Ala Met Ala Arg	810
815	815	820	820	825	825
Arg Phe Leu Lys	830	Leu Lys Arg Gln Thr	835	Lys Val Leu Leu Thr	840
845	845	850	850	855	855
Leu Leu Val Phe	860	Gly Ile Ala Ile Phe	865	Pro Leu Ile Val Glu	870
875	875	880	880	885	885
Ile Ile Tyr Ala	890	Met Leu Asn Glu Lys	895	Ile Asp Trp Glu Phe	900
905	905	910	910	915	915
Asn Glu Leu Tyr	920	Phe Leu Ser Pro Gly	925	Gln Leu Pro Gln Glu	930
935	935	940	940	945	945
Arg Thr Ser Leu	950	Leu Ile Ile Asn Asn	955	Thr Glu Ser Asn Ile	960
965	965	970	970	975	975
Asp Phe Ile Lys	980	Ser Leu Lys His Gln	985	Asn Ile Leu Leu Glu	990
995	995	1000	1000	1005	1005
Asp Asp Phe Glu	1010	Asn Arg Asn Gly Thr	1015	Asp Gly Leu Ser Tyr	1020
1025	1025	1030	1030	1035	1035
Gly Ala Ile Ile	1040	Val Ser Gly Lys Gln	1045	Lys Asp Tyr Arg Phe	1050
1055	1055	1060	1060	1065	1065
Val Val Cys Asn	1070	Thr Lys Arg Leu His	1075	Cys Phe Pro Ile Leu	1080
1085	1085	1090	1090	1095	1095
Asn Ile Ile Ser		Asn Gly Leu Leu Gln		Met Phe Asn His Thr	
860	860	865	865	870	870
His Ile Arg Ile	875	Glu Ser Ser Pro Phe	880	Pro Leu Ser His Ile	885
890	890	895	895	900	900
Leu Trp Thr Gly	905	Leu Pro Asp Gly Ser	910	Phe Phe Leu Phe Leu	915
920	920	925	925	930	930
Leu Cys Ser Ile	935	Ser Pro Tyr Ile Thr	940	Met Gly Ser Ile Ser	945
950	950	955	955	960	960
Tyr Lys Lys Asn	965	Ala Lys Ser Gln Leu	970	Trp Ile Ser Gly Leu	975
980	980	985	985	990	990
Thr Ser Ala Tyr	995	Trp Cys Gly Gln Ala	1000	Leu Val Asp Val Ser	1005
1010	1010	1015	1015	1020	1020
Phe Ile Leu Ile	1025	Leu Leu Leu Met Tyr	1030	Leu Ile Phe Tyr Ile	1035
1040	1040	1045	1045	1050	1050
Asn Met Gln Tyr	1055	Leu Leu Ile Thr Ser	1060	Gln Ile Val Phe Ala	1065
1070	1070	1075	1075	1080	1080
Val Ile Val Thr	1085	Pro Gly Tyr Ala Ala	1090	Ser Leu Val Phe Phe	1095
1095	1095	1100	1100	1105	1105
Tyr Met Ile Ser		Phe Ile Phe Arg Lys		Arg Arg Lys Asn Ser	
995	995	1000	1000	1005	1005
Leu Trp Ser Phe		Tyr Phe Phe Phe Ala		Ser Thr Ile Met Phe	
1010	1010	1015	1015	1020	1020
Ile Thr Leu Ile		Asn His Phe Asp Leu		Ser Ile Leu Ile Thr	
1025	1025	1030	1030	1035	1035
Met Val Leu Val		Pro Ser Tyr Thr Leu		Gly Phe Lys Thr Phe	
1040	1040	1045	1045	1050	1050
Leu Glu Val Arg		Asp Gln Glu His Tyr		Arg Glu Phe Pro Glu	
1055	1055	1060	1060	1065	1065
Asn Phe Glu Leu		Ser Ala Thr Asp Phe		Leu Val Cys Phe Ile	
1070	1070	1075	1075	1080	1080
Tyr Phe Gln Thr		Leu Leu Phe Val Phe		Val Leu Arg Cys Met	
1085	1085	1090	1090	1095	1095

```

Leu Lys Cys Gly Lys Lys Arg Met Arg Lys Asp Pro Val Phe Arg
1100 1105 1110
Ile Ser Pro Gln Ser Arg Asp Ala Lys Pro Asn Pro Glu Glu Pro
1115 1120 1125
Ile Asp Glu Asp Glu Asp Ile Gln Thr Glu Arg Ile Arg Thr Val
1130 1135 1140
Thr Ala Leu Thr Thr Ser Ile Leu Asp Glu Lys Pro Val Ile Ile
1145 1150 1155
Ala Ser Cys Leu His Lys Glu Tyr Ala Gly Gln Lys Lys Ser Cys
1160 1165 1170
Phe Ser Lys Arg Lys Lys Lys Ile Ala Ala Arg Asn Ile Ser Phe
1175 1180 1185
Cys Val Gln Glu Gly Glu Ile Leu Gly Leu Leu Gly Pro Ser Gly
1190 1195 1200
Ala Gly Lys Ser Ser Ser Ile Arg Met Ile Ser Gly Ile Thr Lys
1205 1210 1215
Pro Thr Ala Gly Glu Val Glu Leu Lys Gly Cys Ser Ser Val Leu
1220 1225 1230
Gly His Leu Gly Tyr Cys Pro Gln Glu Asn Val Leu Trp Pro Met
1235 1240 1245
Leu Thr Leu Arg Glu His Leu Glu Val Tyr Ala Ala Val Lys Gly
1250 1255 1260
Leu Arg Glu Ala Asp Ala Arg Leu Ala Ile Ala Arg Leu Val Ser
1265 1270 1275
Ala Phe Lys Leu His Glu Gln Leu Asn Val Pro Val Gln Lys Leu
1280 1285 1290
Thr Ala Gly Ile Thr Arg Lys Leu Cys Phe Val Leu Ser Leu Leu
1295 1300 1305
Gly Asn Ser Pro Val Leu Leu Leu Asp Glu Pro Ser Thr Gly Ile
1310 1315 1320
Asp Pro Thr Gly Gln Gln Gln Met Trp Gln Ala Ile Gln Ala Val
1325 1330 1335
Val Lys Asn Thr Glu Arg Gly Val Leu Leu Thr Thr His Asn Leu
1340 1345 1350
Ala Glu Ala Glu Ala Leu Cys Asp Arg Val Ala Ile Met Val Ser
1355 1360 1365
Gly Arg Leu Arg Cys Ile Gly Ser Ile Gln His Leu Lys Asn Lys
1370 1375 1380
Leu Gly Lys Asp Tyr Ile Leu Glu Leu Lys Val Lys Glu Thr Ser
1385 1390 1395
Gln Val Thr Leu Val His Thr Glu Ile Leu Lys Leu Phe Pro Gln
1400 1405 1410
Ala Ala Gly Gln Gln Arg Tyr Ser Ser Leu Leu Thr Tyr Lys Leu
1415 1420 1425
Pro Val Ala Asp Val Tyr Pro Leu Ser Gln Thr Phe His Lys Leu
1430 1435 1440
Glu Ala Val Lys His Asn Phe Asn Leu Glu Glu Tyr Ser Leu Ser
1445 1450 1455
Gln Cys Thr Leu Glu Lys Val Phe Leu Glu Leu Ser Lys Glu Gln
1460 1465 1470
Glu Val Gly Asn Phe Asp Glu Glu Ile Asp Thr Thr Met Arg Trp
1475 1480 1485
Lys Leu Leu Pro His Ser Asp Glu Pro
1490

```

<210> 25

<211> 774

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4599654CD1

<400> 25

```

Met Glu Ala Glu Gln Arg Pro Ala Ala Gly Ala Ser Glu Gly Ala
1 5 10 15

```

Thr	Pro	Gly	Leu	Glu	Ala	Val	Pro	Pro	Val	Ala	Pro	Pro	Pro	Ala
				20					25					30
Thr	Ala	Ala	Ser	Gly	Pro	Ile	Pro	Lys	Ser	Gly	Pro	Glu	Pro	Lys
				35					40					45
Arg	Arg	His	Leu	Gly	Thr	Leu	Leu	Gln	Pro	Thr	Val	Asn	Lys	Phe
				50					55					60
Ser	Leu	Arg	Val	Phe	Gly	Ser	His	Lys	Ala	Val	Glu	Ile	Glu	Gln
				65					70					75
Glu	Arg	Val	Lys	Ser	Ala	Gly	Ala	Trp	Ile	Ile	His	Pro	Tyr	Ser
				80					85					90
Asp	Phe	Arg	Phe	Tyr	Trp	Asp	Leu	Ile	Met	Leu	Leu	Leu	Met	Val
				95					100					105
Gly	Asn	Leu	Ile	Val	Leu	Pro	Val	Gly	Ile	Thr	Phe	Phe	Lys	Glu
				110					115					120
Glu	Asn	Ser	Pro	Pro	Trp	Ile	Val	Phe	Asn	Val	Leu	Ser	Asp	Thr
				125					130					135
Phe	Phe	Leu	Leu	Asp	Leu	Val	Leu	Asn	Phe	Arg	Thr	Gly	Ile	Val
				140					145					150
Val	Glu	Glu	Gly	Ala	Glu	Ile	Leu	Leu	Ala	Pro	Arg	Ala	Ile	Arg
				155					160					165
Thr	Arg	Tyr	Leu	Arg	Thr	Trp	Phe	Leu	Val	Asp	Leu	Ile	Ser	Ser
				170					175					180
Ile	Pro	Val	Asp	Tyr	Ile	Phe	Leu	Val	Val	Glu	Leu	Glu	Pro	Arg
				185					190					195
Leu	Asp	Ala	Glu	Val	Tyr	Lys	Thr	Ala	Arg	Ala	Leu	Arg	Ile	Val
				200					205					210
Arg	Phe	Thr	Lys	Ile	Leu	Ser	Leu	Leu	Arg	Leu	Leu	Arg	Leu	Ser
				215					220					225
Arg	Leu	Ile	Arg	Tyr	Ile	His	Gln	Trp	Glu	Glu	Ile	Phe	His	Met
				230					235					240
Thr	Tyr	Asp	Leu	Ala	Ser	Ala	Val	Val	Arg	Ile	Phe	Asn	Leu	Ile
				245					250					255
Gly	Met	Met	Leu	Leu	Leu	Cys	His	Trp	Asp	Gly	Cys	Leu	Gln	Phe
				260					265					270
Leu	Val	Pro	Met	Leu	Gln	Asp	Phe	Pro	Pro	Asp	Cys	Trp	Val	Ser
				275					280					285
Ile	Asn	His	Met	Val	Asn	His	Ser	Trp	Gly	Arg	Gln	Tyr	Ser	His
				290					295					300
Ala	Leu	Phe	Lys	Ala	Met	Ser	His	Met	Leu	Cys	Ile	Gly	Tyr	Gly
				305					310					315
Gln	Gln	Ala	Pro	Val	Gly	Met	Pro	Asp	Val	Trp	Leu	Thr	Met	Leu
				320					325					330
Ser	Met	Ile	Val	Gly	Ala	Thr	Cys	Tyr	Ala	Met	Phe	Ile	Gly	His
				335					340					345
Ala	Thr	Ala	Leu	Ile	Gln	Ser	Leu	Asp	Ser	Ser	Arg	Arg	Gln	Tyr
				350					355					360
Gln	Glu	Lys	Tyr	Lys	Gln	Val	Glu	Gln	Tyr	Met	Ser	Phe	His	Lys
				365					370					375
Leu	Pro	Ala	Asp	Thr	Arg	Gln	Arg	Ile	His	Glu	Tyr	Tyr	Glu	His
				380					385					390
Arg	Tyr	Gln	Gly	Lys	Met	Phe	Asp	Glu	Glu	Ser	Ile	Leu	Gly	Glu
				395					400					405
Leu	Ser	Glu	Pro	Leu	Arg	Glu	Glu	Ile	Ile	Asn	Phe	Thr	Cys	Arg
				410					415					420
Gly	Leu	Val	Ala	His	Met	Pro	Leu	Phe	Ala	His	Ala	Asp	Pro	Ser
				425					430					435
Phe	Val	Thr	Ala	Val	Leu	Thr	Lys	Leu	Arg	Phe	Glu	Val	Phe	Gln
				440					445					450
Pro	Gly	Asp	Leu	Val	Val	Arg	Glu	Gly	Ser	Val	Gly	Arg	Lys	Met
				455					460					465
Tyr	Phe	Ile	Gln	His	Gly	Leu	Leu	Ser	Val	Leu	Ala	Arg	Gly	Ala
				470					475					480
Arg	Asp	Thr	Arg	Leu	Thr	Asp	Gly	Ser	Tyr	Phe	Gly	Glu	Ile	Cys
				485					490					495
Leu	Leu	Thr	Arg	Gly	Arg	Arg	Thr	Ala	Ser	Val	Arg	Ala	Asp	Thr
				500					505					510
Tyr	Cys	Arg	Leu	Tyr	Ser	Leu	Ser	Val	Asp	His	Phe	Asn	Ala	Val

Leu	Glu	Glu	Phe	515	Pro	Met	Met	Arg	Arg	520	Ala	Phe	Glu	Thr	Val	525	Ala
				530	Leu	Arg	Ile	Gly	Lys	535	Lys	Asn	Ser	Ile	Leu	540	Gln
Met	Asp	Arg	Leu	545	Glu	Pro	Ser	Pro	Gly	550	Ser	Ser	Gly	Gly	Ile	555	Met
Arg	Lys	Arg	Ser	560	Val	Gln	His	Asp	Arg	565	Asp	Met	Ala	Arg	Gly	570	Val
Glu	Gln	His	Leu	575	Pro	Ser	Thr	Gly	Ala	580	Gln	Leu	Ser	Gly	Lys	585	Pro
Arg	Gly	Arg	Ala	590	Pro	Leu	Val	His	Ala	595	Pro	Leu	Gln	Ala	Ala	600	Ala
Val	Leu	Trp	Glu	605	Val	Ala	Ile	Ala	Leu	610	Thr	His	Gln	Arg	Gly	615	Pro
Val	Thr	Ser	Asn	620	Pro	Asp	Ser	Pro	Ala	625	Leu	Leu	Ala	Arg	Ser	630	Ser
Leu	Pro	Leu	Ser	635	Ala	Gly	Ser	Pro	Ala	640	Ser	Pro	Leu	Val	Pro	645	Val
Ala	Trp	Arg	Ser	650	Trp	Ala	Ser	Thr	Ser	655	Arg	Leu	Pro	Ala	Pro	660	Pro
Arg	Ala	Gly	Pro	665	His	Ala	Ser	Leu	Ser	670	Arg	Ala	Gly	Arg	Ser	675	Gln
Ala	Arg	Thr	Leu	680	Gly	Pro	Pro	Pro	Gly	685	Gly	Gly	Gly	Arg	Arg	690	Leu
Val	Ser	Leu	Leu	695	Arg	Pro	Leu	Ser	Ala	700	Ser	Gln	Pro	Ser	Leu	705	Pro
Gly	Pro	Arg	Gly	710	Gly	Asp	Gly	Ser	Pro	715	Gly	Arg	Lys	Gly	Ser	720	Gly
Gln	Arg	Ala	Thr	725	Pro	Pro	Ser	Gly	Leu	730	Leu	Ala	Lys	Pro	Pro	735	Arg
Ser	Glu	Arg	Leu	740	Pro	Arg	Pro	Pro	Val	745	Pro	Glu	Pro	Ala	Thr	750	Pro
Thr	Ala	Gln	Pro	755	Leu	Ser	Ala	Asn	Met	760						765	
Arg	Gly	Leu	Gln	770													

<210> 26

<211> 614

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5047435CD1

<400> 26

Met	Ala	Glu	Gly	Glu	Arg	Gly	Ala	Asp	Val	Pro	His	Gly	Leu	Gly			
1				5					10					15			
Ala	Trp	Leu	Ala	Asp	Val	Ala	Leu	Ala	Ala	Leu	Arg	Ala	Gly	Gly			
				20					25					30			
Gln	Gly	Arg	Arg	Asp	Arg	Gly	Gly	Gly	Gly	Pro	Glu	Ser	Leu	Ser			
				35					40					45			
Gly	Gly	Ser	Gly	Val	Gly	Asp	Ser	Gly	Gly	Gly	Cys	Ala	Pro	Gly			
				50					55					60			
Pro	Ser	Ala	Pro	Pro	Ala	Arg	Arg	Arg	Val	Pro	Leu	Ala	Met	Gly			
				65					70					75			
His	Ser	Pro	Pro	Val	Leu	Pro	Leu	Cys	Ala	Ser	Val	Ser	Leu	Leu			
				80					85					90			
Gly	Gly	Leu	Thr	Phe	Gly	Tyr	Glu	Leu	Ala	Val	Ile	Ser	Gly	Ala			
				95					100					105			
Leu	Leu	Pro	Leu	Gln	Leu	Asp	Phe	Gly	Leu	Ser	Cys	Leu	Glu	Gln			
				110					115					120			
Glu	Phe	Leu	Val	Gly	Ser	Leu	Leu	Leu	Gly	Ala	Leu	Leu	Ala	Ser			
				125					130					135			
Leu	Val	Gly	Gly	Phe	Leu	Ile	Asp	Cys	Tyr	Gly	Arg	Lys	Gln	Ala			
				140					145					150			
Ile	Leu	Gly	Ser	Asn	Leu	Val	Leu	Leu	Ala	Gly	Ser	Leu	Thr	Leu			

Gly	Leu	Ala	Gly	155	Ser	Leu	Ala	Trp	Leu	160	Val	Leu	Gly	Arg	Ala	165	Val
Val	Gly	Phe	Ala	170	Ile	Ser	Leu	Ser	Ser	175	Met	Ala	Cys	Cys	Ile	180	Tyr
Val	Ser	Glu	Leu	185	Val	Gly	Pro	Arg	Gln	190	Arg	Gly	Val	Leu	Val	195	Ser
Leu	Tyr	Glu	Ala	200	Gly	Ile	Thr	Val	Gly	205	Ile	Leu	Leu	Ser	Tyr	210	Ala
Leu	Asn	Tyr	Ala	215	Leu	Ala	Gly	Thr	Pro	220	Trp	Gly	Trp	Arg	His	225	Met
Phe	Gly	Trp	Ala	230	Thr	Ala	Pro	Ala	Val	235	Leu	Gln	Ser	Leu	Ser	240	Leu
Leu	Phe	Leu	Pro	245	Ala	Gly	Thr	Asp	Glu	250	Thr	Ala	Thr	His	Lys	255	Asp
Leu	Ile	Pro	Leu	260	Gln	Gly	Gly	Glu	Ala	265	Pro	Lys	Leu	Gly	Pro	270	Gly
Arg	Pro	Arg	Tyr	275	Ser	Phe	Leu	Asp	Leu	280	Phe	Arg	Ala	Arg	Asp	285	Asn
Met	Arg	Gly	Arg	290	Thr	Thr	Val	Gly	Leu	295	Gly	Leu	Val	Leu	Phe	300	Gln
Gln	Leu	Thr	Gly	305	Gln	Pro	Asn	Val	Leu	310	Cys	Tyr	Ala	Ser	Thr	315	Ile
Phe	Ser	Ser	Val	320	Gly	Phe	His	Gly	Gly	325	Ser	Ser	Ala	Val	Leu	330	Ala
Ser	Val	Gly	Leu	335	Gly	Ala	Val	Lys	Val	340	Ala	Ala	Thr	Leu	Thr	345	Ala
Met	Gly	Leu	Val	350	Asp	Arg	Ala	Gly	Arg	355	Arg	Ala	Leu	Leu	Leu	360	Ala
Gly	Cys	Ala	Leu	365	Met	Ala	Leu	Ser	Val	370	Ser	Gly	Ile	Gly	Leu	375	Val
Ser	Phe	Ala	Val	380	Pro	Met	Asp	Ser	Gly	385	Pro	Ser	Cys	Leu	Ala	390	Val
Pro	Asn	Ala	Thr	395	Gly	Gln	Thr	Gly	Leu	400	Pro	Gly	Asp	Ser	Gly	405	Leu
Leu	Gln	Asp	Ser	410	Ser	Leu	Pro	Pro	Ile	415	Pro	Arg	Thr	Asn	Glu	420	Asp
Gln	Arg	Glu	Pro	425	Ile	Leu	Ser	Thr	Ala	430	Lys	Lys	Thr	Lys	Pro	435	His
Pro	Arg	Ser	Gly	440	Asp	Pro	Ser	Ala	Pro	445	Pro	Arg	Leu	Ala	Leu	450	Ser
Ser	Ala	Leu	Pro	455	Gly	Pro	Pro	Leu	Pro	460	Ala	Arg	Gly	His	Ala	465	Leu
Leu	Arg	Trp	Thr	470	Ala	Leu	Leu	Cys	Leu	475	Met	Val	Phe	Val	Ser	480	Ala
Phe	Ser	Phe	Gly	485	Phe	Gly	Pro	Val	Thr	490	Trp	Leu	Val	Leu	Ser	495	Glu
Ile	Tyr	Pro	Val	500	Glu	Ile	Arg	Gly	Arg	505	Ala	Phe	Ala	Phe	Cys	510	Asn
Ser	Phe	Asn	Trp	515	Ala	Ala	Asn	Leu	Phe	520	Ile	Ser	Leu	Ser	Phe	525	Leu
Asp	Leu	Ile	Gly	530	Thr	Ile	Gly	Leu	Ser	535	Trp	Thr	Phe	Leu	Leu	540	Tyr
Gly	Leu	Thr	Ala	545	Val	Leu	Gly	Leu	Gly	550	Phe	Ile	Tyr	Leu	Phe	555	Val
Pro	Glu	Thr	Lys	560	Gly	Gln	Ser	Leu	Ala	565	Glu	Ile	Asp	Gln	Gln	570	Phe
Gln	Lys	Arg	Arg	575	Phe	Thr	Leu	Ser	Phe	580	Gly	His	Arg	Gln	Asn	585	Ser
Thr	Gly	Ile	Pro	590	Tyr	Ser	Arg	Ile	Glu	595	Ile	Ser	Ala	Ala	Ser	600	
				605						610							

<210> 27
 <211> 2180
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7475603CD1

<400> 27

Met	Arg	Phe	Arg	Lys	Gly	Gln	Glu	Leu	Pro	Ala	Ala	Ala	Pro	His
1				5					10					15
Val	Phe	Ser	Pro	Thr	Val	Val	Leu	Thr	Ser	Leu	Ser	Arg	Pro	Leu
				20					25					30
Pro	Ser	Leu	Thr	Met	Ala	Phe	Trp	Thr	Gln	Leu	Met	Leu	Leu	Leu
				35					40					45
Trp	Lys	Asn	Phe	Met	Tyr	Arg	Arg	Arg	Gln	Pro	Val	Gln	Leu	Leu
				50					55					60
Val	Glu	Leu	Leu	Trp	Pro	Leu	Phe	Leu	Phe	Phe	Ile	Leu	Val	Ala
				65					70					75
Val	Arg	His	Ser	His	Pro	Pro	Leu	Glu	His	His	Glu	Cys	His	Phe
				80					85					90
Pro	Asn	Lys	Pro	Leu	Pro	Ser	Ala	Gly	Thr	Val	Pro	Trp	Leu	Gln
				95					100					105
Gly	Leu	Ile	Cys	Asn	Val	Asn	Asn	Thr	Cys	Phe	Pro	Gln	Leu	Thr
				110					115					120
Pro	Gly	Glu	Glu	Pro	Gly	Arg	Leu	Ser	Asn	Phe	Asn	Asp	Ser	Leu
				125					130					135
Val	Ser	Arg	Leu	Leu	Ala	Asp	Ala	Arg	Thr	Val	Leu	Gly	Gly	Ala
				140					145					150
Ser	Ala	His	Arg	Thr	Leu	Ala	Gly	Leu	Gly	Lys	Leu	Ile	Ala	Thr
				155					160					165
Leu	Arg	Ala	Ala	Arg	Ser	Thr	Ala	Gln	Pro	Gln	Pro	Thr	Lys	Gln
				170					175					180
Ser	Pro	Leu	Glu	Pro	Pro	Met	Leu	Asp	Val	Ala	Glu	Leu	Leu	Thr
				185					190					195
Ser	Leu	Leu	Arg	Thr	Glu	Ser	Leu	Gly	Leu	Ala	Leu	Gly	Gln	Ala
				200					205					210
Gln	Glu	Pro	Leu	His	Ser	Leu	Leu	Glu	Ala	Ala	Glu	Asp	Leu	Ala
				215					220					225
Gln	Glu	Leu	Leu	Ala	Leu	Arg	Ser	Leu	Val	Glu	Leu	Arg	Ala	Leu
				230					235					240
Leu	Gln	Arg	Pro	Arg	Gly	Thr	Ser	Gly	Pro	Leu	Glu	Leu	Leu	Ser
				245					250					255
Glu	Ala	Leu	Cys	Ser	Val	Arg	Gly	Pro	Ser	Ser	Thr	Val	Gly	Pro
				260					265					270
Ser	Leu	Asn	Trp	Tyr	Glu	Ala	Ser	Asp	Leu	Met	Glu	Leu	Val	Gly
				275					280					285
Gln	Glu	Pro	Glu	Ser	Ala	Leu	Pro	Asp	Ser	Ser	Leu	Ser	Pro	Ala
				290					295					300
Cys	Ser	Glu	Leu	Ile	Gly	Ala	Leu	Asp	Ser	His	Pro	Leu	Ser	Arg
				305					310					315
Leu	Leu	Trp	Arg	Arg	Leu	Lys	Pro	Leu	Ile	Leu	Gly	Lys	Leu	Leu
				320					325					330
Phe	Ala	Pro	Asp	Thr	Pro	Phe	Thr	Arg	Lys	Leu	Met	Ala	Gln	Val
				335					340					345
Asn	Arg	Thr	Phe	Glu	Glu	Leu	Thr	Leu	Leu	Arg	Asp	Val	Arg	Glu
				350					355					360
Val	Trp	Glu	Met	Leu	Gly	Pro	Arg	Ile	Phe	Thr	Phe	Met	Asn	Asp
				365					370					375
Ser	Ser	Asn	Val	Ala	Met	Leu	Gln	Arg	Leu	Leu	Gln	Met	Gln	Asp
				380					385					390
Glu	Gly	Arg	Arg	Gln	Pro	Arg	Pro	Gly	Gly	Arg	Asp	His	Met	Glu
				395					400					405
Ala	Leu	Arg	Ser	Phe	Leu	Asp	Pro	Gly	Ser	Gly	Gly	Tyr	Ser	Trp
				410					415					420
Gln	Asp	Ala	His	Ala	Asp	Val	Gly	His	Leu	Val	Gly	Thr	Leu	Gly
				425					430					435
Arg	Val	Thr	Glu	Cys	Leu	Ser	Leu	Asp	Lys	Leu	Glu	Ala	Ala	Pro
				440					445					450
Ser	Glu	Ala	Ala	Leu	Val	Ser	Arg	Ala	Leu	Gln	Leu	Leu	Ala	Glu
				455					460					465

His	Arg	Phe	Trp	Ala	Gly	Val	Val	Phe	Leu	Gly	Pro	Glu	Asp	Ser
				470					475					480
Ser	Asp	Pro	Thr	Glu	His	Pro	Thr	Pro	Asp	Leu	Gly	Pro	Gly	His
				485					490					495
Val	Arg	Ile	Lys	Ile	Arg	Met	Asp	Ile	Asp	Val	Val	Thr	Arg	Thr
				500					505					510
Asn	Lys	Ile	Arg	Asp	Arg	Phe	Trp	Asp	Pro	Gly	Pro	Ala	Ala	Asp
				515					520					525
Pro	Leu	Thr	Asp	Leu	Arg	Tyr	Val	Trp	Gly	Gly	Phe	Val	Tyr	Leu
				530					535					540
Gln	Asp	Leu	Val	Glu	Arg	Ala	Ala	Val	Arg	Val	Leu	Ser	Gly	Ala
				545					550					555
Asn	Pro	Arg	Ala	Gly	Leu	Tyr	Leu	Gln	Gln	Met	Pro	Tyr	Pro	Cys
				560					565					570
Tyr	Val	Asp	Asp	Val	Phe	Leu	Arg	Val	Leu	Ser	Arg	Ser	Leu	Pro
				575					580					585
Leu	Phe	Leu	Thr	Leu	Ala	Trp	Ile	Tyr	Ser	Val	Thr	Leu	Thr	Val
				590					595					600
Lys	Ala	Val	Val	Arg	Glu	Lys	Glu	Thr	Arg	Leu	Arg	Asp	Thr	Met
				605					610					615
Arg	Ala	Met	Gly	Leu	Ser	Arg	Ala	Val	Leu	Trp	Leu	Gly	Trp	Phe
				620					625					630
Leu	Ser	Cys	Leu	Gly	Pro	Phe	Leu	Leu	Ser	Ala	Ala	Leu	Leu	Val
				635					640					645
Leu	Val	Leu	Lys	Leu	Gly	Asp	Ile	Leu	Pro	Tyr	Ser	His	Pro	Gly
				650					655					660
Val	Val	Phe	Leu	Phe	Leu	Ala	Ala	Phe	Ala	Val	Ala	Thr	Val	Thr
				665					670					675
Gln	Ser	Phe	Leu	Leu	Ser	Ala	Phe	Phe	Ser	Arg	Ala	Asn	Leu	Ala
				680					685					690
Ala	Ala	Cys	Gly	Gly	Leu	Ala	Tyr	Phe	Ser	Leu	Tyr	Leu	Pro	Tyr
				695					700					705
Val	Leu	Cys	Val	Ala	Trp	Arg	Asp	Arg	Leu	Pro	Ala	Gly	Gly	Arg
				710					715					720
Val	Ala	Ala	Ser	Leu	Leu	Ser	Pro	Val	Ala	Phe	Gly	Phe	Gly	Cys
				725					730					735
Glu	Ser	Leu	Ala	Leu	Leu	Glu	Glu	Gln	Gly	Glu	Gly	Ala	Gln	Trp
				740					745					750
His	Asn	Val	Gly	Thr	Arg	Pro	Thr	Ala	Asp	Val	Phe	Ser	Leu	Ala
				755					760					765
Gln	Val	Ser	Gly	Leu	Leu	Leu	Leu	Asp	Ala	Ala	Leu	Tyr	Gly	Leu
				770					775					780
Ala	Thr	Trp	Tyr	Leu	Glu	Ala	Val	Cys	Pro	Gly	Gln	Tyr	Gly	Ile
				785					790					795
Pro	Glu	Pro	Trp	Asn	Phe	Pro	Phe	Arg	Arg	Ser	Tyr	Trp	Cys	Gly
				800					805					810
Pro	Arg	Pro	Pro	Lys	Ser	Pro	Ala	Pro	Cys	Pro	Thr	Pro	Leu	Asp
				815					820					825
Pro	Lys	Val	Leu	Val	Glu	Glu	Ala	Pro	Pro	Gly	Leu	Ser	Pro	Gly
				830					835					840
Val	Ser	Val	Arg	Ser	Leu	Glu	Lys	Arg	Phe	Pro	Gly	Ser	Pro	Gln
				845					850					855
Pro	Ala	Leu	Arg	Gly	Leu	Ser	Leu	Asp	Phe	Tyr	Gln	Gly	His	Ile
				860					865					870
Thr	Ala	Phe	Leu	Gly	His	Asn	Gly	Ala	Gly	Lys	Thr	Thr	Thr	Leu
				875					880					885
Ser	Ile	Leu	Ser	Gly	Leu	Phe	Pro	Pro	Ser	Gly	Gly	Ser	Ala	Phe
				890					895					900
Ile	Leu	Gly	His	Asp	Val	Arg	Ser	Ser	Met	Ala	Ala	Ile	Arg	Pro
				905					910					915
His	Leu	Gly	Val	Cys	Pro	Gln	Tyr	Asn	Val	Leu	Phe	Asp	Met	Leu
				920					925					930
Thr	Val	Asp	Glu	His	Val	Trp	Phe	Tyr	Gly	Arg	Leu	Lys	Gly	Leu
				935					940					945
Ser	Ala	Ala	Val	Val	Gly	Pro	Glu	Gln	Asp	Arg	Leu	Leu	Gln	Asp
				950					955					960
Val	Gly	Leu	Val	Ser	Lys	Gln	Ser	Val	Gln	Thr	Arg	His	Leu	Ser

	965		970		975
Gly Gly Met Gln Arg	Lys Leu Ser Val Ala	Ile Ala Phe Val Gly			
	980		985		990
Gly Ser Gln Val Val	Ile Leu Asp Glu Pro	Thr Ala Gly Val Asp			
	995		1000		1005
Pro Ala Ser Arg Arg	Gly Ile Trp Glu Leu	Leu Leu Lys Tyr Arg			
	1010		1015		1020
Glu Gly Arg Thr Leu	Ile Leu Ser Thr His	His Leu Asp Glu Ala			
	1025		1030		1035
Glu Leu Leu Gly Asp	Arg Val Ala Val Val	Ala Gly Gly Arg Leu			
	1040		1045		1050
Cys Cys Cys Gly Ser	Pro Leu Phe Leu Arg	Arg His Leu Gly Ser			
	1055		1060		1065
Gly Tyr Tyr Leu Thr	Leu Val Lys Ala Arg	Leu Pro Leu Thr Thr			
	1070		1075		1080
Asn Glu Lys Ala Asp	Thr Asp Met Glu Gly	Ser Val Asp Thr Arg			
	1085		1090		1095
Gln Glu Lys Lys Asn	Gly Ser Gln Gly Ser	Arg Val Gly Thr Pro			
	1100		1105		1110
Gln Leu Leu Ala Leu	Val Gln His Trp Val	Pro Gly Ala Arg Leu			
	1115		1120		1125
Val Glu Glu Leu Pro	His Glu Leu Val Leu	Val Leu Pro Tyr Thr			
	1130		1135		1140
Gly Ala His Asp Gly	Ser Phe Ala Thr Leu	Phe Arg Glu Leu Asp			
	1145		1150		1155
Thr Arg Leu Ala Glu	Leu Arg Leu Thr Gly	Tyr Gly Ile Ser Asp			
	1160		1165		1170
Thr Ser Leu Glu Glu	Ile Phe Leu Lys Val	Val Glu Glu Cys Ala			
	1175		1180		1185
Ala Asp Thr Asp Met	Glu Asp Gly Ser Cys	Gly Gln His Leu Cys			
	1190		1195		1200
Thr Gly Ile Ala Gly	Leu Asp Val Thr Leu	Arg Leu Lys Met Pro			
	1205		1210		1215
Pro Gln Glu Thr Ala	Leu Glu Asn Gly Glu	Pro Ala Gly Ser Ala			
	1220		1225		1230
Pro Glu Thr Asp Gln	Gly Ser Gly Pro Asp	Ala Val Gly Arg Val			
	1235		1240		1245
Gln Gly Trp Ala Leu	Thr Arg Gln Gln Leu	Gln Ala Leu Leu Leu			
	1250		1255		1260
Lys Arg Phe Leu Leu	Ala Arg Arg Ser Arg	Arg Gly Leu Phe Ala			
	1265		1270		1275
Gln Ile Val Leu Pro	Ala Leu Phe Val Gly	Leu Ala Leu Val Phe			
	1280		1285		1290
Ser Leu Ile Val Pro	Pro Phe Gly His Tyr	Pro Ala Leu Arg Leu			
	1295		1300		1305
Ser Pro Thr Met Tyr	Gly Ala Gln Val Ser	Phe Phe Ser Glu Asp			
	1310		1315		1320
Ala Pro Gly Asp Pro	Gly Arg Ala Arg Leu	Leu Glu Ala Leu Leu			
	1325		1330		1335
Gln Glu Ala Gly Leu	Glu Glu Pro Pro Val	Gln His Ser Ser His			
	1340		1345		1350
Arg Phe Ser Ala Pro	Glu Val Pro Ala Glu	Val Ala Lys Val Leu			
	1355		1360		1365
Ala Ser Gly Asn Trp	Thr Pro Glu Ser Pro	Ser Pro Ala Cys Gln			
	1370		1375		1380
Cys Ser Arg Pro Gly	Ala Arg Arg Leu Leu	Pro Asp Cys Pro Ala			
	1385		1390		1395
Ala Ala Gly Gly Pro	Pro Pro Pro Gln Ala	Val Thr Gly Ser Gly			
	1400		1405		1410
Glu Val Val Gln Asn	Gln Thr Gly Arg Asn	Leu Ser Asp Phe Leu			
	1415		1420		1425
Val Lys Thr Tyr Pro	Arg Leu Val Arg Gln	Gly Leu Lys Thr Lys			
	1430		1435		1440
Lys Trp Val Asn Glu	Val Arg Tyr Gly Gly	Phe Ser Leu Gly Gly			
	1445		1450		1455
Arg Asp Pro Gly Leu	Pro Ser Gly Gln Glu	Leu Gly Arg Ser Val			
	1460		1465		1470

Glu	Glu	Leu	Trp	Ala	Leu	Leu	Ser	Pro	Leu	Pro	Gly	Gly	Ala	Leu
				1475					1480					1485
Asp	Arg	Val	Leu	Lys	Asn	Leu	Thr	Ala	Trp	Ala	His	Ser	Leu	Asp
				1490					1495					1500
Ala	Gln	Asp	Ser	Leu	Lys	Ile	Trp	Phe	Asn	Asn	Lys	Gly	Trp	His
				1505					1510					1515
Ser	Met	Val	Ala	Phe	Val	Asn	Arg	Ala	Ser	Asn	Ala	Ile	Leu	Arg
				1520					1525					1530
Ala	His	Leu	Pro	Pro	Gly	Pro	Ala	Arg	His	Ala	His	Ser	Ile	Thr
				1535					1540					1545
Thr	Leu	Asn	His	Pro	Leu	Asn	Leu	Thr	Lys	Glu	Gln	Leu	Ser	Glu
				1550					1555					1560
Ala	Ala	Leu	Met	Ala	Ser	Ser	Val	Asp	Val	Leu	Val	Ser	Ile	Cys
				1565					1570					1575
Val	Val	Phe	Ala	Met	Ser	Phe	Val	Pro	Ala	Ser	Phe	Thr	Leu	Val
				1580					1585					1590
Leu	Ile	Glu	Glu	Arg	Val	Thr	Arg	Ala	Lys	His	Leu	Gln	Leu	Met
				1595					1600					1605
Gly	Gly	Leu	Ser	Pro	Thr	Leu	Tyr	Trp	Leu	Gly	Asn	Phe	Leu	Trp
				1610					1615					1620
Asp	Met	Cys	Asn	Tyr	Leu	Val	Pro	Ala	Cys	Ile	Val	Val	Leu	Ile
				1625					1630					1635
Phe	Leu	Ala	Phe	Gln	Gln	Arg	Ala	Tyr	Val	Ala	Pro	Ala	Asn	Leu
				1640					1645					1650
Pro	Ala	Leu	Leu	Leu	Leu	Leu	Leu	Leu	Tyr	Gly	Trp	Ser	Ile	Thr
				1655					1660					1665
Pro	Leu	Met	Tyr	Pro	Ala	Ser	Phe	Phe	Phe	Ser	Val	Pro	Ser	Thr
				1670					1675					1680
Ala	Tyr	Val	Val	Leu	Thr	Cys	Ile	Asn	Leu	Phe	Ile	Gly	Ile	Asn
				1685					1690					1695
Gly	Ser	Met	Ala	Thr	Phe	Val	Leu	Glu	Leu	Phe	Ser	Asp	Gln	Lys
				1700					1705					1710
Leu	Gln	Glu	Val	Ser	Arg	Ile	Leu	Lys	Gln	Val	Phe	Leu	Ile	Phe
				1715					1720					1725
Pro	His	Phe	Cys	Leu	Gly	Arg	Gly	Leu	Ile	Asp	Met	Val	Arg	Asn
				1730					1735					1740
Gln	Ala	Met	Ala	Asp	Ala	Phe	Glu	Arg	Leu	Gly	Asp	Arg	Gln	Phe
				1745					1750					1755
Gln	Ser	Pro	Leu	Arg	Trp	Glu	Val	Val	Gly	Lys	Asn	Leu	Leu	Ala
				1760					1765					1770
Met	Val	Ile	Gln	Gly	Pro	Leu	Phe	Leu	Leu	Phe	Thr	Leu	Leu	Leu
				1775					1780					1785
Gln	His	Arg	Ser	Gln	Leu	Leu	Pro	Gln	Pro	Arg	Val	Arg	Ser	Leu
				1790					1795					1800
Pro	Leu	Leu	Gly	Glu	Glu	Asp	Glu	Asp	Val	Ala	Arg	Glu	Arg	Glu
				1805					1810					1815
Arg	Val	Val	Gln	Gly	Ala	Thr	Gln	Gly	Asp	Val	Leu	Val	Leu	Arg
				1820					1825					1830
Asn	Leu	Thr	Lys	Val	Tyr	Arg	Gly	Gln	Arg	Met	Pro	Ala	Val	Asp
				1835					1840					1845
Arg	Leu	Cys	Leu	Gly	Ile	Pro	Pro	Gly	Glu	Cys	Phe	Gly	Leu	Leu
				1850					1855					1860
Gly	Val	Asn	Gly	Ala	Gly	Lys	Thr	Ser	Thr	Phe	Arg	Met	Val	Thr
				1865					1870					1875
Gly	Asp	Thr	Leu	Ala	Ser	Arg	Gly	Glu	Ala	Val	Leu	Ala	Gly	His
				1880					1885					1890
Ser	Val	Ala	Arg	Glu	Pro	Ser	Ala	Ala	His	Leu	Ser	Met	Gly	Tyr
				1895					1900					1905
Cys	Pro	Gln	Ser	Asp	Ala	Ile	Phe	Glu	Leu	Leu	Thr	Gly	Arg	Glu
				1910					1915					1920
His	Leu	Glu	Leu	Leu	Ala	Arg	Leu	Arg	Gly	Val	Pro	Glu	Ala	Gln
				1925					1930					1935
Val	Ala	Gln	Thr	Ala	Gly	Ser	Gly	Leu	Ala	Arg	Leu	Gly	Leu	Ser
				1940					1945					1950
Trp	Tyr	Ala	Asp	Arg	Pro	Ala	Gly	Thr	Tyr	Ser	Gly	Gly	Asn	Lys
				1955					1960					1965
Arg	Lys	Leu	Ala	Thr	Ala	Leu	Ala	Leu	Val	Gly	Asp	Pro	Ala	Val

1970	1975	1980
Val Phe Leu Asp Glu	Pro Thr Thr Gly Met	Asp Pro Ser Ala Arg
1985	1990	1995
Arg Phe Leu Trp Asn	Ser Leu Leu Ala Val	Val Arg Glu Gly Arg
2000	2005	2010
Ser Val Met Leu Thr	Ser His Ser Met Glu	Glu Cys Glu Ala Leu
2015	2020	2025
Cys Ser Arg Leu Ala	Ile Met Val Asn Gly	Arg Phe Arg Cys Leu
2030	2035	2040
Gly Ser Pro Gln His	Leu Lys Gly Arg Phe	Ala Ala Gly His Thr
2045	2050	2055
Leu Thr Leu Arg Val	Pro Ala Ala Arg Ser	Gln Pro Ala Ala Ala
2060	2065	2070
Phe Val Ala Ala Glu	Phe Pro Gly Ala Glu	Leu Arg Glu Ala His
2075	2080	2085
Gly Gly Arg Leu Arg	Phe Gln Leu Pro Pro	Gly Gly Arg Cys Ala
2090	2095	2100
Leu Ala Arg Val Phe	Gly Glu Leu Ala Val	His Gly Ala Glu His
2105	2110	2115
Gly Val Glu Asp Phe	Ser Val Ser Gln Thr	Met Leu Glu Glu Val
2120	2125	2130
Phe Leu Tyr Phe Ser	Lys Asp Gln Gly Lys	Asp Glu Asp Thr Glu
2135	2140	2145
Glu Gln Lys Glu Ala	Gly Val Gly Val Asp	Pro Ala Pro Gly Leu
2150	2155	2160
Gln His Pro Lys Arg	Val Ser Gln Phe Leu	Asp Asp Pro Ser Thr
2165	2170	2175
Ala Glu Thr Val Leu		
2180		

<210> 28
 <211> 1737
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7477845CD1

<400> 28
Met Leu Lys Arg Lys Gln Ser Ser Arg Val Glu Ala Gln Pro Val
1 5 10 15
Thr Asp Phe Gly Pro Asp Glu Ser Leu Ser Asp Asn Ala Asp Ile
20 25 30
Leu Trp Ile Asn Lys Pro Trp Val His Ser Leu Leu Arg Ile Cys
35 40 45
Ala Ile Ile Ser Val Ile Ser Val Cys Met Asn Thr Pro Met Thr
50 55 60
Phe Glu His Tyr Pro Pro Leu Gln Tyr Val Thr Phe Thr Leu Asp
65 70 75
Thr Leu Leu Met Phe Leu Tyr Thr Ala Glu Met Ile Ala Lys Met
80 85 90
His Ile Arg Gly Ile Val Lys Gly Asp Ser Ser Tyr Val Lys Asp
95 100 105
Arg Trp Cys Val Phe Asp Gly Phe Met Val Phe Cys Leu Trp Val
110 115 120
Ser Leu Val Leu Gln Val Phe Glu Ile Ala Asp Ile Val Asp Gln
125 130 135
Met Ser Pro Trp Gly Met Leu Arg Ile Pro Arg Pro Leu Ile Met
140 145 150
Ile Arg Ala Phe Arg Ile Tyr Phe Arg Phe Glu Leu Pro Arg Thr
155 160 165
Arg Ile Thr Asn Ile Leu Lys Arg Ser Gly Glu Gln Ile Trp Ser
170 175 180
Val Ser Ile Phe Leu Leu Phe Phe Leu Leu Tyr Gly Ile Leu
185 190 195
Gly Val Gln Met Phe Gly Thr Phe Thr Tyr His Cys Val Val Asn

Asp Thr Lys Pro	200	Asn Val Thr Trp	205	Ser Leu Ala Ile	210
215	Gly	220	Asn	Pro	225
Asp Thr His Cys	230	Pro Glu Leu Glu	235	Gly Tyr Gln Cys	240
245	Ser	250	Asp	Leu Gly Leu Ser	255
Pro Gly Phe Lys	260	Met Asp Leu Glu	265	Ile Gly Thr Ser	270
275	Tyr	280	Glu	Phe Leu	285
Gln Glu Leu Gly	290	Ser Gly Phe Asn	295	Gly Trp Val Phe	300
305	Glu	310	Arg	Tyr Phe	315
Phe Thr Val Tyr	320	Ala Ala Ser Gln	325	Trp Val Phe	330
335	Ile	340	Ser	Thr Thr Ser	345
Met Tyr Arg Ala	350	Asp Ser Phe Pro	355	Thr Thr Ser	360
365	Leu	370	Ala	Thr Thr Ser	375
Tyr Phe Ile Thr	380	Ile Phe Phe Leu	385	Thr Thr Ser	390
395	Val	400	Val	Thr Thr Ser	405
Val Phe Ile Ala	410	Ile Ile Glu Thr	415	Thr Thr Ser	420
425	Gln	430	Asp	Thr Thr Ser	435
Gln Phe Gln Gln	440	Met Trp Gly Ser Arg	445	Thr Thr Ser	450
455	Ala	460	Ala	Thr Thr Ser	465
Ala Thr Thr Gln	470	Met Phe His Glu Asp	475	Thr Thr Ser	480
485	Leu	490	Val	Thr Thr Ser	495
Leu Val Ala Val	500	Asp Val Asn Lys Pro	505	Thr Thr Ser	510
515	Cys	520	Val	Thr Thr Ser	525
530	Leu	535	Ala	Thr Thr Ser	540
Cys Leu Gln Lys	545	Met Met Arg Ser Ser	550	Thr Thr Ser	555
560	Leu	565	Ala	Thr Thr Ser	570
Leu Ser Met Val	575	Thr Val Asp Val Ile	580	Thr Thr Ser	585
590	Tyr	595	Val	Thr Thr Ser	600
Tyr Lys Gly Glu	605	Asn Phe Arg Arg Gln	610	Thr Thr Ser	615
620	Ala	625	Leu	Thr Thr Ser	625
Ala Glu Val Ala	635	Phe Thr Val Leu Phe	640	Thr Thr Ser	645
650	Lys	655	Val	Thr Thr Ser	655
Lys Ile Trp Cys	665	Leu Gly Phe Thr Gly	670	Thr Thr Ser	665
680	His	685	Ile	Thr Thr Ser	675
His Lys Phe Glu	695	Leu Leu Val Ile	700	Thr Thr Ser	685
455	Tyr	460	Gly	Thr Thr Ser	690
470	Pro	475	Thr	Thr Thr Ser	695
485	Arg	490	Pro	Thr Thr Ser	700
495	Val	500	Ala	Thr Thr Ser	705
505	Val	510	Leu	Thr Thr Ser	
515	Val	520	Leu	Thr Thr Ser	
525	Gln	530	Met	Thr Thr Ser	
535	Pro	540	Arg	Thr Thr Ser	
545	Trp	550	Ala	Thr Thr Ser	
555	Trp	560	Val	Thr Thr Ser	
565	Trp	570	Ala	Thr Thr Ser	
575	Ala	580	Thr	Thr Thr Ser	
585	Ala	590	Thr	Thr Thr Ser	
595	Asn	600	Leu	Thr Thr Ser	
605	Asn	610	Leu	Thr Thr Ser	
615	Gln	620	Leu	Thr Thr Ser	
625	Gln	630	Leu	Thr Thr Ser	
635	Leu	640	Leu	Thr Thr Ser	
645	Ile	650	Leu	Thr Thr Ser	
655	Ser	660	Leu	Thr Thr Ser	
665	Cys	670	Leu	Thr Thr Ser	
675	Cys	680	Leu	Thr Thr Ser	
685	His	690	Leu	Thr Thr Ser	
695	His	700	Leu	Thr Thr Ser	

Lys	Leu	Arg	Lys	Ser	Val	Phe	Ser	Ile	Arg	Ala	Arg	Asn	Leu	Leu
				710					715					720
Glu	Lys	Glu	Thr	Ala	Val	Thr	Lys	Ile	Leu	Arg	Ala	Cys	Thr	Arg
				725					730					735
Gln	Arg	Met	Leu	Ser	Gly	Ser	Phe	Glu	Gly	Gln	Pro	Ala	Lys	Glu
				740					745					750
Arg	Ser	Ile	Leu	Ser	Val	Gln	His	His	Ile	Arg	Gln	Glu	Arg	Arg
				755					760					765
Ser	Leu	Arg	His	Gly	Ser	Asn	Ser	Gln	Arg	Ile	Ser	Arg	Gly	Lys
				770					775					780
Ser	Leu	Glu	Thr	Leu	Thr	Gln	Asp	His	Cys	Asn	Thr	Val	Ile	Tyr
				785					790					795
Arg	Asn	Ala	Gln	Arg	Glu	Val	Ser	Glu	Ile	Lys	Met	Ile	Gln	Glu
				800					805					810
Lys	Lys	Glu	Leu	Ala	Glu	Met	Leu	Gln	Gly	Lys	Cys	Lys	Lys	Glu
				815					820					825
Leu	Arg	Glu	Ser	His	Pro	Tyr	Phe	Asp	Lys	Pro	Leu	Phe	Ile	Val
				830					835					840
Gly	Arg	Glu	His	Arg	Phe	Arg	Asn	Phe	Cys	Arg	Val	Val	Val	Arg
				845					850					855
Ala	Arg	Phe	Asn	Ala	Ser	Lys	Thr	Asp	Pro	Val	Thr	Gly	Ala	Val
				860					865					870
Lys	Asn	Thr	Lys	Tyr	His	Leu	Leu	Tyr	Asp	Leu	Leu	Gly	Leu	Val
				875					880					885
Thr	Tyr	Leu	Asp	Trp	Val	Met	Ile	Ile	Val	Thr	Ser	Asp	Ser	Cys
				890					895					900
Ile	Ser	Met	Met	Phe	Glu	Ser	Pro	Phe	Arg	Arg	Val	Met	His	Ala
				905					910					915
Pro	Thr	Leu	Gln	Ile	Ala	Glu	Tyr	Val	Phe	Val	Ile	Phe	Met	Ser
				920					925					930
Ile	Glu	Leu	Asn	Leu	Lys	Ile	Met	Ala	Asp	Gly	Leu	Phe	Phe	Thr
				935					940					945
Pro	Thr	Ala	Val	Ile	Arg	Asp	Phe	Gly	Gly	Val	Met	Asp	Ile	Phe
				950					955					960
Ile	Tyr	Leu	Val	Ser	Leu	Ile	Phe	Leu	Cys	Trp	Met	Pro	Gln	Asn
				965					970					975
Val	Pro	Ala	Glu	Ser	Gly	Ala	Gln	Leu	Leu	Met	Val	Leu	Arg	Cys
				980					985					990
Leu	Arg	Pro	Leu	Arg	Ile	Phe	Lys	Leu	Val	Pro	Gln	Met	Arg	Lys
				995					1000					1005
Val	Val	Arg	Glu	Leu	Phe	Ser	Gly	Phe	Lys	Glu	Ile	Phe	Leu	Val
				1010					1015					1020
Ser	Ile	Leu	Leu	Leu	Thr	Leu	Met	Leu	Val	Phe	Ala	Ser	Phe	Gly
				1025					1030					1035
Val	Gln	Leu	Phe	Ala	Gly	Lys	Leu	Ala	Lys	Cys	Asn	Asp	Pro	Asn
				1040					1045					1050
Ile	Ile	Arg	Arg	Glu	Asp	Cys	Asn	Gly	Ile	Phe	Arg	Ile	Asn	Val
				1055					1060					1065
Ser	Val	Ser	Lys	Asn	Leu	Asn	Leu	Lys	Leu	Arg	Pro	Gly	Glu	Lys
				1070					1075					1080
Lys	Pro	Gly	Phe	Trp	Val	Pro	Arg	Val	Trp	Ala	Asn	Pro	Arg	Asn
				1085					1090					1095
Phe	Asn	Phe	Asp	Asn	Val	Gly	Asn	Ala	Met	Leu	Ala	Leu	Phe	Glu
				1100					1105					1110
Val	Leu	Ser	Leu	Lys	Gly	Trp	Val	Glu	Val	Arg	Asp	Val	Ile	Ile
				1115					1120					1125
His	Arg	Val	Gly	Pro	Ile	His	Gly	Ile	Tyr	Ile	His	Val	Phe	Val
				1130					1135					1140
Phe	Leu	Gly	Cys	Met	Ile	Gly	Leu	Thr	Leu	Phe	Val	Gly	Val	Val
				1145					1150					1155
Ile	Ala	Asn	Phe	Asn	Glu	Asn	Lys	Gly	Thr	Ala	Leu	Leu	Thr	Val
				1160					1165					1170
Asp	Gln	Arg	Arg	Trp	Glu	Asp	Leu	Lys	Ser	Arg	Leu	Lys	Ile	Ala
				1175					1180					1185
Gln	Pro	Leu	His	Leu	Pro	Pro	Arg	Pro	Asp	Asn	Asp	Gly	Phe	Arg
				1190					1195					1200
Ala	Lys	Met	Tyr	Asp	Ile	Thr	Gln	His	Pro	Phe	Phe	Lys	Arg	Thr

Ile Ala Leu Leu Val	1205	Leu Ala Gln Ser Val	1210	Leu Leu Ser Val Lys	1215
Trp Asp Val Glu Asp	1220	Pro Val Thr Val Pro	1225	Leu Ala Thr Met Ser	1230
Val Val Phe Thr Phe	1235	Ile Phe Val Leu Glu	1240	Val Thr Met Lys Ile	1245
Ile Ala Met Ser Pro	1250	Ala Gly Phe Trp Gln	1255	Ser Arg Arg Asn Arg	1260
Tyr Asp Leu Leu Val	1265	Thr Ser Leu Gly Val	1270	Val Trp Val Val Leu	1275
His Phe Ala Leu Leu	1280	Asn Ala Tyr Thr Tyr	1285	Met Met Gly Ala Cys	1290
Val Ile Val Phe Arg	1295	Phe Phe Ser Ile Cys	1300	Gly Lys His Val Thr	1305
Leu Lys Met Leu Leu	1310	Leu Thr Val Val Val	1315	Ser Met Tyr Lys Ser	1320
Phe Phe Ile Ile Val	1325	Gly Met Phe Leu Leu	1330	Leu Leu Cys Tyr Ala	1335
Phe Ala Gly Val Val	1340	Leu Phe Gly Thr Val	1345	Lys Tyr Gly Glu Asn	1350
Ile Asn Arg His Ala	1355	Asn Phe Ser Ser Ala	1360	Gly Lys Ala Ile Thr	1365
Val Leu Phe Arg Ile	1370	Val Thr Gly Glu Asp	1375	Trp Asn Lys Ile Met	1380
His Asp Cys Met Val	1385	Gln Pro Pro Phe Cys	1390	Thr Pro Asp Glu Phe	1395
Thr Tyr Trp Ala Thr	1400	Asp Cys Gly Asn Tyr	1405	Ala Gly Ala Leu Met	1410
Tyr Phe Cys Ser Phe	1415	Tyr Val Ile Ile Ala	1420	Tyr Ile Met Leu Asn	1425
Leu Leu Val Ala Ile	1430	Ile Val Glu Asn Phe	1435	Ser Leu Ile Tyr Ser	1440
Thr Glu Glu Asp Gln	1445	Leu Leu Ser Tyr Asn	1450	Asp Leu Arg His Phe	1455
Gln Ile Ile Trp Asn	1460	Met Val Asp Asp Lys	1465	Arg Glu Val Phe Pro	1470
Thr Phe Arg Val Lys	1475	Phe Leu Leu Arg Leu	1480	Leu Arg Gly Arg Leu	1485
Glu Val Asp Leu Asp	1490	Lys Asp Lys Leu Leu	1495	Phe Lys His Met Cys	1500
Tyr Glu Met Glu Arg	1505	Leu His Asn Gly Gly	1510	Asp Val Thr Phe His	1515
Asp Val Leu Ser Met	1520	Leu Ser Tyr Arg Ser	1525	Val Asp Ile Arg Lys	1530
Ser Leu Gln Leu Glu	1535	Glu Leu Leu Ala Arg	1540	Glu Gln Leu Glu Tyr	1545
Thr Ile Glu Glu Glu	1550	Val Ala Lys Gln Thr	1555	Ile Arg Met Trp Leu	1560
Lys Lys Cys Leu Lys	1565	Arg Ile Arg Ala Lys	1570	Gln Gln Gln Ser Cys	1575
Ser Ile Ile His Ser	1580	Leu Arg Glu Ser Gln	1585	Gln Gln Glu Leu Ser	1590
Arg Phe Leu Asn Pro	1595	Pro Ser Ile Glu Thr	1600	Thr Gln Pro Ser Glu	1605
Asp Thr Asn Ala Asn	1610	Ser Gln Asp Asn Ser	1615	Met Gln Pro Glu Thr	1620
Ser Ser Gln Gln Gln	1625	Leu Leu Ser Pro Thr	1630	Leu Ser Asp Arg Gly	1635
Gly Ser Arg Gln Asp	1640	Ala Ala Asp Ala Gly	1645	Lys Pro Gln Arg Lys	1650
Phe Gly Gln Trp Arg	1655	Leu Pro Ser Ala Pro	1660	Lys Pro Ile Ser His	1665
Ser Val Ser Ser Val	1670	Asn Leu Arg Phe Gly	1675	Gly Arg Thr Thr Met	1680
Lys Ser Val Val Cys	1685	Lys Met Asn Pro Met	1690	Thr Asp Ala Ala Ser	1695
	1700		1705		1710

Cys Gly Ser Glu Val Lys Lys Trp Trp Thr Arg Gln Leu Thr Val
 1715 1720 1725
 Glu Ser Asp Glu Ser Gly Asp Asp Leu Leu Asp Ile
 1730 1735

<210> 29
 <211> 547
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 168827CD1

<400> 29
 Met Ala Phe Gln Asp Leu Leu Asp Gln Val Gly Gly Leu Gly Arg
 1 5 10 15
 Phe Gln Ile Leu Gln Met Val Phe Leu Ile Met Phe Asn Val Ile
 20 25 30
 Val Tyr His Gln Thr Gln Leu Glu Asn Phe Ala Ala Phe Ile Leu
 35 40 45
 Asp His Arg Cys Trp Val His Ile Leu Asp Asn Asp Thr Ile Pro
 50 55 60
 Asp Asn Asp Pro Gly Thr Leu Ser Gln Asp Ala Leu Leu Arg Ile
 65 70 75
 Ser Ile Pro Phe Asp Ser Asn Leu Arg Pro Glu Lys Cys Arg Arg
 80 85 90
 Phe Val His Pro Gln Trp Lys Leu Ile His Leu Asn Gly Thr Phe
 95 100 105
 Pro Asn Thr Ser Glu Pro Asp Thr Glu Pro Cys Val Asp Gly Trp
 110 115 120
 Val Tyr Asp Gln Ser Ser Phe Pro Ser Thr Ile Val Thr Lys Trp
 125 130 135
 Asp Leu Val Cys Glu Ser Gln Pro Leu Asn Ser Val Ala Lys Phe
 140 145 150
 Leu Phe Met Ala Gly Met Met Val Gly Gly Asn Leu Tyr Gly His
 155 160 165
 Leu Ser Asp Arg Phe Gly Arg Lys Phe Val Leu Arg Trp Ser Tyr
 170 175 180
 Leu Gln Leu Ala Ile Val Gly Thr Cys Ala Phe Ala Pro Thr
 185 190 195
 Ile Leu Val Tyr Cys Ser Leu Arg Phe Leu Ala Gly Ala Ala Thr
 200 205 210
 Phe Ser Ile Ile Val Asn Thr Val Leu Leu Ile Val Glu Trp Ile
 215 220 225
 Thr His Gln Phe Cys Ala Met Ala Leu Thr Leu Thr Leu Cys Ala
 230 235 240
 Ala Ser Ile Gly His Ile Thr Leu Gly Ser Leu Ala Phe Val Ile
 245 250 255
 Arg Asp Gln Cys Ile Leu Gln Leu Val Met Ser Ala Pro Cys Phe
 260 265 270
 Val Phe Phe Leu Phe Ser Arg Trp Leu Ala Glu Ser Ala Arg Trp
 275 280 285
 Leu Ile Ile Asn Asn Lys Pro Glu Glu Gly Leu Lys Glu Leu Thr
 290 295 300
 Lys Ala Ala His Arg Asn Gly Met Lys Asn Ala Glu Asp Ile Leu
 305 310 315
 Thr Met Glu Val Leu Lys Ser Thr Met Lys Gln Glu Leu Glu Ala
 320 325 330
 Ala Gln Lys Lys His Ser Leu Cys Glu Leu Leu Arg Ile Pro Asn
 335 340 345
 Ile Cys Lys Arg Ile Cys Phe Leu Ser Phe Val Arg Phe Ala Ser
 350 355 360
 Thr Ile Pro Phe Trp Gly Leu Thr Leu His Leu Gln His Leu Gly
 365 370 375
 Asn Asn Val Phe Leu Leu Gln Thr Leu Phe Gly Ala Val Thr Leu
 380 385 390

Leu	Ala	Asn	Cys	Val	Ala	Pro	Trp	Ala	Leu	Asn	His	Met	Ser	Arg
				395					400					405
Arg	Leu	Ser	Gln	Met	Leu	Leu	Met	Phe	Leu	Leu	Ala	Thr	Cys	Leu
				410					415					420
Leu	Ala	Ile	Ile	Phe	Val	Pro	Gln	Glu	Met	Gln	Thr	Leu	Arg	Val
				425					430					435
Val	Leu	Ala	Thr	Leu	Gly	Val	Gly	Ala	Ala	Ser	Leu	Gly	Ile	Thr
				440					445					450
Cys	Ser	Thr	Ala	Gln	Glu	Asn	Glu	Leu	Ile	Pro	Ser	Ile	Ile	Arg
				455					460					465
Gly	Arg	Ala	Thr	Gly	Ile	Thr	Gly	Asn	Phe	Ala	Asn	Ile	Gly	Gly
				470					475					480
Ala	Leu	Ala	Ser	Leu	Met	Met	Ile	Leu	Ser	Ile	Tyr	Ser	Arg	Pro
				485					490					495
Leu	Pro	Trp	Ile	Ile	Tyr	Gly	Val	Phe	Ala	Ile	Leu	Ser	Gly	Leu
				500					505					510
Val	Val	Leu	Leu	Leu	Pro	Glu	Thr	Arg	Asn	Gln	Pro	Leu	Leu	Asp
				515					520					525
Ser	Ile	Gln	Asp	Val	Glu	Asn	Glu	Gly	Val	Asn	Ser	Leu	Ala	Ala
				530					535					540
Pro	Gln	Arg	Ser	Ser	Val	Leu								
				545										

<210> 30

<211> 547

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472734CD1

<400> 30

Met	Gly	Phe	Asp	Val	Leu	Leu	Asp	Gln	Val	Gly	Gly	Met	Gly	Arg
1				5					10					15
Phe	Gln	Ile	Cys	Leu	Ile	Ala	Phe	Phe	Cys	Ile	Thr	Asn	Ile	Leu
				20					25					30
Leu	Phe	Pro	Asn	Ile	Val	Leu	Glu	Asn	Phe	Thr	Ala	Phe	Thr	Pro
				35					40					45
Ser	His	Arg	Cys	Trp	Val	Pro	Leu	Leu	Asp	Asn	Asp	Thr	Val	Ser
				50					55					60
Asp	Asn	Asp	Thr	Gly	Thr	Leu	Ser	Lys	Asp	Asp	Leu	Leu	Arg	Ile
				65					70					75
Ser	Ile	Pro	Leu	Asp	Ser	Asn	Leu	Arg	Pro	Gln	Lys	Cys	Gln	Arg
				80					85					90
Phe	Ile	His	Pro	Gln	Trp	Gln	Leu	Leu	His	Leu	Asn	Gly	Thr	Phe
				95					100					105
Pro	Asn	Thr	Asn	Glu	Pro	Asp	Thr	Glu	Pro	Cys	Val	Asp	Gly	Trp
				110					115					120
Val	Tyr	Asp	Arg	Ser	Ser	Phe	Leu	Ser	Thr	Ile	Val	Thr	Glu	Trp
				125					130					135
Asp	Leu	Val	Cys	Glu	Ser	Gln	Ser	Leu	Lys	Ser	Met	Val	Gln	Ser
				140					145					150
Leu	Phe	Met	Ala	Gly	Ser	Leu	Leu	Gly	Gly	Leu	Ile	Tyr	Gly	His
				155					160					165
Leu	Ser	Asp	Arg	Phe	Gly	Arg	Lys	Phe	Val	Leu	Arg	Trp	Ser	Tyr
				170					175					180
Leu	Gln	Leu	Ala	Ile	Val	Gly	Thr	Cys	Ala	Ala	Phe	Ala	Pro	Thr
				185					190					195
Ile	Leu	Val	Tyr	Cys	Ser	Leu	Arg	Phe	Leu	Ala	Gly	Ala	Ala	Thr
				200					205					210
Phe	Ser	Ile	Ile	Val	Asn	Thr	Val	Leu	Leu	Ile	Val	Glu	Trp	Ile
				215					220					225
Thr	His	Gln	Phe	Cys	Ala	Met	Ala	Leu	Thr	Leu	Thr	Leu	Cys	Ala
				230					235					240
Ala	Ser	Ile	Gly	His	Ile	Thr	Leu	Gly	Ser	Leu	Ala	Phe	Val	Ile
				245					250					255

Arg	Asp	Gln	Cys	Ile	Leu	Gln	Leu	Val	Met	Ser	Ala	Pro	Cys	Phe
				260					265					270
Val	Phe	Phe	Leu	Phe	Ser	Arg	Trp	Leu	Ala	Glu	Ser	Ala	Arg	Trp
				275					280					285
Leu	Ile	Ile	Asn	Asn	Lys	Pro	Glu	Glu	Gly	Leu	Lys	Glu	Leu	Arg
				290					295					300
Lys	Ala	Ala	His	Arg	Asn	Gly	Met	Lys	Asn	Ala	Glu	Asp	Ile	Leu
				305					310					315
Thr	Met	Glu	Val	Leu	Lys	Ser	Thr	Met	Lys	Gln	Glu	Leu	Glu	Ala
				320					325					330
Ala	Gln	Lys	Lys	His	Ser	Leu	Cys	Glu	Leu	Leu	Arg	Ile	Pro	Asn
				335					340					345
Ile	Cys	Lys	Arg	Ile	Cys	Phe	Leu	Ser	Phe	Val	Arg	Phe	Ala	Ser
				350					355					360
Thr	Ile	Pro	Phe	Trp	Gly	Leu	Thr	Leu	His	Leu	Gln	His	Leu	Gly
				365					370					375
Asn	Asn	Val	Phe	Leu	Leu	Gln	Thr	Leu	Phe	Gly	Ala	Val	Thr	Leu
				380					385					390
Leu	Ala	Asn	Cys	Val	Ala	Pro	Trp	Ala	Leu	Asn	His	Met	Ser	Arg
				395					400					405
Arg	Leu	Ser	Gln	Met	Leu	Leu	Met	Phe	Leu	Leu	Ala	Thr	Cys	Leu
				410					415					420
Leu	Ala	Ile	Ile	Phe	Val	Pro	Gln	Glu	Met	Gln	Thr	Leu	Arg	Val
				425					430					435
Val	Leu	Ala	Thr	Leu	Gly	Val	Gly	Ala	Ala	Ser	Leu	Gly	Ile	Thr
				440					445					450
Cys	Ser	Thr	Ala	Gln	Glu	Asn	Glu	Leu	Ile	Pro	Ser	Ile	Ile	Arg
				455					460					465
Gly	Arg	Ala	Thr	Gly	Ile	Thr	Gly	Asn	Phe	Ala	Asn	Ile	Gly	Gly
				470					475					480
Ala	Leu	Ala	Ser	Leu	Met	Met	Ile	Leu	Ser	Ile	Tyr	Ser	Arg	Pro
				485					490					495
Leu	Pro	Trp	Ile	Ile	Tyr	Gly	Val	Phe	Ala	Ile	Leu	Ser	Gly	Leu
				500					505					510
Val	Val	Leu	Leu	Leu	Pro	Glu	Thr	Arg	Asn	Gln	Pro	Leu	Leu	Asp
				515					520					525
Ser	Ile	Gln	Asp	Val	Glu	Asn	Glu	Gly	Val	Asn	Ser	Leu	Ala	Ala
				530					535					540
Pro	Gln	Arg	Ser	Ser	Val	Leu								
				545										

<210> 31
 <211> 988
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7473473CD1

<400> 31
Met Pro Gly Gly Lys Arg Gly Leu Val Ala Pro Gln Asn Thr Phe
1 5 10 15
Leu Glu Asn Ile Val Arg Arg Ser Ser Glu Ser Ser Phe Leu Leu
20 25 30
Gly Asn Ala Gln Ile Val Asp Trp Pro Val Val Tyr Ser Asn Asp
35 40 45
Gly Phe Cys Lys Leu Ser Gly Tyr His Arg Ala Asp Val Met Gln
50 55 60
Lys Ser Ser Thr Cys Ser Phe Met Tyr Gly Glu Leu Thr Asp Lys
65 70 75
Lys Thr Ile Glu Lys Val Arg Gln Thr Phe Asp Asn Tyr Glu Ser
80 85 90
Asn Cys Phe Glu Val Leu Leu Tyr Lys Lys Asn Arg Thr Pro Val
95 100 105
Trp Phe Tyr Met Gln Ile Ala Pro Ile Arg Asn Glu His Glu Lys
110 115 120

Val	Val	Leu	Phe	Leu	Cys	Thr	Phe	Lys	Asp	Ile	Thr	Leu	Phe	Lys
				125					130					135
Gln	Pro	Ile	Glu	Asp	Asp	Ser	Thr	Lys	Gly	Trp	Thr	Lys	Phe	Ala
				140					145					150
Arg	Leu	Thr	Arg	Ala	Leu	Thr	Asn	Ser	Arg	Ser	Val	Leu	Gln	Gln
				155					160					165
Leu	Thr	Pro	Met	Asn	Lys	Thr	Glu	Val	Val	His	Lys	His	Ser	Arg
				170					175					180
Leu	Ala	Glu	Val	Leu	Gln	Leu	Gly	Ser	Asp	Ile	Leu	Pro	Gln	Tyr
				185					190					195
Lys	Gln	Glu	Ala	Pro	Lys	Thr	Pro	Pro	His	Ile	Ile	Leu	His	Tyr
				200					205					210
Cys	Ala	Phe	Lys	Thr	Thr	Trp	Asp	Trp	Val	Ile	Leu	Ile	Leu	Thr
				215					220					225
Phe	Tyr	Thr	Ala	Ile	Met	Val	Pro	Tyr	Asn	Val	Ser	Phe	Lys	Thr
				230					235					240
Lys	Gln	Asn	Asn	Ile	Ala	Trp	Leu	Val	Leu	Asp	Ser	Val	Val	Asp
				245					250					255
Val	Ile	Phe	Leu	Val	Asp	Ile	Val	Leu	Asn	Phe	His	Thr	Thr	Phe
				260					265					270
Val	Gly	Pro	Gly	Gly	Glu	Val	Ile	Ser	Asp	Pro	Lys	Leu	Ile	Arg
				275					280					285
Met	Asn	Tyr	Leu	Lys	Thr	Trp	Phe	Val	Ile	Asp	Leu	Leu	Ser	Cys
				290					295					300
Leu	Pro	Tyr	Asp	Ile	Ile	Asn	Ala	Phe	Glu	Asn	Val	Asp	Glu	Gly
				305					310					315
Ile	Ser	Ser	Leu	Phe	Ser	Ser	Leu	Lys	Val	Val	Arg	Leu	Leu	Arg
				320					325					330
Leu	Gly	Arg	Val	Ala	Arg	Lys	Leu	Asp	His	Tyr	Leu	Glu	Tyr	Gly
				335					340					345
Ala	Ala	Val	Leu	Val	Leu	Leu	Val	Cys	Val	Phe	Gly	Leu	Val	Ala
				350					355					360
His	Trp	Leu	Ala	Cys	Ile	Trp	Tyr	Ser	Ile	Gly	Asp	Tyr	Glu	Val
				365					370					375
Ile	Asp	Glu	Val	Thr	Asn	Thr	Ile	Gln	Ile	Asp	Ser	Trp	Leu	Tyr
				380					385					390
Gln	Leu	Ala	Leu	Ser	Ile	Gly	Thr	Pro	Tyr	Arg	Tyr	Asn	Thr	Ser
				395					400					405
Ala	Gly	Ile	Trp	Glu	Gly	Gly	Pro	Ser	Lys	Asp	Ser	Leu	Tyr	Val
				410					415					420
Ser	Ser	Leu	Tyr	Phe	Thr	Met	Thr	Ser	Leu	Thr	Thr	Ile	Gly	Phe
				425					430					435
Gly	Asn	Ile	Ala	Pro	Thr	Thr	Asp	Val	Glu	Lys	Met	Phe	Ser	Val
				440					445					450
Ala	Met	Met	Met	Val	Gly	Ala	Leu	Leu	Tyr	Ala	Thr	Ile	Phe	Gly
				455					460					465
Asn	Val	Thr	Thr	Ile	Phe	Gln	Gln	Met	Tyr	Ala	Asn	Thr	Asn	Arg
				470					475					480
Tyr	His	Glu	Met	Leu	Asn	Asn	Val	Arg	Asp	Phe	Leu	Lys	Leu	Tyr
				485					490					495
Gln	Val	Pro	Lys	Gly	Leu	Ser	Glu	Arg	Val	Met	Asp	Tyr	Ile	Val
				500					505					510
Ser	Thr	Trp	Ser	Met	Ser	Lys	Gly	Ile	Asp	Thr	Glu	Lys	Val	Leu
				515					520					525
Ser	Ile	Cys	Pro	Lys	Asp	Met	Arg	Ala	Asp	Ile	Cys	Val	His	Leu
				530					535					540
Asn	Arg	Lys	Val	Phe	Asn	Glu	His	Pro	Ala	Phe	Arg	Leu	Ala	Ser
				545					550					555
Asp	Gly	Cys	Leu	Arg	Ala	Leu	Ala	Val	Glu	Phe	Gln	Thr	Ile	His
				560					565					570
Cys	Ala	Pro	Gly	Asp	Leu	Ile	Tyr	His	Ala	Gly	Glu	Ser	Val	Asp
				575					580					585
Ala	Leu	Cys	Phe	Val	Val	Ser	Gly	Ser	Leu	Glu	Val	Ile	Gln	Asp
				590					595					600
Asp	Glu	Val	Val	Ala	Ile	Leu	Gly	Lys	Gly	Asp	Val	Phe	Gly	Asp
				605					610					615
Ile	Phe	Trp	Lys	Glu	Thr	Thr	Leu	Ala	His	Ala	Cys	Ala	Asn	Val

Arg	Ala	Leu	Thr	620	Tyr	Cys	Asp	Leu	His	625	Ile	Lys	Arg	Glu	630	Ala	
Leu	Leu	Lys	Val	635	Leu	Asp	Phe	Tyr	Thr	640	Ala	Phe	Ala	Asn	Ser	645	Phe
Ser	Arg	Asn	Leu	650	Thr	Leu	Thr	Cys	Asn	655	Leu	Arg	Lys	Arg	Ile	660	Ile
Phe	Arg	Lys	Ile	665	Ser	Asp	Val	Lys	Lys	670	Glu	Glu	Glu	Glu	Arg	675	Leu
Arg	Gln	Lys	Asn	680	Glu	Val	Thr	Leu	Ser	685	Ile	Pro	Val	Asp	His	690	Pro
Val	Arg	Lys	Leu	695	Phe	Gln	Lys	Phe	Lys	700	Gln	Gln	Lys	Glu	Leu	705	Arg
Asn	Gln	Gly	Ser	710	Thr	Gln	Gly	Asp	Pro	715	Glu	Arg	Asn	Gln	Leu	720	Gln
Val	Glu	Ser	Arg	725	Ser	Leu	Gln	Asn	Gly	730	Ala	Ser	Ile	Thr	Gly	735	Thr
Ser	Val	Val	Thr	740	Val	Ser	Gln	Ile	Thr	745	Pro	Ile	Gln	Thr	Ser	750	Leu
Ala	Tyr	Val	Lys	755	Thr	Ser	Glu	Ser	Leu	760	Lys	Gln	Asn	Asn	Arg	765	Asp
Ala	Met	Glu	Leu	770	Lys	Pro	Asn	Gly	Gly	775	Ala	Asp	Gln	Lys	Cys	780	Leu
Lys	Val	Asn	Ser	785	Pro	Ile	Arg	Met	Lys	790	Asn	Gly	Asn	Gly	Lys	795	Gly
Trp	Leu	Arg	Leu	800	Lys	Asn	Asn	Met	Gly	805	Ala	His	Glu	Glu	Lys	810	Lys
Glu	Asp	Trp	Asn	815	Asn	Val	Thr	Lys	Ala	820	Glu	Ser	Met	Gly	Leu	825	Leu
Ser	Glu	Asp	Pro	830	Lys	Ser	Ser	Asp	Ser	835	Glu	Asn	Ser	Val	Thr	840	Lys
Asn	Pro	Leu	Arg	845	Lys	Thr	Asp	Ser	Cys	850	Asp	Ser	Gly	Ile	Thr	855	Lys
Ser	Asp	Leu	Arg	860	Leu	Asp	Lys	Ala	Gly	865	Glu	Ala	Arg	Ser	Pro	870	Leu
Glu	His	Ser	Pro	875	Ile	Gln	Ala	Asp	Ala	880	Lys	His	Pro	Phe	Tyr	885	Pro
Ile	Pro	Glu	Gln	890	Ala	Leu	Gln	Thr	Thr	895	Leu	Gln	Glu	Val	Lys	900	His
Glu	Leu	Lys	Glu	905	Asp	Ile	Gln	Leu	Leu	910	Ser	Cys	Arg	Met	Thr	915	Ala
Leu	Glu	Lys	Gln	920	Val	Ala	Glu	Ile	Leu	925	Lys	Ile	Leu	Ser	Glu	930	Lys
Ser	Val	Pro	Gln	935	Ala	Ser	Ser	Pro	Lys	940	Ser	Gln	Met	Pro	Leu	945	Gln
Val	Pro	Pro	Gln	950	Ile	Pro	Cys	Gln	Asp	955	Ile	Phe	Ser	Val	Ser	960	Arg
Pro	Glu	Ser	Pro	965	Glu	Ser	Asp	Lys	Asp	970	Glu	Ile	His	Phe		975	
				980						985							

<210> 32
 <211> 533
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7477725CD1

<400> 32
 Met Ala Phe Glu Glu Leu Leu Ser Gln Val Gly Gly Leu Gly Arg
 1 5 10 15
 Phe Gln Met Leu His Leu Val Phe Ile Leu Pro Ser Leu Met Leu
 20 25 30
 Leu Ile Pro His Leu Leu Glu Asn Phe Ala Ala Ala Ile Pro
 35 40 45
 Gly His Arg Cys Trp Val His Met Leu Asp Asn Asn Thr Gly Ser

Gly Asn Glu Thr	50	Gly Ile Leu Ser Glu	55	Ala Leu Leu Arg	60
	65		70		75
Ser Ile Pro Leu Asp	80	Ser Asn Leu Arg	85	Pro Glu Lys Cys Arg	90
	95		100		105
Phe Val His Pro Gln	110	Trp Gln Leu Leu	115	His Leu Asn Gly Thr	120
	125		130		135
His Ser Thr Ser Glu	140	Ala Asp Thr Glu	145	Pro Cys Val Asp Gly	150
	155		160		165
Val Tyr Asp Gln Ser	170	Tyr Phe Pro Ser	175	Thr Ile Val Thr Lys	180
	185		190		195
Asp Leu Val Cys Asp	200	Tyr Gln Ser Leu	205	Lys Ser Val Val Gln	210
	215		220		225
Leu Leu Leu Thr Gly	230	Met Leu Val Gly	235	Gly Ile Ile Gly Gly	240
	245		250		255
Val Ser Asp Arg Phe	260	Gly Arg Arg Phe	265	Ile Leu Arg Trp Cys	270
	275		280		285
Leu Gln Leu Ala Ile	290	Thr Asp Thr Cys	295	Ala Ala Phe Ala Pro	300
	305		310		315
Phe Pro Val Tyr Cys	320	Val Leu Arg Phe	325	Leu Ala Gly Phe Ser	330
	335		340		345
Met Ile Ile Ile Ser	350	Asn Asn Ser Leu	355	Pro Ile Thr Glu Trp	360
	365		370		375
Arg Pro Asn Ser Lys	380	Ala Leu Val Val	385	Ile Leu Ser Ser Gly	390
	395		400		405
Leu Ser Ile Gly Gln	410	Ile Ile Leu Gly	415	Leu Ala Tyr Val Phe	420
	425		430		435
Arg Asp Trp Gln Thr	440	Leu His Val Val	445	Ala Ser Val Pro Phe	450
	455		460		465
Val Phe Phe Leu Leu	470	Ser Arg Trp Leu	475	Val Glu Ser Ala Arg	480
	485		490		495
Leu Ile Ile Thr Asn	500	Lys Leu Asp Glu	505	Gly Leu Lys Ala Leu	510
	515		520		525
Lys Val Ala Arg Thr	530	Asn Gly Ile Lys		Asn Ala Glu Glu Thr	
Asn Ile Glu Val Val		Arg Ser Thr Met		Gln Glu Leu Asp Ala	
Ala Gln Thr Lys Thr		Thr Val Cys Asp		Leu Phe Arg Asn Pro	
Met Arg Lys Arg Ile		Cys Ile Leu Val		Phe Leu Arg Phe Ala	
Thr Ile Pro Phe Tyr		Gly Thr Met Val		Asn Leu Gln His Val	
Ser Asn Ile Phe Leu		Leu Gln Val Leu		Tyr Gly Ala Val Ala	
Ile Val Arg Cys Leu		Ala Leu Leu Thr		Leu Asn His Met Gly	
Arg Ile Ser Gln Ile		Leu Phe Met Phe		Leu Val Gly Leu Ser	
Leu Ala Asn Thr Phe		Val Pro Lys Glu		Met Gln Thr Leu Arg	
Ala Leu Ala Cys Leu		Gly Ile Gly Cys		Ser Ala Ala Thr Phe	
Ser Val Ala Val His		Phe Ile Glu Leu		Ile Pro Thr Val Leu	
Ala Arg Ala Ser Gly		Ile Asp Leu Thr		Ala Ser Arg Ile Gly	
Ala Leu Ala Pro Leu		Leu Met Thr Leu		Thr Val Phe Phe Thr	
Leu Pro Trp Ile Ile		Tyr Gly Ile Phe		Pro Ile Ile Gly Gly	
Ile Val Phe Leu Leu		Pro Glu Thr Lys		Asn Leu Pro Leu Pro	
Thr Ile Lys Asp Val		Glu Asn Gln			

<210> 33

<211> 1775

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3474673CB1

<400> 33

```

atttcaggaa atgtgagggg gctctggggc ccttcctca ggcctgagg tcaccagca 60
gttctctct tctccctggc ctaggcctag caggtagggc cccgcacac atttgaggc 120
gggccagatg cccacagttc agagcctctt ttgtgccgg ggattggatc ccagggtgg 180
gtggggccag gctgtcccat tccccaacac tcctctctcc cggcgaaacc gggcaccagc 240
aggcgtttgc gagaggagat acgagctgga cgctggccc ttcctccca ccgggtccta 300
gtccaccgct ccgggcgcg gctccccgcc tctcccgcta tgtaccgacc gcgagcccg 360
gcggctcccg agggcagggt ccggggctgc gcggtgccca gcaccgtgct cctgtgctc 420
gcctacctgg cttacctggc gctgggcacc ggcgtgttct ggacgtgga gggccgcgc 480
gcgcaggact ccagcgcgag cttccagcgc gacaagtggg agctgttgca gaacttcacg 540
tgtctggacc gccggcgct ggactcgctg atccgggatg tcgtccaagc atacaaaaac 600
ggagccagcc tctcagcaa caccaccagc atggggcgct gggagctcgt gggctccttc 660
ttctttctg tgtccaccat caccaccatt ggctatggca acctgagccc caacacgatg 720
gctgcccgc tcttctgcat cttctttgcc cttgtggga tccactcaa cctcgtggtg 780
ctcaaccgac tggggcatct catgcagcag ggagtaaacc actgggccag caggctgggg 840
ggcacctggc aggatcctga caaggcgcg tggctggcgg gctctggcgc cctcctctcg 900
ggcctcctgc tcttctgct gctgccaccg ctgctcttct cccacatgga gggctggagc 960
tacacagagg gcttctactt cgccttcate accctcagca ccgtgggctt cggcgactac 1020
gtgattggaa tgaaccctc ccagaggtag ccactgtggt acaagaacat ggtgtccctg 1080
tggatcctct ttgggatggc atggctggcc ttgatcatca aactcatcct ctcccagctg 1140
gagacgccag ggagggtatg ttctgctgc caccacagct ctaaggaaga cttcaagtc 1200
caaagctgga gacagggacc tgaccgggag ccagagtccc actcccaca gcaaggatgc 1260
tatccagagg gacccatggg aatcatacag catctggaaac cttctgctca cgtgcaggc 1320
tgtggcaagg acagctagtt ataactccatt ctttggctgt cgtcctcggt agcaagacc 1380
ctgattttaa gctttgcaca tgtccacca aactaaagac tacattttcc atccacccta 1440
gaggtgggt gcagctatat gattaattct gccaatagg gtatacagag acatgtcctg 1500
ggtgacatgg gatgtgactt tcgggtgtcg gggcagcatg ccttctctcc ccacttcct 1560
actttagcgg gatgcaatgc cgcgatatg atggctggga gctctggcag ccatagcca 1620
ccatgaagta gcggcaatgt ttgagcggca caataagata ggaagagtct ggatctctga 1680
tgatcacaga gccatcctaa caaacggaat atcacccgac ctcttttatg tgagagagaa 1740
ataaacatct tatgtaaaat accaaaaaaa aaaaa 1775

```

<210> 34

<211> 1545

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4588877CB1

<400> 34

```

aatgagggcc ctgggggttg ggcccaggag tggggctgtg gtggtgagtg gacagggctg 60
ggctggaaat gtcccctgag tgccccctct cactcagge tatggcaacc tgagcccaa 120
cacgatggct gccgcctct tctgcatctt ctttgcctt gtggggatcc cactcaacct 180
cgtggtgctc aaccgactgg ggcatctcat gcagcaggga gtaaaccact gggccagcag 240
gctggggggc acctggcagg tgagggggct gctggacggg gtggggatgg gtcacttcta 300
gaatgagggg ctgtggtggg aattgggggt actaatgaca agaggtggga gcaagtgtta 360
ctggtgaggt tgtgttggga ttgggggtca ctgctcagaa tagggctcct agtgaaaaag 420
ggcattaatg gtggagatgg ggtgggactg ggcagacagg aaggacatga ggcacaggct 480
ccaggcaggg aactgggaga acacagacca ggtgaagagc ccccttctta ctggggacag 540
ctctggcctg cctccagctc cctcggctcc cagcctggg gtgaaggcct caggaggcct 600
ggggacaata ttgcacccac aggatcctga caaggcgcg tggctggcgg gctctggcgc 660
cctcctctcg ggctcctgc tcttctgct gctgccaccg ctgctcttct cccacatgga 720
gggctggagc tacacagagg gcttctactt cgccttcate accctcagca ccgtgggctt 780
cggcgactac gtgattggaa tgaaccctc ccagaggtag cactgtggt acaagaacat 840
ggtgtccctg ttggatcctt ttgggatggc atggctggcc ttgatcatca aactcatcct 900
ctcccagctg gagacgccag ggagggtatg ttctgctgc caccacagct ctaaggaaga 960
cttcaagtc caaagctgga gacagggacc tgaccgggag ccagagtccc actcccaca 1020

```

```

gcaaggatgc tatccagagg gacccatggg aatcatacag catctggaac cttctgctca 1080
cgctgcaggc tgtggcaagg acagctagtt atactccatt ctttggctgt cgtcctcggg 1140
agcaagaccc ctgatttttaa gctttgcaca tgtccaccca aactaaagac tacattttcc 1200
atccacccta gaggtctgggt gcagctatat gattaattct gcccaatagg gtatacagag 1260
acatgtcctg ggtgacatgg gatgtgactt tcgggtgtcg gggcagcatg cccttctccc 1320
ccacttcctt acttttagcgg gctgcaatgc cgccgatatg atggctggga gctctggcag 1380
ccatacggca ccatgaagta gcggcaatgt ttgagcggca caataagata ggaagagtct 1440
ggatctctga tgatcacaga gccatcctaa caaacggaat atcaccgcac ctcctttatg 1500
tgagagagaa ataaacatct tatgtaaaaa accaaaaaaa aaaaa 1545

```

<210> 35

<211> 1941

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472214CB1

<400> 35

```

atggcggaga aggcgctgga ggccgtgggc tgtggactag ggccgggggc tgtggccatg 60
gccgtgacgc tggaggacgg ggccggaaccc cctgtgctga ccacgcacct gaagaagggt 120
gagaaccaca tcaactgaagc ccagcgettc tcccacctac ccaagcgtct agccgtggac 180
atcgagttcg tggagctgtc ctattccgtg cgggaggggc cctgctggcg caaaaggggt 240
tataagaccc ttctcaagtg cctctcaggt aaattctgcc gccgggagct gattggcatc 300
atgggcccc caggggctgg caagtctaca ttcatgaaca tcttggcagg atacaggagg 360
tctggaatga aggggcagat cctggttaat ggaaggccac gggagctgag gaccttccgc 420
aagatgtcct gctacatcat gcaagatgac atgctgtctg cgcacctcac ggtgttggaa 480
gccatgatgg tgtctgctaa cctgaatctt actgagaatc ccgatgtgaa aaacgatctc 540
gtgacagaga tectgacggc actgggcctg atgtcgtgct cccacacgag gacagccctg 600
ctctctggcg ggcaaggaa gcgtctggcc atcgccctgg agctggtcaa caaccgcct 660
gtcatgttct ttgatgagcc caccagtggg ctggatagcg cctcttgttt ccaagtgggt 720
tccctcatga agtcctggc acaggggggc cgtaccatca tctgcacat ccaccagccc 780
agtccaagc tctttgagat gtttgacaag ctctacatcc tgagccaggg tcagtgcac 840
ttcaaaggcg tggtcaccaa cctgatcccc tatctaaagg gactcggctt gcattgcccc 900
acctaccaca acccgctga cctcgtcate gaggtggcct ctggcagta tggagacctg 960
aaccccatgt tgttcagggc tgtgcagaat gggctgtgcg ctatggctga gaagaagagc 1020
agccctgaga agaacgaggt ccctgcccc ctcacacagt tctgcacct cttcaagagg 1140
attgaaagcc acacctttgc caccagcacc ctgacccacc tacggttcac gtcccacgtg 1200
accttctgt ccactctcag ggacacgggt cctcctctac ctgcatattg gcgacgatgc cagcaaggtc 1260
gttatggcg tgcctatcgg cctcctctcc atgctgttcc tcatgttcgc cgccctcatg 1320
ttcaacaaca ccggtgcct ctttagagat gcggtcttca tgaggagagc cctcaactac 1380
ccaactgtgc tcaccgtccc ttacctggcc aagaccatgg ctgacgtgcc ctttcagggtg 1440
tggtacagcc tcaaagcgta ttagcttggt cagcattgtg cgggccagcc cgctgagacc 1500
gtgtgtccgg tggctctcgc agcctggcc ccttgggtgg ccacttttgt gggcccagtt 1620
agccgcttcc tgctctctc ttccaactcc ctacaggtgg ccacttttgt gcttcaagac catccccact 1680
gggtgctga tggagctcta tctctcctat gtcaggatag gctttgaggg tgtgatcctg 1740
accgcatcc ctgtcctctt gttctccggc ttctttgtca gcttcaagac catccccact 1800
tacctgcaat ggagctccta tctctcctat acatgtttag aggaacgctg cccgttccgg 1860
acgatctatg gcatggagcg aggagacctg gtggaggatg ccaagctcta catggacttc 1920
gagccacaga gcacccctcc agcgtggatg gtggaggatg cctaccttgt gctgcgttac 1941
ctggctcttg gcactctctt cctagccctg cggctgctgg
cgggtcaagt cagagagata g

```

<210> 36

<211> 4971

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473053CB1

<400> 36

```

caaagtagcg ggccgaggcc cgggggagcg gggccgcagc tgggggggcg ggagcccggt 60
gggagccgag ccgagcgcgc cccgccccag cccccgcat gggcagtagc gggccgcccg 120
ggcgggcgccc gaggcgtgag cgctgagggt ctcccatggg attgctggga tcttgcgtgg 180

```

tgagatggca	gtgtgtgcaa	aaaagcgccc	cccagaagaa	gaaaggaggg	cgcgggctaa	240
tgaccgagaa	tacaatgaga	aattccagta	tgcgagtaac	tgcatcaaga	cctccaagta	300
caatattctc	accttcctgc	ctgtcaacct	ctttgagcag	ttccaggaag	ttgccaacac	360
ttactttcctg	ttcctcctca	ttctgcagtt	gatccccag	atctcttccc	tgtcctgggt	420
caccaccatt	gtgccttttg	ttctttgtct	caccatcaca	gctgttaaag	atgccactga	480
tgactatttc	cgccacaaga	gcgataacca	ggtgaataac	cgccagtctc	aggtgctgat	540
caacggaatc	ctccagcagg	agcagtggat	gaatgtctgt	gttgggtgata	ttatcaagct	600
agaaaataac	cagtttgttg	cgggcgatct	cctcctcctt	tccagcagtg	agcccatgg	660
gctgtgttac	atagagacag	cagaacttga	tggcgagacc	aacatgaaag	tacgtcaggc	720
gattccagtc	acctcagaat	tgggagacat	cagtaagctt	gccaagtttg	acggtgaagt	780
gatctgtgaa	cctcccaaca	acaaactgga	caaattcagc	ggaaccctct	actggaagga	840
aaataagttc	cctctgagca	accagaacat	gctgtgcgg	ggctgtgtgc	tgcgaaacac	900
cgagtgggtgc	ttcgggctgg	tcatctttgc	aggtcccagc	actaagctga	tgcaaaacag	960
cggcagaaca	aagttcaaaa	gaacgagtat	cgatcgctta	atgaataccc	tggtgctctg	1020
gattttttgga	ttcctggttt	gcatgggggt	gatcctggcc	attggcaatg	ccatctggga	1080
gcacgaggtg	gggatgcgtt	tccaggtcta	cctgcctggg	gatgaggcag	tgagagtgac	1140
cttcttctct	ggcttctctc	ccttctgggt	ctacatcctc	atcctcaaca	cctgtgtgct	1200
catttctactc	tatgtcagtg	tggaggtcat	ccgtctgggc	cacagctact	tcatcaactg	1260
ggataagaag	atgttctgca	tgaagaagcg	gacgcctgca	gaagcccgc	ccaccacct	1320
aaacgaggag	ctggggccagg	tggagtacat	cctctccgac	aagacgggca	ccctcaccca	1380
gaacatcatg	gttttcaaca	agtgtcccat	caatggccac	agctatggtg	atgtgtttga	1440
cgctctggga	cacaagctg	aattgggaga	gaggcctgaa	cctgttgact	tctccttcaa	1500
tcctctgggt	gacaagaagt	tcttattttg	ggaccccagc	ctgctggagg	ctgtcaagat	1560
cggggacccc	cacacgcctg	agttcttccg	cctcctttcc	ctgtgtcata	ctgtcatgtc	1620
agaagaaaag	aacgaaggag	agctgtacta	caaagctcag	tccccagatg	agggggccct	1680
ggtcacgcga	gcccaggaact	ttgggtttgt	tttccgctct	cgcaccccca	aaacaatcac	1740
cgtccatgag	atgggacacag	ccatcaccta	ccagctgctg	gccatcctgg	acttcaacaa	1800
catccgcaag	cggatgtcgg	tcatagtgcg	gaatccagag	gggaagatcc	gactctactg	1860
caaagggggt	gacactatcc	tactggacag	actgcaccac	tccactcaag	agctgctcaa	1920
caccaccatg	gaccacetta	atgagtacgc	aggggaaggg	ctgaggaccc	tggtgctggc	1980
ctacaaggat	ctggatgaag	agtactatga	ggagtgggct	gagcgacgcc	tccaggccag	2040
cctggcccag	gacagccggg	aggacaggct	ggctagcatc	tatgaggagg	ttgagaacaa	2100
catgatgctg	ctgggtgcaa	cggccattga	ggacaaactt	cagcaagggg	ttccagagac	2160
cattgcccctc	ctgacactgg	ccaacatcaa	gatttgggtg	ctaaccggag	acaagcaaga	2220
catggctgtg	aacatcggct	attcctgcaa	gatgctgacg	gatgacatga	ctgaggtttt	2280
catagtcact	ggccatactg	tcttggaggt	cggggaggag	ctcaggaaag	cccgggagaa	2340
gatgatggac	tcatcccgct	ctgtaggcaa	cggcttcacc	tatcaggaca	agctttcttc	2400
ttccaagcta	acttctgtcc	tggaggccgt	tgtctggggg	tacgcccctg	tcataaatgg	2460
tcacagcctg	gcccacgcac	tggaggcaga	catggagctg	gagtttctgg	agacagcgtg	2520
tgcctgcaaa	gctgtcatct	gctgccgggt	gacccccttg	cagaaggcac	aggtgcttag	2580
actgtgtcaag	aagtacaaga	aggctgtgac	gcttgccatt	ggagacggag	ccaatgatgt	2640
cagcatgatc	aaaacggctc	acattggtgt	ggggatcagt	gggcaggaag	ggatccaggc	2700
tgtcttggcc	tccgattact	ccttctccca	gttcaagttc	ctgcagcgcc	tcctgtctgg	2760
gcattgggcgc	tggctctacc	tgcgaatgtg	caagtttctt	tgctatttct	tctacaaaaa	2820
ctttgtcttc	acttctgtcc	acttctgtgt	ttgtctcttc	tgtggcttct	cagcccagac	2880
cgtctatgac	cagtatttca	tcaccctgta	taacatcgctg	tacacctccc	tgccagtcct	2940
ggctatgggg	gtctttgatc	aggatgtccc	cgagcagcgg	agcatggagt	accctaagct	3000
gtatgagccg	ggccagctga	accttctctt	caacaagcgg	gagttcttca	tctgcatcgc	3060
ccagggcac	tacacctccg	tgtcatgttt	cttcatctcc	tatgggggtg	ttgtgtatgc	3120
caccgggat	gatggcactc	agctggctga	ctaccagtc	tttgagtgca	ctgtggccac	3180
atccttgggtc	attgtgggtta	gcgtgcagat	tgggctcgac	acaggctact	ggacggccat	3240
caaccacttc	ttcatctggg	gaagccttgc	tgtttacttt	gccatcctct	ttgccatgca	3300
cagcaatggg	ctcttcgaca	tgtttcccaa	ccagttccgg	tttgtgggga	atgccagaaa	3360
caccttggcc	cagcccacgg	tgtggctgac	cattgtgctc	accacagtgc	tctgcatcat	3420
gcccgtggtt	gccttccgat	tcctcaggct	caacctgaag	cggatctctc	ccgacacggg	3480
ccgctacaca	cagctcgtga	ggaagaagca	gaaggcccag	caccgctgca	tgcggcgggt	3540
tggecgact	ggctcccggc	gctccggcta	tgccttctcc	catcaggagg	gcttcgggga	3600
gctcatcatg	tctggcaaga	acatgcggct	gagctctctc	gcgctctcca	gcttcaccac	3660
ccgctccagc	tccagctgga	ttgagagcct	ggcgaggaag	aagagtga	gtgccagtag	3720
cccagtggtc	gggtccgaca	agcccccaa	ggcttgagg	cagaggtg	atgccctggt	3780
ccagtgaaca	gagcaccag	ggctggccag	tactgagggt	aacagcgtct	cggaaactgt	3840
ggtcctcatt	ccttgcctcc	cgccccccg	gtagactctg	tcctgctggt	cccaccacac	3900
atggctggga	catctgttcc	cagctgtagg	cccttccacc	agctggggag	ctagagggag	3960
caggcccaag	ggcagagcag	aggctgaggc	acggggagcc	agccccctc	ggggaccaga	4020
agtgggaacca	aaaacaagaa	aaaactgtga	gagatgtgtg	ctgcccctgc	cctgccctgg	4080
accacaggg	agactataat	ctccttattt	ttttactcct	actcccaga	ggggccctag	4140
tgcctctggt	cctgaattac	ataagaatgt	accatgccgg	gaagccagag	acctgcaggg	4200


```

gcctcggccc ctcacatcgt gtatgtctct ccttgatttg tgttgtgtcc agtttggttt 4260
tgtctttttt tatttgcaa gtggaggagg cttttatgtg acttttatgt tgtggttgg 4320
gtcttaactc tcctgggaaa aggaggctgg cacacactgg gatgccgcag cctggccggc 4380
tgtgggggtg tttgggagga tccatgtcgg ctctgcctgc agtgaccagt gctctgtggg 4440
gcagaggagc tgaccaggga gggagggtacc catgagcaga gggtagtggg agagtgtaaa 4500
ggagggtttg gtccctgtctg ctctctcacc ttgagagtaa agtgctgccc tctgccccca 4560
acacacacac atatcaattc ctggattcct tagtctgtct ggccctgggc tggagcctag 4620
gaaagtggcc cccaaatcct tagtgagcta aagctgggtc tgaaatttgg tcagtgggga 4680
ggggtagtgt tcttttcttt tttctttttt ttttttctt tttttttttg agatggagtc 4740
tcaactctgt cacctaggca agagtgtcaat ggcacaatct cagctcactg caacctccac 4800
ctcctggggt caagcgattc tcttgctctc ccggacccaa cacttggaact taatctcact 4860
ttcttaaatc tttctattct cagacacggg tctagtacca ttccttctct ttagccccag 4920
ggagcaaatt aaagaggtta cgagttaaaa tcttaaaaaa aaaaaaaaaa a 4971

```

<210> 37

<211> 1404

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473347CB1

<400> 37

```

atggtcctgg ctttccagtt agtctccttc acctacatct ggatcatatt gaaaccaaat 60
gtttgtgctg cttctaacat caagatgaca caccagcggg gctcctcttc aatgaaacaa 120
acctgcaaac aagaaactag aatgaagaaa gatgacagta ccaaagcgcg gcctcagaaa 180
tatgagcaac ttctccatat agaggacaac gatttctgcaa tgagacctgg atttggaggg 240
tctccagttg cagtaggtat agatgtccat gttgaaagca ttgacagcat ttcagagact 300
aacatggact ttacaatgac tttttatctc aggcattact ggaaagacga gaggtctctc 360
tttcttagca cagcaaacaa aagcatgaca tttgatcata gattgaccag aaagatctgg 420
gtgcctgata tcttttttgg ccactctaaa agatccttca tccatgatac aactatggag 480
aatatcatgc tgcgcgtaca ccctgatgga aacgtcctcc taagtctcag gataacgggt 540
tcggccatgt gctttatgga tttcagcagg tttcctcttg acactcaaaa ttgttctctt 600
gaactggaaa gctatgccta caatgaggat gacctaatgc tatactggaa acacggaaac 660
aagtccctaa atactgaaga acatatgtcc ctttctcagt tcttcattga agacttcagt 720
gcatctagtg gattagcttt ctatagcagc acaggctggg acaataggct tttcatcatc 780
tctgtgctaa ggaggcatgt tttcttcttt gtgtgcccac cctattacce agccatattg 840
atggtgatgc tttcatgggt ttcatttttg attgaccgaa gagctgttcc tgcaagaggt 900
tcctcgggaa tcaccacagt gctgaccatg tccacaatca tcaactgtgt gagegcctcc 960
atgccccagg tgtcctacct caaggctgtg tgtgtgtacc tggggtcag ctccctcttt 1020
gtgttctctg cagtcattga gtatgcagct gtgaactacc tcaccacagt ggaagagcgg 1080
aaacaattca agaagacagg aaagatttct aggatgtaca atattgatgc agttcaagct 1140
atggcctttg atggttggtt ccatgacagc gagattgaca tggaccagac ttcctctctt 1200
ctaaactcag aagacttcat gagaagaaaa ctgatatgca gcccagcac cgatttcact 1260
cggataaaga gaagaaaatc cctaggagga catgttggtg gaatcattct ggaaaacaac 1320
catgtcattg acacctattc taggatttta tccccattg tgtatatttt atttaatttg 1380
ttttactggg gtgtatatgt atga 1404

```

<210> 38

<211> 4048

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474240CB1

<400> 38

```

cttccatccc ccctcagcca ttccttactg ctctgggcaa ccgccagggt aagcccattt 60
gcactgggaa attggcgctg tttgggagaa gagaaacaga tcgattgccc ttgtgactcc 120
ccgccccctt cccatcccca cccccaccgc tctctccctc tttccctccc ccgccacctc 180
ccctcaccct gcctccttcc cgttccccac ccccaaacc cctcaccgcg ggcagtccgg 240
tgcgaggccc cctccggaag gtgaggggaa tggattggac tccgggtggg aaagcgggtg 300
tctagaagtg gtgctaattg gaagagaatt ctggtttcaa aagaggatgc tctgccaca 360
agagcggctc gcgcgctggc ctgggctcta gccgaggaga gatcccgga ggactccaga 420
gctccggggg agcgcctctc ggaagaccgg ggccaacatg cctgtgcgca gggggcatgt 480

```

```

ggcaccacaa aatacatttc tgggggacct cattcggaat tttgaagggc aaaataaaaa 540
atztatcatt gcaaattgcca gagtgcagaa ctgtgccatc atttattgca acgatgggtt 600
ctgtgagatg actggtttct ccaggccaga tgatcatgcaa aagccatgca cctgcgactt 660
tctccatgga cccgagacca agaggcatga tattgcccac attgcccagg cattgctggg 720
gtcagaagag aggaaagtgg aggtcaccta ctatcacaaa aatgggtcca cttttatttg 780
taacactcac ataattccag tgaaaaacca agagggcgctg gctatgatgt tcatcattaa 840
ttttgaatat gtgacggata atgaaaacgc tgccacccca gagagggtaa acccaatatt 900
accaatcaaa actgtaaacc ggaaatTTTT tgggttcaaa ttccttggtc tgagagtctt 960
catttacaga aagcagtcct taccacaaga agaccccgat gtggtggtca tgcattcatc 1020
taaacacagt gatgattcag tagccatgaa gcattttaag tctcctacaa aagaaagctg 1080
cagccctctt gaagcagatg acacaaaagg tttgatacag cccagcaaat gttctccctt 1140
ggtgaaatata tccggacctc ttgaccattc ctctcccaaa aggcaatggg accgactcta 1200
ccctgacatg ctgcagtcaa gttccagct gtccatttc agatcaaggg aaagcttatg 1260
tagtatacgg agagcatctt cgggccatga tatagaagga ttcggcgctc accccaagaa 1320
catattttaga gaccgacatg ccagcgaaga caatggtcgc aatgtcaaag ggccttttaa 1380
tcataacaag tcaagctacc tgggatccac atcagattca aacctcaaca aatacagcac 1440
cattaacaag attccacagc tcaacttgaa tttctcagag gtcaaaaactg agaaaaagaa 1500
ttcatcacct cttcttcag ataaaaacat tattgcacc aagggttaaag atcgaacaca 1560
caatgtgact gagaaagtga cccagggtct ctcttttaga gcagatgtcc tacctgaata 1620
caaactgcag acaccacgca tcaacaagtt tacgatattg cactacagcc ctttcaaggc 1680
agtcctgggac tggcttatcc tgctgttggt catatacact gctatattta ctctctactc 1740
tgcagccttc ctctcaatg acagagaaga acagaaaaga cgagaatgtg gctattcttg 1800
tagccctttg aatgtggtag acttgattgt ggatattatg tttatcatag atattttaat 1860
aaacttcaga acaacatatg taaatcagaa tgaagaagtg gtaagtgatc cgcctaaaat 1920
agcaatacac tacttcaaag gctggttctt gattgacatg gttgcagcaa ttccttttga 1980
cttgctgatt tttggtcag gttctgtaga gacaacaaca ttaattgggtc ttttgaagac 2040
tgcccgactc ctccgtcttg tgcgctggc caggaaactg gatcgatatt cagaatagg 2100
cgctgctgtt ctaatgctct taatgtgcat ctttgccctg attgtcact ggctggcttg 2160
catttggtat gcgattggga atgtagaaag gccttacctg actgacaaaa tccgatgggt 2220
ggattcctta ggacagcaaa ttgggaaacg ttacaatgac agtgactcaa gttctggacc 2280
atccattaaa gacaaatagc tcacagcact ttattttacc ttcagcagtt taaccagtgt 2340
aggattcggg aatgtgtctc ctaacacgaa ttcggagaaa atcttttcaa tttgtgtcat 2400
gttgattggc tcaataatgt atgcaagcat ttttggaat gtatctgcaa ttatccaaag 2460
actatactcg ggaactgcca ggtaccacat gcagatgctg cgagtaaaag agttcattcg 2520
ctttcaccaa atcccgaacc ctctgaggca acgtcttgaa gaatatttcc agcacgcatg 2580
gacttacacc aatggctatg acatgaacat ggtccataag ggtttcccag aatgcttaca 2640
agcagacatt tgtctacatc tcaaccagac attgtgcaa aactgcaaag cctttcgggg 2700
ggcaagtaaa ggttgccctt gagctttggc aatgaagttc aaaaccaccc atgcacctcc 2760
aggagacacc ctggttcact gtggggatgt cctcactgca ctttatttct tatccagagg 2820
ctccattaaa atctcctaaag atgacattgt ggtggtctatt ctgggaaaaa atgatattg 2880
tgagagaaatg gttcatcttt atgccaaaacc tggaaagtct aatgcagatg taagagcct 2940
cacatactgt gacttgcata agattcagcg agaagacttg ttagaggttt tggatatgta 3000
tccctgagttt tctgatcact ttctaacaaa cctagagttg actttcaacc taaggcatga 3060
gagcgcaaaag gctgatctcc tacgatcaca atccatgaat gattcagaag gagacaactg 3120
taaacataag agaaggaaat tgtcatttga aagtgaagga gagaaagaaa acagtacaaa 3180
tgatcctgaa gactctgcag ataccataag acattatcag agttccaaga gacactttga 3240
agagaaaaaaa agcagatcct catctttcat ctctccatt gatgatgaac aaaagccgct 3300
cttctcagga atagtagact cttctccagg aatagggaaa gcactctggg ccatcttgga 3360
agaaacagtg cccacctcag gaagaatgca catagataaa agaagtcact cttgcaaaga 3420
tatcactgac atgcgaagct gggaacgaga aaatgcacat cccagcctg aagactccag 3480
tccatctgca cttcagcgag ctgctgggg tatctctgaa accgaaagcg acctcaccta 3540
cggggaagtg gaacaaagat tagatctgct ccaggagcaa cttaacaggc ttgaatccca 3600
aatgaccact gacatccaga ccatcttaca gttgctgcag aaacaaacca ctgtggtccc 3660
cccagcctac agtatggtaa cagcaggatc agaatalcag agacccatca tccagctgat 3720
gagaaccagt caaccggaag catccatcaa aactgaccga agtttcagcc cttcctcaca 3780
atgtcctgaa tttctagacc ttgaaaaatc taaacttaaa tccaaagaat ccctttcaag 3840
tggggtgcat ctgaacacag cttcagaaga caacttgact tcacttttaa aacaagacag 3900
tgatctctct ttagagcttc acctgcgga tacgttcac caattaggca 3960
tcttcttttg ccagattcat ccctaagcac tgtaggaatc gtgggtcttc ataggcatgt 4020
ttctgatcct ggtcttccag ggaaataa

```

<210> 39
 <211> 1539
 <212> DNA
 <213> Homo sapiens
 <220>

<221> misc_feature
 <223> Incyte ID No: 7475338CB1

<400> 39
 atggagaaca aagaggcggg aacccctcca cccattccat ccagggaggg gcggctccag 60
 ccgacgctgt tgctggcgac actgagcgcg gcctttggct cagccttcca gtacggctac 120
 aacctctctg tggtaaacac gccgcacaag gtgttcaagt cattttacaa cgaaacctac 180
 tttgagcgac acgcaacatt catggacggg aagctcatgc tgcttctatg gtcttgacc 240
 gtctccatgt ttcctctggg cggcctgttg gggtcattgc tcgtgggcct gctggttgat 300
 agctgcggca ggaaggggac cctgctgatc aacaacatct ttgccatcat ccccgccatc 360
 ctgatgggag tcagcaaagt ggccaaggct tttgagctga tcgtcttttc ccgagtgggtg 420
 ctgggagctt gtgcaggat ctctacagc gcccttccca tgtacctggg agaactggcc 480
 cccaagaacc tgagaggcat ggtgggaaca atgaccgagg ttttcgtcat cgttgaggtc 540
 ttcttagcac agatcttcag cctccaggcc atcttgggca acccggcagg ttggccgggtg 600
 cttctggcgc tcacaggggt gcccgccctg ctgcagctgc tgacctgcc cttcttcccc 660
 gaaagcccc gctactccct gattcagaaa ggagatgaag ccacagcgcg acaagctctg 720
 aggaggctga gaggccacac ggacatggag gccgagctgg aggacatgcy tgcggaggcc 780
 cgggccgagc gcgccgaggg ccacctgtct gtgctgcacc tctgtgccct gcggctccctg 840
 cgctggcagc tctctccat catcgtgctc atggccggcc agcagctgtc gggcatcaat 900
 gcgatcaact actatgcgga caccatctac acatctgcgg gcgtggaggc cgctcactcc 960
 caatatgtaa cgggtggctc tggcgtgctc aacatagtga tgaccatcac ctcggtctg 1020
 cttgtggagc ggctgggacg gcggcacctc ctgctggccg gctacggcat ctgcggtct 1080
 gcctgcctgg tgctgacggg ggtgctccta ttccagaaca ggggtcccca gctgtcctac 1140
 ctcggcacatc tctgtgtctt tgccatcac gcgggacatt ccattggggc cagtcctgtc 1200
 cctcgggtgg tgaggaccga gatcttctct cagtcctccc ggccgggcagc tttcatgggt 1260
 gacggggcag tgcactggct caccactctc atcataggct tctgttccc atccatccag 1320
 gaggccatcg gtgcctacag tttcatcatc tttgccggaa tctgcctcct cactgcgatt 1380
 tacatctacg tggttattcc ggagaccaag ggcaaacat ttgtggagat aaaccgcatt 1440
 tttgccaaga gaaacagggt gaagcttcca gaggagaaag aagaaaccat tgatgctggg 1500
 cctcccacag cctctctctg caaggaaact tccttttag 1539

<210> 40
 <211> 3114
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7476747CB1

<400> 40
 ccaagcagtg cctcacttct gccttgtcta gctgtactct ggaaaattaa gaaatttatg 60
 agtgtagcac caagtatacc aatgggaagg atgggagtc gaagtcaagt gaactcagcc 120
 cgctctgtg tactttgcac ttttccattt cccttggtag caggcacttt catacttaat 180
 ccatagtgga tgagcaactc tgacaatgac agcttctacc ccagaggcga 240
 ccccaaacat ggagctaaag gctccagctg caggaggtct taatgctggc cctgtcccc 300
 cagctgccat gtccacgcag agacttcgga atgaagacta ccacgactac agctccacgg 360
 acgtgagccc tgaggagagc ccgtcggaag gcctcaacaa cctctcctcc ccgggctcct 420
 accagcgctt tggtaaaagc aatagcacia catggttcca gacctgatc cacctgttaa 480
 aaggcaacat tggcacagga ctcttgggac tccctctggc ggtgaaaaat gcaggcatcg 540
 tgatgggtcc catcagcctg ctgatcatag gcactgtggc cgtgcactgc atgggtatcc 600
 tgggtgaaatg tgctcaccac ttctgccgca ggctgaataa atcctttgtg gattatgggtg 660
 atactgtgat gtatggacta gaatccagcc cctgtcctg gctccggaac cacgcacact 720
 ggggaagacg tgttgtggac ttcttcttga ttgtcaccca gctgggatc tgcgtgtgtct 780
 attttgtgtt tctggctgac aactttaaac aggtgataga agcggccaat gggaccacca 840
 ataactgcc aacaatgag acggtgattc tgacgcctac catggactcg cgactctaca 900
 tgctctcctt cctgcccttc ctgggtgctg ttggttttcat caggaacctc cgagccctgt 960
 ccattcttct cctgttggcc aacatcacca tgctggctag cttgggtcat atctaccagt 1020
 tcattgttca gaggatccca gacccagcc acctcccctt ggtggcccc ttggaagacct 1080
 accctctctt ctttggcaca gcgatttttt catttgaagg catttgcccc gttctgcccc 1140
 tggaaaacaa aatgaaggat cctcggaagt tccactcat cctgtacctg ggcattggta 1200
 tcgtcaccat cctctacatc agcctggggg gtctggggta cctgcaattt ggagctaata 1260
 tccaaggcag cataaccttc aacctgccca actgctgggt gtaccagtca gtttaagctgc 1320
 tgtactccat cgggatcttt ttacctaag cactccagtt ctacgtcccc gctgagatca 1380
 tcatccctct ctttgtgtcc cgagcgcccg agcctgtgta gttagtgggtg gacctgtttg 1440
 tgcgcccagt gctgggtctg ctgacatcac tgtctggcag tgttgacaat ggctgggtatg 1500
 gcacggaagc cgatggcacc tcctgcggca gtgcaccatt ggtcttctgc agttcctcct 1560

tccctggctca	cccgtggctg	agtttcagat	gtgagagcca	gtgggtgtcc	tgacacagag	1620
atacgggtcgt	cgtgtggggc	ttcgccagg	gcaccttggc	catcctcatc	ccccgcctgg	1680
acctgggtcat	ctccctgggtg	ggctccgtga	gcagcagcgc	cctggccctc	atcatcccac	1740
cgctcctgga	ggtcaccacc	ttctactcag	agggcatgag	ccccctcacc	atctttaagg	1800
acgccctgat	cagcatcctg	ggcttcgtgg	gctttgtggg	ggggacctat	gaggctctct	1860
atgagctgat	ccagccaagc	aatgctccca	tcttcatcaa	ttccacctgt	gccttcata	1920
agggatctgg	gttcgtctct	gcagctgcct	acccctgccc	catgtgtccc	ccgttacctg	1980
tcctcagagc	ctcaggtatg	gtccaggctc	tgaggaaagt	cagggttgct	gtgtgggaac	2040
ccctctgcct	ggcacctgga	tacctggggc	caggtaacct	gagggcagtg	gagaggtggg	2100
gtggcagaca	cgcagaagtg	ctactagtga	cagggtgcgc	atcgctcacc	tgtacctatt	2160
tacaccaga	actttccagc	tccccctcat	catgctcctc	ccttcttacc	tgcctcccc	2220
ctgctgggtg	acctgcacca	actcattctt	actgcacagt	tcactttatt	taacaatttt	2280
catgtccccc	acctcatgtt	ttcacctttt	ctggggccagg	catagattaa	gtaactggga	2340
acgccccctc	tttataaagc	tgggcttctt	tctcatctct	ctcccaaagt	ttgtatctca	2400
gtattcttcc	tattcgagtc	tccagggggg	ggctggacct	acctgggtcat	ttgaaacagg	2460
cccccaagct	ggagttttta	atctggactc	tctggcttgc	tgtgaccttc	aaggcaatgc	2520
ttctcttccc	tggattcctt	agtgtgggtc	acagtactgt	gttcttagtt	gctttagctc	2580
ttaaaacata	cgaagtgttg	cctaaactga	aaatatattat	cttttattta	aaatcagatt	2640
tttgttttta	gactgtctta	gatctggggc	tattacgaat	cacttcttct	tcagtaaaact	2700
ttgactcaac	ttctcctgct	gaaaagaagc	tcgctccaga	tgtctgcatg	ggtcctcggc	2760
actcttggct	gaggattcaa	aggttttaat	caggatctgc	taaaaatgta	cctcggtagg	2820
gaggcacaga	ttttgcctcc	tgttgaccag	cctggtttca	taccgaaaag	acattgaagg	2880
actgcagaaa	tgtatgggtg	caccggggcg	aggggaagggt	ggctgagtga	gaggcgata	2940
aaatggggct	gtgtgcatgc	aggcccatgt	ttcagcctca	gcccacgcca	ggtgaaagga	3000
tcagcaatgc	tctgttgcca	tcgtgctggg	acgacaccag	ctctattgcc	accgatgagt	3060
agctgagggtc	agtgtgcaca	gagtttgaaa	ttaagttaat	agactttaca	gcag	3114

<210> 41

<211> 2877

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477898CB1

<400> 41

atgccgggtcc	gcagggggcca	cgctcgctccc	caaaacactt	acctggacac	catcatccgc	60
aagttcgagg	gccaaagtgc	gaagttcctg	attgccaatg	ctcagatgga	gaactgcgcc	120
atcatttact	gcaacgacgg	cttctgcgaa	ctcttcggct	actcccgagt	ggaggtgatg	180
cagcaaccct	gcacctgcga	cttcttcaca	ggccccaca	caccaagcag	cgccgtgtcc	240
cgcctagcgc	aggccctgct	gggggctgag	gagtgcgaagg	tggacatcct	ctactaccgc	300
aaggatgcct	ccagcttccg	ctgcctggta	gatgtgggtg	ccgtgaagaa	cgaggacggg	360
gctgtcatca	tgttcattct	caacttcgag	gacctggccc	agctcctggc	caagtgcagc	420
agccgcagct	tgtcccagcg	cctgttgctc	cagagcttcc	tgggctccga	gggctctcat	480
ggcaggccag	gcggaccagg	gccaggcaca	ggcaggggca	agtacaggac	catcagccag	540
atcccacagt	tcacgctcaa	cttcgtggag	ttcaacttgg	agaagcaccg	ctccagctcc	600
accacggaga	ttgagatcat	cgcgccccat	aagggtggtg	agcggacaca	gaacgtcact	660
gagaagggtca	cccaggctct	gtccctgggc	gcgatgtgct	tgccggagta	caagctgcag	720
gcgcgcgcga	tccaccgctg	gaccatcctg	cactacagcc	ccttcaaggc	cgtgtgggac	780
tggctcatcc	tgtctgtggg	catctacacg	gctgtcttca	cgccttactc	agccgccttc	840
ctgctcagcg	atcaggacga	atcagggcgt	ggggcctgca	gctatacctg	cagtccccctc	900
actgtgggtg	atctcatcgt	ggacatcatg	ttcgctgctg	acatcgctcat	caacttccgc	960
accacctatg	tcaacaccaa	tgatgagggtg	gtcagccacc	ccgcgcgcct	cgccgtccac	1020
tacttcaagg	gctggttcct	cattgacatg	gtggccgcga	tccctttcga	cctcctgac	1080
ttccgcactg	gctccgatga	gaccacaacc	ctgattgggg	tattgaagac	agcgcggctg	1140
ctgcccgtgg	tgcgcgtagc	acggaagctg	gaccgctact	ctgagtatgg	ggcggctgtg	1200
ctcttcttgc	tcatgtgcac	ctttccgctc	atagcgcact	ggctggcctg	catctgggtac	1260
gccatcgggc	atgtggagcg	gccctaccta	gaacacaaga	tccgctggct	ggacagcctg	1320
gggtgtgcag	ttgggacagc	ctacaacggc	agcgacccag	cctcggggccc	ctcgggtgac	1380
gacaagtatg	tcacagccct	ctacttcacc	ttcagcagcc	tcaccagcgt	gggcttcggc	1440
aatgtctcgc	ccaacaccaa	ctccgagaag	gtcttctcca	tctgcgtcat	gctcatcggc	1500
tccctgatgt	acgccagcat	cttcgggaac	gtgtccgcga	tcattccagcg	cctgtactcg	1560
ggcacccgcg	gtaccacac	gcagatgtcg	cgtgtcaagg	agttcatccg	cttccaccag	1620
atccccaacg	cactgcgcca	gcgcctggag	gagtatcttc	agcacgcctg	gtccctacac	1680
aatggcattg	acatgaacgc	gggtgctgaag	ggcttccccg	agtgccctgca	ggctgacatc	1740
tgcctgcacc	tgcaccgcgc	actgctgcag	cactgcccag	ctttcagcgg	cgccggcaag	1800

ggctgcctgc	gcgcgctagc	cgtcaagttc	aagaccaccc	acgcgcgcgc	tggggacacg	1860
ctgggtgcacc	tcggcgacgt	gctctccacc	ctctacttca	tctcccagag	ctccatcgag	1920
atcctgcgcg	acgacgtggt	cgtggccatc	ctaggaaaga	atgacatctt	tggggaaacc	1980
gtcagcctcc	atgccagacc	aggcaagtcc	agtgcagacg	tgcgggctct	gacctactgc	2040
gacctgcaca	agatccagcg	ggcagatctg	ctggagggtg	tggacatgta	cccggccttt	2100
gcggagagct	tctggagtaa	gctggagggtc	accttcaacc	tgcgggacgt	aaccgggggt	2160
ctccactcat	ccccccgaca	ggctcctggc	agccaagacc	accaaggttt	ctttctcagt	2220
gacaaccagt	cagatgcagc	ccctcccctg	agcatctcag	atgcattctg	gctctggcct	2280
gagctactgc	aggaaatgcc	cccaaagcac	agcccccaaa	gccctcagga	agaccagat	2340
tgctggcctc	tgaagctggg	ctccaggcta	gagcagctcc	aggcccagat	gaacaggctg	2400
gagtcccgcg	tgctctcaga	cctcagccgc	atcttgcagc	tcctccagaa	gccccatgcc	2460
cagggccacg	ccagctacat	tctggaagcc	cctgcctcca	atgacctggc	cttgggtcct	2520
atagcctcgg	agacgcagag	tccaggggccc	agggtgcccc	agggttttct	gcctcctgca	2580
cagaccccaa	gctatggaga	cttggatgac	tgtagtccaa	agcacaggaa	ctcctcccc	2640
aggatgcctc	acctggctgt	ggcaatggac	aaaactctgg	caccatcctc	agaacaggaa	2700
cagcctgagg	ggctctggcc	accctatgcc	tcacctctac	atccctgga	agtacaagga	2760
ctcatctgtg	gtccctgctt	ctcctccctc	cctgaacacc	ttggctctgt	tcccaagcag	2820
ctggacttcc	agagacatgg	ctcagatcct	ggatttgcag	ggagttgggg	ccactga	2877

<210> 42

<211> 2820

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7472728CB1

<400> 42

atggggcatc	aaggggcatt	tgaagaagga	aatggtggac	tgagagtgat	agcgacctgg	60
aggaggaagg	aggcttgag	aagggaactgt	cttttaggag	ccctgcccag	tgtttctgt	120
ggagggtggg	gccatcgtgg	aagacagacc	tatggtaggg	cttgtggggg	gaaagaaaag	180
cccttttagtc	ttttgggtcc	tcaaatacaca	gtttatgcag	tttggcccca	gtcagaggga	240
ccccaggaag	gcagactcag	ggtaaattct	gcctgtcttc	caccagagag	gggactcacc	300
aacgcttgta	caaaccatga	agaactctct	ttggactggt	tgctttttga	gaatgttaac	360
accttgactc	tggatttctg	cctatgggaa	aaaaccacaa	tagtgccagg	ggtgcttcca	420
tatgcaggat	taactctgca	gtcaaagttt	ctggtgggca	gagcattggt	agcaggggtc	480
catgtgatca	cactgacacc	tgagagagtg	acacaccatg	tacatggctg	gtatatggag	540
gatggattta	aggggggacag	gactgaaggc	tgctgcagtg	attcagtggc	cgttcccgcg	600
gcagcacccg	tgtgccagcc	caagagcgcc	actaacgggc	aacccccggc	tcgggtcccg	660
actccaactc	cgcgcctgtc	catttccctc	cgagccacag	tggtagccag	gatggaaagg	720
acctcccaag	ggggccttgca	gaccgtcatg	aagtgggaaga	cgggtggttg	catctttgtg	780
gttgtggtgg	tctaccttgt	cactggcggt	cttgtcttcc	gggcatggga	gcagcccttt	840
gagagcagcc	agaagaatac	catcgccctg	gagaaggcgg	aattcctgcg	ggatcatgtc	900
tgtgtgagcc	cccaggagct	ggagacgttg	atccagcatg	ctcttgatgc	tgacaatgcg	960
ggagtcagtc	caataggaaa	ctcttccacc	aacagcagcc	actgggacct	cggcagtgcc	1020
tttttctttg	ctggaactgt	cattacgacc	atgtatggga	atattgctcc	gagcactgaa	1080
ggaggcaaaa	tcttttgtat	tttatatgcc	atctttggaa	ttccactctt	tggtttctta	1140
ttggctggaa	ttggagacca	acttggaaac	atctttggga	aaagcattgc	aagagtggag	1200
aaggctcttt	gaaaaaagca	agtgaagtcag	accaagatcc	gggtcatctc	aaccatcctg	1260
ttcatcttgg	cgggctgcat	tgtgtttgtg	acgatccctg	ctgtcatctt	taagtacatc	1320
gagggttgga	cggccttgga	gtccattttac	tttgtgtgtg	tcactctgac	cacggtgggc	1380
tttggtgatt	ttgtggcagt	ggttgttttc	aggggaaacg	ctggcatcaa	ttatcgggag	1440
tggtataagc	ccctagtgtg	gttttggate	cttgtttggc	ttgcctactt	tcgactgtc	1500
ctcagtatga	tcggagattg	gctacgggtt	ctgtccaaaa	agacaaaaga	agaggtgggt	1560
gaaatcaagg	cccatgcggc	agagtgggaag	gccaatgtca	cggctgagtt	ccgggagaca	1620
cggcgaaggc	tcagcgtgga	gatccacgat	aagctgcagc	gggcagccac	catccgcagc	1680
atggagcgcc	ggcggctggg	cctggaccag	cgggcccact	cactggacat	gctgtcccc	1740
gagaagcgct	ctgtctttgc	tgccctggag	accggccgct	tcaaggcctc	atcccaggag	1800
agcatcaaca	acggcccca	caacctgcgc	ctgaaggggc	cggagcagct	gaaccaagcat	1860
gggcagggtg	cgtccgagga	caacatcatc	aacaagtctg	ggteccacct	cagactcacc	1920
aagaggaaaa	acaaggacct	caaaaagacc	ttgcccagag	acgttcagaa	aatctacaag	1980
accttccgga	attactccct	ggacgaggag	aagaaagagg	aggagacgga	aaagatgtgt	2040
aactcagaca	actccagcac	agccatgctg	acggactgta	tccagcagca	cgctgagttg	2100
gagaacggaa	tgatacccac	ggacaccaaa	gaccgggagc	cggagaacaa	ctcattactt	2160
gaagacagaa	actaaatgtg	aaggacattg	gtcttggact	gagcgttgtg	tgtgtgtgtg	2220
tgtgtgtgtt	tttaatatcc	acactgagac	atgtgcctta	aacagacttt	ttagtccaaa	2280

```

attacatagc attgaagaat atattttcact gtgccataaa caactgaaag cttgctctgc 2340
caaaaggaat cagagaacaa gaacttcatt tcagatagca aacgcaggac acaccaagag 2400
tgtccgtgca cgtagccggt tctggccgta catgttaagg gcatttcagt ggcagtgcgtg 2460
tacccttggg cagtgcctacc tgggcacaca cgtagacaag ggcagctatt ccttagacca 2520
gcctcctgaa agaaacaggt gtgtcttttt agtggagtcg tagtaatatg tgcacacaca 2580
gaaggggacc tgattgggtg ggagctgggt atgtgtaact agcgttggag ttgacatttt 2640
ggcatgtgct ctgagcttga attttgatac caaccattca gtgcatcata cctagtcttt 2700
ctatgctcca aatgaatgtc tgtggggacc tgagagcacc tgggaatttg tgggaagcaga 2760
tcagagcaca cgtacgaaaa ggtgcaattc cttttctcat gacaaaaggg aaaaaataa 2820

```

```

<210> 43
<211> 1440
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 7474322CB1

```

```

<220>
<221> unsure
<222> 757
<223> a, t, c, g, or other

```

```

<400> 43
atgtacaatg agattctgat gctggggggcc aaactgcacc cgacgctgaa gctggaggag 60
ctcaccaaca agaaggggaat gacgccgctg gctctggcag ctgggaccgg gaagatcggg 120
aatcgccacg acatgctctt ggtggagccg ctgaaccgac tcctgcagga caagtgggac 180
agattcgctc agcgcctctt ctacttcaac ttcttggtct actgcctgta catgatcatc 240
ttcaccatgg ctgcctacta caggcccgtg gatggcttgc ctccctttaa gatggaaaaa 300
actggagact atttccgagt tactggagag atcctgtctg tgttaggagg agtctacttc 360
tttttccgag ggattcagta tttcctgcag aggcggccgt cgatgaagac cctgtttgtg 420
gacagctaca gtgagatgct tttgtttctg cagtcactgt tcatgctggc caccgtgggtg 480
ctgtacttca gccacctcaa ggagtatgtg gcttccatgg tattctcctt ggccttgggc 540
tggaccaaca tgctctacta caccgcggtt ttccagcaga tgggcatcta tgccgtcatg 600
atagagaaga tgatcctgag agacctgtgc cgtttcatgt ttgtctacat cgtcttcttg 660
ttcgggtttt ccacagcggg ggtgacgctg attgaagacg ggaagaatga ctccctgcctg 720
tctgagtgca cgtcgcacag gtggcggggg cctgctngca ggcccaatag ctccctacaac 780
agcctgtact ccacctgcct ggagctgttc aagttcacca tcggcatggg cgacctggag 840
ttcactgaga actatgactt caaggctgtc ttcatcatcc tgctgctggc ctatgtaatt 900
ctcacctaca tagttctcct cctcaacatg ctcatgtctc tgatgggcga gactgtggag 960
aacgtctcca aggagagcga acgcatctgg cgctgcaga gagccatcac catcctggac 1020
acggagaaga gcttccctaa gtgcatgagg aaggccttcc gctcaggcaa gctgctgcag 1080
gtgggggtaca cactgatgg caaggacgac taccggtggt gcttcgtgga cgaggtgaac 1140
tggaccacct ggaacaccaa cgtgggcac atcaacgaag acccgggcaa ctgtgagggc 1200
gtcaagcgca cctgagctt ctccctgcgg tcaagcagag tttcaggcag aacttggaa 1260
aactttgccc tgggtcccc ttttaagagag gcaagtgtct gagataggca gtctgtcag 1320
cccgagggaag tttatctgct acagttttca gggctctctg agccagagga cgctgaggtt 1380
ttcaagagtc ctgccgcttc cggggagaag tgaggacgtc acgcagacag cactgtcaac 1440

```

```

<210> 44
<211> 2394
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 5455621CB1

```

```

<400> 44
atttcagaac acatctgaat tccttctctg tggcatatgc tttaggagag gagcagacag 60
ctcttagcta gggtcagatt tcaaattctc atctcttggg gccaatacca ccaccagatt 120
cttctttgaa gtcaactttt gagatcttca ctaagtacac gttggtgtct gaagattcac 180
acgagtgcct ctggtaatca ttttcttcag ggaatcacag tctctcctct cagcaaagca 240
tccactgtac tgaactttgc ttttggaaac atcttcttcc tgagacctcg ttgaaagaaa 300
ctctctgggt tcatactttc caatatggag gtgaagaact ttgcagtttg ggattatgtt 360
gtatttgcag cctctttttt catttctctt ggaattgggg tgttctttgc cattaaggag 420

```

```

agaaaaaagg caacttcccg agagtctctg gttgggggaa ggcaaatgag ctttggccct 480
gtcggcttgt ctctgacagc cagcttcatg tcagctgtca cgggtcctggg gacctcttct 540
gaagtctacc gctttggggc atccttccca gtcttcttca ttgcttacct atttgtcatc 600
ctcttaacat cagagctctt tctccctgtg ttctacagat ctggtatcac cagcacttat 660
gagtacttac aactacgatt caacaaacca gttcgctatg ctgccacagt catctacatt 720
gtacagacga ttctctacac aggagtgggtg gtgtatgctc ctgccctggc actcaatcaa 780
gtgactgggt ttgatctctg gggctctgtg ttgcaacag gaattgtttg cacattctac 840
tgtaccctgg gaggattaaa agcagtgggtg tggacagatg catttcagat ggttgtcatg 900
atttgtgggt tcttaacggg tctcattcaa ggatcaactc atgctggggg attccacaat 960
gtattagagc aatcaacaaa tggatctcga ctacatatat ttgactttga tgtagatcct 1020
ctcagggcag acactttttg gactatcaca gtgggaggaa cttttacttg gctcggaaac 1080
tatgggggtc atcaatcaac tattcagcga tgcactctct gcaaaaacaga aaagcatgct 1140
aagcttgcc tgtatttttaa cttgctgggt ctctggatca ttctgggtgtg tgcgtctctc 1200
tctggcttaa tcatgtactc tcaactttaaa gactgtgacc cttggacttc tggcatcatc 1260
tcagcaccag accagctgat gccgtacttt gtcatggaga tatttgccac aatgccagga 1320
ctgccaggag tttttgtggc ttgtgccttc agtggaaact tgagcaccgt ggcttcagc 1380
atcaatgcct tggcaacagt gacctttgag gattttgtca agagctgttt tcctcatctc 1440
tccgacaagc tgagcacctg gatcagtaaa ggcttatgtc tcttatttgg cgtgatgtgt 1500
acctctatgg ctgtggctgc atctgtcatg ggaggtgttg tgcaggcttc cctcagcatt 1560
cacggcatgt gtggaggacc aatgctgggc ttattctccc tgggaatcgt gttccctttt 1620
gtgaactgga aggggtgcact aggggtctct cttactggaa tcacctgtgc attttgggtg 1680
gccattgggg ccttcattta ccttcacca gcctctaaga catggccttt gcctctatca 1740
acagaccaat gtatcaaata aaatgtgaca gcaacagggc ctccagtact atccagcaga 1800
cctggaatag ctgatacctg gtactcgatc tctaccttt actacagtgc actgggctgc 1860
ttaggatgca ttgttgctgg agtaatcatc agcctcataa caggtcgcca aagaggtgag 1920
gatattcaac cactgttaat tagaccagtt tgaattttat ttgtcttttg gtctaagaag 1980
tacaaaacac tatgctgggtg tggagttcag catgacagtg ggacagagca ggaaaacctt 2040
gagaatggca gtgcccgaa acaggggggt gaatctgtct tacagaacgg actcagaaga 2100
gaaagcctgg tacatgttcc aggctatgat cctaaggaca aaagctacaa caatatggca 2160
tttgagacta cccatttcta aggcaatacc tgtatgaatg cacacacaca cgtgaagaag 2220
acacacacac cacaacttc cactactctt ttgcctactt gttagtagat atgtatagtt 2280
gccattgcta gaagacaggg atgtctgggt cctattttcta cttattttata actacatgca 2340
aatgactgt ctctcgggat attctttgaa agactccaac tttcacagag aaaa 2394

```

<210> 45

<211> 2890

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477248CB1

<400> 45

```

gaataactaag ccaggggcaga atgcttgtga agtagcaact aaagtggcag tgtttcttct 60
gaaattctca ggcagtcaga ctgtcttagg caaatcttga taaaatagcc cttatccagg 120
tttttatcta aggaatccca agaagactgg ggaatggaga gacagtcaag gggtatgtca 180
gaaaaggatg agtatcagtt tcaacatcag ggagcgggtg agctgcttgt cttcaatttt 240
ttgtcatcct ttaccatttt gacaatctgg ttatttaaaa atcatcgatt ccgcttcttg 300
catgaaactg gaggagcaat ggtgtatggc gactaatttt gactaatttt acgatatgct 360
acagcaccaa ctgatattga aagtggact gtctatgact gtgtaaaact aactttcagt 420
ccatcaactc tgctgggttaa tatcactgac caagtttatg aatataaata caaaagagaa 480
ataagtacgc acaacatcaa tctcatcaa ggaaatgcta tacttgaaaa gatgacattt 540
gatccagaaa tcttcttcaa tgttttactg ccaccaatta tatttcatgc aggatatagt 600
ctaaagaaga gacacttttt tcaaaactta ggatctatct taacgtatgc cttcttggga 660
actgccatct cctgcacgtg cataggggta attatgtatg gttttgtgaa ggctatgata 720
catgctggcc agctgaaaaa tggagacttt catttcaact actgtttatt ttttggttca 780
ctgatgtctg ctacagatcc agtgacagtg ctggccattt tccatgaact gcacgtcgac 840
cctgacctgt acacactctt gtttggagag agtgtgttga atgatgcagt ggccatagtc 900
cttacatatt ctatagtcct ttacagtcct aaggagaatc caaatgcatt tggatgccga 960
gcattcttcc agtctgtggg gaatttctct ggaatcttct ctggctcatt tgcaatgggg 1020
tctgcgtatg ccatcatcac agcactgttg accaaaatta ccaagctgtg tgagttccc 1080
atgctggaaa cgggectgtt tttcctgctt tcttggagtg ccttctgtgc tgccgaggct 1140
gccggcctaa cagggatagt tgcgtgtctc ttctgtggag tcacacaagc acattatacc 1200
tacaacaatc tgtcttcgga ttccaaaata agaactaaac agttgtttga atttatgaac 1260
tttttggcgg agaacgtcat cttctgttac atgggcctgg cactgttcac gttccagaat 1320
catatcttta atgctctttt tatacttggga gcctttctag caatttttgt tgccagagcc 1380

```


tgcaacatat	atccccctctc	cttctctctg	aatctaggcc	gaaaacagaa	gatccccctgg	1440
aacttttcagc	acatgatgat	gttttcagggt	ttgcgaggag	cgatcgcat	tgcccttagct	1500
attcggaaca	cagaatctca	gccccaaaca	atgatgttta	ccactacgct	gctcctcgtg	1560
ttctttcactg	tctgggtatt	tggaggaggga	acaaccccc	tgttgacttg	gcttcagatc	1620
agagttggcg	tggacctgga	tgaaaatctg	aaggaggacc	cctcctcaca	acaccaggaa	1680
gcaaataact	tgataaaaa	catgacgaaa	gcagagagtg	ctcggctctt	cagaatgtgg	1740
tatagctttg	accacaagta	tctgaaacca	attttaaccc	actctgggtcc	tccgctgact	1800
acaacattac	ctgaatggtg	tggtecgatt	tccaggctgc	ttaccagtcc	tcaagcctat	1860
ggggaacagc	taaaagagga	tgatgtggaa	tgcattgtaa	accaggatga	actagccata	1920
aattaccagg	agcaagcctc	ctcaccctgc	agtcctcctg	caaggctagg	tctggaccag	1980
aaagcttcac	ccagacgccc	aggcaaggaa	aacatttatg	agggagacct	cggccctggg	2040
aggctatgaa	ctcaagcttg	agcaaacctt	gggtcaatcc	cagttgaatt	aattggcatg	2100
aagagtacag	atgtaatcac	aagtaatgca	agactcactg	aggaatacaa	gccaaagctga	2160
tgaggcagta	caggggagag	gctggaaaac	atattaagag	cataaattgg	agagaatcaa	2220
agccttgctca	catggatcct	ctgggtgcctg	aagaaatgag	atttttattat	ccctctctat	2280
tatgcaaatg	aatttagttt	tttgacagca	gccattctga	ttactggatt	ggctgggggtg	2340
gggatggagg	tatcaggagt	ctagctgctg	gaggatggga	cagctgtgct	gggtcttcag	2400
ggcattttctg	ctgcgaatgc	ggctctccag	gcccttcact	tctattctgg	atttttattcc	2460
ctccattaag	gagagtttta	aaataaaaaga	aagcttctga	gagtaaacad	tttgctccta	2520
agctgaaggg	aatgcccgag	tatttagtaa	gtgataagtt	tcttattttg	aggacttgac	2580
tcccatttgc	tctcagtgac	cccagggcag	agccagaga	agtgttccgt	accactgct	2640
gatggtttcc	cagagcccac	actgagttga	agaacctatt	gttcttcttg	gcacccctct	2700
tatgctactt	ctcccatcgc	tcaaaggggt	tgcctatggc	tgggtgtgcc	ctgccctaaa	2760
tgcagcacca	ctttcaagca	gcttctagct	atagctttcc	accagggtatt	tttaattcca	2820
ttcacctcc	tccccagca	attcaccagt	caggagtgat	ttttactgta	aagatgggtg	2880
cttagtaaaa						2890

<210> 46

<211> 3926

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2944004CB1

<400> 46

ctggaccttt	aatccactgt	aggatatggac	aggggaagaaa	ggaagaccat	caatcagggt	60
caagaagatg	aaatggagat	ttatgggttac	aatttgagtc	gctggaagct	tgccatagtt	120
tctttaggag	tgatttgctc	tgggtgggggt	tctcctcctc	ctctctattg	gatgcctgag	180
tggcgggtga	aagcgacctg	tgctcagagct	gcaattaaag	actgtgaagt	agtgtgctg	240
aggactactg	atgaattcaa	aatgtgggtt	tgtgcaaaaa	ttcgcgttct	ttctttggaa	300
acttaccag	tttcaagctc	aaaatctatg	tctaataagc	tttcaaatgg	ccatgcagtt	360
tgtttaattg	agaatcccac	tgaagaaaat	aggcacagga	tcagtaaata	ttcacagact	420
gaatcacaa	agatttcgtta	tttcacccac	catagtgtaa	aatatttctg	gaatgatacc	480
attcacaatt	ttgatttctt	aaagggactg	gatgaagggtg	tttcttgtac	gtcaatttat	540
gaaaagcata	gtgcaggact	gacaaaagggg	atgcatgcct	acagaaaact	gctttatgga	600
gtaaatgaaa	ttgctgtaaa	agtgccttct	gtttttaagc	ttctaattaa	agaggttctc	660
aaccattttt	acattttcca	gctgttcagt	gttatactgt	ggagcactga	tgaatactat	720
tactatgctc	tagctattgt	ggttatgtcc	atagtatcaa	tcgtaagctc	actatatctc	780
attagaaagc	aatatgttat	gttgcatgac	atgggtggcaa	ctcatagtac	cgtaagagtt	840
tcagtttgta	gagtaaatga	agaaatagaa	gaaatctttt	ctaccgacct	tgtgccagga	900
gatgtcatgg	tcatccatt	aaatgggaca	ataatgcctt	gtgatgctgt	gcttattaat	960
ggtacctgca	ttgtaaacga	aagcatgtta	acaggagaaa	gtgttccagt	gacaaagact	1020
aatgtgcaa	atccttcagt	ggatgtgaaa	ggaataggag	atgaattata	taatccagaa	1080
acacataaac	gacatacttt	gttttgtggg	acaactgtta	ttcagactcg	tttctacact	1140
ggagaactcg	tcaaagccat	agttgttaga	acaggattta	gtacttccaa	aggacagctt	1200
gttcgttcca	tattgtatcc	caaaccaact	gattttaaac	tctacagaga	tgccacttg	1260
tttctactat	gtcttgtggc	agttgtggc	attgggttta	tctacactat	tattaaatgc	1320
attttaaatg	aggtaacaag	tggggtcata	attatcgagt	ctcttgatat	tatcacaatt	1380
actgtgcccc	ctgcacttcc	tgctgcaatg	actgctggta	ttgtgtatgc	tcagagaaga	1440
ctgaaaaaaa	tcggtatttt	ctgtatcagt	cctcaaagaa	taaatatattg	tggacagctc	1500
aatcttggtt	gctttgacaa	gactggaaat	ctaactgaag	atgggtttaga	tctttggggg	1560
attcaacgag	tggaaaatgc	acgatttctt	tcaccagaag	aaaatgtgtg	caatgagatg	1620
ttggtaaaat	cccagtttgt	tgcttgatg	gtcacttgct	attcacttac	aaaaattgaa	1680
ggagtgtctc	ctgggtgatcc	acttgatctg	aaaatgtttg	aggctatttg	atggattctg	1740
gaagaagcaa	ctgaagaaga	aacagcactt	cataatcgaa	ttatgcccac	agtggttcgt	1800

cctcccaaac	aactgcttcc	tgaatctacc	cctgcaggaa	accaagaaat	ggagctgttt	1860
gaacttccag	ctacttatga	gatagggaatt	gttcgccagt	tcccattttc	ttctgctttg	1920
caacgtatga	gtgtggttgc	caggggtgctg	ggggatagga	aaatggacgc	ctacatgaaa	1980
ggagcgcccg	aggccattgc	cggctctctgt	aaacctgaaa	cagttcctgt	cgattttcaa	2040
aacgtttttg	aagacttcac	taaacagggc	ttccgtgtga	ttgctcttgc	acacagaaaa	2100
ttggagtcaa	aactgacatg	gcataaagta	cagaatatta	gcagagatgc	aattgagaac	2160
aacatggatt	ttatgggatt	aattataatg	cagaacaaat	taaagcaaaa	aacccttgca	2220
gtacttgaag	atttgcataa	agccaacatt	cgcaccgtca	tggtcacagg	tgacagtatg	2280
ttgactgctg	tctctgtggc	cagagattgt	ggaatgattc	tacctcagga	taaagtgatt	2340
attgctgaag	cattacctcc	aaaggatggg	aaagttgcca	aaataaattg	gcattatgca	2400
gactccctca	cgcagtgcag	tcatccatca	gcaattgacc	cagaggctat	tccggttaaa	2460
ttggtccatg	atagcttaga	ggatcttcaa	atgactcggt	atcattttgc	aatgaatgga	2520
aaatcattct	cagtataact	ggagcatttt	caagaccttg	ttcctaagtt	gatgttgcat	2580
ggcaccgtgt	ttgcccgat	ggcacctgat	cagaagacac	agttgataga	agcattgcaa	2640
aatgttgatt	attttgttgg	gatgtgtggt	gatggcgcaa	atgattgtgg	tgctttgaag	2700
agggcacacg	gaggcatttc	cttatcggag	ctcgaagctt	cagtggcatc	tccctttacc	2760
tctaagactc	ctagtatttc	ctgtgtgcca	aaccttatca	gggaaggccg	tgctgcttta	2820
ataacttcct	tctgtgtgtt	taaattcatg	gcattgtaca	gcattatcca	gtacttcagt	2880
gttactctgc	tgtattctat	cttaagtaac	ctaggagact	tccagtttct	cttcattgat	2940
ctggcaatca	ttttggtagt	ggtattttaca	atgagtttaa	atcctgcctg	gaaagaactt	3000
gtggcacaaa	gaccaccttc	gggtcttata	tctggggccc	ttctcttctc	cgttttgtct	3060
cagattatca	tctgcatttg	atttcaatct	ttgggttttt	tttgggtcaa	acagcaacct	3120
tggtatgaag	tgtggcatcc	aaaatcagat	gcttctaata	caacaggaag	cgggtttttg	3180
aattcttcac	acgtagacaa	tgaaccgaa	cttgatgaac	ataatataca	aaattatgaa	3240
aataccacag	tgttttttat	ttccagtttt	cagtaacctc	tagtggcaat	tgcccttttca	3300
aaaggaaaaa	ccttcaggca	accttgctac	aaaaattatt	tttttgtttt	ttctgtgatt	3360
tttttatata	tttttatatt	attcatcatg	ttgtatccag	ttgcctctgt	tgaccagggt	3420
cttcagatag	tgtgtgtacc	atatcagtg	cgtgtaacta	tgctcatcat	tgttcttgct	3480
aatgcctttg	tgtctatcac	agtggagaac	ttcttccctg	acatggctct	ttggaaagtt	3540
gtgttcaacc	gagacaaaca	aggagagtat	cggttcagca	ccacacagcc	accgcaggag	3600
tcagtggatc	ggtggggaaa	atgctgctta	ccctgggccc	tgggctgtag	aaagaagaca	3660
ccaaaggcaa	agtacatgta	tctggcgcag	gagctcttgg	ttgatccaga	atggccacca	3720
aaacctcaga	caaccacaga	agctaaagct	ttagttaagg	agaatggatc	atgtcaaatc	3780
atcaccataa	catagcagtg	aatcagtcct	agtggtattg	ctgatagcag	tattcaggaa	3840
tatgtgattt	taggagtttc	tgatcctgtg	tgtcagaatg	gcactagttc	agtttatgtc	3900
ccttctgata	tagtagctta	tttgac				3926

<210> 47

<211> 2135

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3046849CB1

<400> 47

cgctcaggcc	cctcttttga	atgctccacg	ccctcctgcg	atctagaacg	attcagggca	60
ggatcctgct	cctgaccatc	tgcgtgcgcg	gcattggtgg	gacttttcag	tttggctata	120
acctctctat	catcaatgcc	ccgaccttgc	acattcagga	attcaccaat	gagacatggc	180
aggcgcgtag	tggagagcca	ctgcccgcac	acctagtcc	gcttatgtgg	tccctcatcg	240
tgtctctgta	tcccctggga	ggcctctttg	gagcactgct	tgccaggtccc	ttggccatca	300
cgctgggaag	gaagaagtcc	ctcctggtga	ataacatctt	tgtggtgtca	gcagcaatcc	360
tgtttggatt	cagccgcaaa	gcaggtctct	ttgagatgat	catgctggga	agactgctcg	420
tgggagtcaa	tgcagggtgtg	agcatgaaca	tccagcccat	gtacctgggg	gagagcgccc	480
ctaaggagct	ccgaggagct	gtggccatga	gctcagccat	ctttacggct	ctggggatcg	540
tgatgggaca	ggtgggtcga	ctcagggagc	tcttaggtgg	ccctcaggcc	tggcccctgc	600
tgctggccag	ctgcctggtg	cccggggcgc	tccagctcgc	ctccctgcct	ctgctccctg	660
aaagcccgcg	ctacctcttc	attgactgtg	gagacaccga	ggcctgctg	gcagcactac	720
ggcagctacg	gggctccggg	gacttggcag	gggagctgga	ggagctggag	gaggagcgcg	780
ctgcctgcca	gggctgccgt	gcccggcgcc	catgggagct	gttccagcat	cgggccctga	840
ggagacaggt	gacaagcctc	gtggttcttg	gcagtgccat	ggagctctgc	gggaatgact	900
cgggtgtacgc	ctacgcctcc	tccgtgttcc	ggaaggcagg	agtgccggaa	gcgaagatcc	960
agtacgcgat	categggagc	gggagctgcg	agctgctcac	ggcggttgtt	agttgtgtgg	1020
taatcgagag	tgtgggtcgg	cgcgtgctgc	tcatcggtgg	gtacagcctg	atgacctgct	1080
gggggagcat	cttcactgtg	gccctgtgcc	tgcagagctc	cttcccctgg	acactctacc	1140
tggccatggc	ctgcactctt	gccttcaccc	tcagcttttg	cattggccct	gccggagtga	1200

cggggatacct	ggccacagag	ctgtttgacc	agatggccag	gcctgctgcc	tgcattggtct	1260
gcgggggcgct	catgtggatc	atgtctcatcc	tggtcgccct	gggatttccc	tttatcatgg	1320
aggccttgctc	ccacttcctc	tatgtccctt	tccttggtgt	ctgtgtctgt	ggggccatct	1380
acactggcct	gttccttccct	gagaccaaag	gcaagacctt	ccaagagatc	tccaaggaat	1440
tacacagact	caacttcccc	aggcggggccc	aggggccccc	gtggaggagc	ctggagggtta	1500
tccagtcaac	agaactctag	tcccaaagggt	gtggccagag	ccaaagccag	ctactgtcct	1560
gtcctctgct	tcctgccagg	gccctgggtcc	tcactccctc	ctgcattcct	catttaagga	1620
gtgtttattg	agcacccctt	gtgtgcagac	atggctccag	gtgcttagca	atcaatggtg	1680
agcgtgggtat	tccaggctaa	aggtaattaa	ctgacagaaa	atcagtaaca	acataattac	1740
aggctgggtt	tggcagctca	tgactgtaat	cccagcaact	tgggaggcca	agggtgggagg	1800
atcaattgag	gccagagttt	gaaaccagcc	taggtaacat	agttagagccc	cctatctcta	1860
caaaaaattt	taaacattag	ctgggcagtg	tggtatgtgc	taacagctct	agctactcag	1920
gaggctgagg	cagcaggatc	acttgagtc	caagagttca	aggtagcagt	aagctaacaa	1980
ttcacaccac	tgcattgcca	gactgggggtg	acagagggag	acttcatctc	tttaaaaaca	2040
taataataat	aattacggac	tccggaaatg	cgttgacaac	gaaacatacc	ggtggccccg	2100
tgagggtggtg	atccccgtatc	ccagccttgg	gaagc			2135

<210> 48

<211> 2637

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4538363CB1

<400> 48

atgggctgga	gatgccactg	tccgcttggt	ttaatgatca	atgagctccc	tgccaggaaa	60
ccctttctga	cctggtttgc	ccctcagtc	ctcgggctca	tacctagtgc	ctgcggcagg	120
acagccatgg	ccgccaactc	caccagcgac	ctccacactc	ccgggacgca	gctgagcgtg	180
gctgacatca	tcgtcatcac	tgtgtatttt	gctctgaacg	tggccgtggg	catatggctc	240
tcttgtcggg	ccagtaggaa	cacgggtgaat	ggctacttcc	tggcaggccg	ggacatgacg	300
tgggtggccga	ttggagcctc	cctcttcgcc	agcagcgagg	gctctggcct	cttcatttga	360
ctggcgggct	caggcgcggc	aggaggctctg	gccgtggcag	gcttcgagtg	gaatgccacg	420
tacgtgctgc	tggcactggc	atgggtgttc	gtgcccctct	acatctcctc	agagatcgtc	480
accttacctg	agtacattca	gaagcgctac	gggggccagc	ggatccgcat	gtacctgtct	540
gtcctgtccc	tgtactgttc	tgtcttcacc	aagatatcgc	tggacctgta	cgcgggggct	600
ctgtttgtgc	acatctgcct	gggctggaac	ttctacctct	ccaccatcct	cacgctcggc	660
atcacagccc	tgtacaccat	cgcagggggc	ctggctgctg	taatctacac	ggacgccttg	720
cagacgctca	tcattggtgg	gggggctgtc	atcctgacaa	tcaaagcttt	tgaccagatc	780
ggtggttacg	ggcagctgga	ggcagcctac	gcccaggcca	ttccctccag	gacctatgcc	840
aacaccacct	gccacctgcc	acgtacagac	gccatgcaca	tgtttcgaga	ccccacaca	900
ggggacctgc	cgtggaccgg	gatgaccttt	ggcctgacca	tcattggccac	ctggtactgg	960
tgcaccgacc	aggctcatcgt	gcagcgatca	ctgtcagccc	gggacctgaa	ccatgccaag	1020
gcgggctcga	tccttgcagc	ctacctcaag	atgctcccca	tgggcctgat	catcatgccg	1080
ggcatgatca	gccgcgcatt	gttcccagat	gatgtgggct	gcgtgggtgc	gtccgagtgc	1140
ctgcgggcct	gcggggccga	ggtcggctgc	tccaacatcg	cctaccccaa	gctgggtcatg	1200
gaactgatgc	ccatcggtct	gcgggggctg	atgatcgcag	tgatgctggc	ggcgctcatg	1260
tcgtcgtcga	cctccatctt	caacagcagc	agcaccctct	tcactatgga	catctggagg	1320
cggctgcgtc	ccgctccggg	cgagcgggag	ctcctgctgg	tgggacggct	ggtcatagtg	1380
gcactcatcg	gcgtgagtgt	ggcctggatc	ccgctcctgc	aggactccaa	cagcgggcaa	1440
ctcttcatct	acatgcagtc	agtgaccagc	tccttggtcc	caccagtgc	tgcagtcttt	1500
gtcctgggctg	tcttctggcg	acgtgccaac	gagcaggggg	ccttctgggg	cctgatagca	1560
gggctgggtgg	tggggggccac	gaggctggtc	ctggaattcc	tgaaccagc	cccaccgtgc	1620
ggagagccag	acacgcggcc	agccgtcctg	gggagcatcc	actacctgca	cttcgctgtc	1680
ggcctctttg	cactcagttg	tgtgttgtg	gtggctggaa	gcctgctgac	cccaccccca	1740
cagagtgtcc	agattgagaa	ccttacctgg	tggaccctgg	ctcaggatgt	gcccttggga	1800
actaaagcag	gtgatggcca	aacaccccag	aaacacgcct	tctgggcccg	tgtctgtggc	1860
ttcaatgcca	tctcctcat	gtgtgtcaac	atattctttt	atgcctactt	cgctcgaaca	1920
ctgccatcct	ggacagaag	gcaggagctc	ttagtctctc	ggtccaccca	tttccctcat	1980
ggggatcccc	aggccccaag	aggggcagat	tccctcaca	gctgcacagc	agctcgggtg	2040
ccaagaactg	gccaagccag	caaagcggga	gcctgaaaac	attagggggg	aaactgggac	2100
gaaacataag	tgtgactttt	tccaaacaac	agcaccacaa	gcaagtcaag	catttgggac	2160
gcgacaaact	tagattttcc	tgaccggggc	caccacaccc	caacctcttc	acctcccaa	2220
ctaccaaac	agctcacac	catactcaca	ccaccacag	cggcccggcc	ccactccaat	2280
cagaaaggca	ccccccact	ctcaagacgc	gacggcgcaa	tcgactgcaa	ctccataacg	2340
atgccaaaac	gacacaagcc	aggacacggc	actgtatata	gcacgagggt	gatctgcaac	2400

gttgtggccg	aatgcagaaa	atacactggg	tgctggcgta	aggaagatcc	gcgagtaaac	2460
aacgggtctt	taaaacttact	gcatccacca	aggtacactt	ccagaacgag	accagacaac	2520
tacactccac	acaacctgca	gccacaccct	atctctgcta	tcataaagag	ccccgcacc	2580
acataataat	gccggcagac	tcagtgcgcg	aaacccttgt	gctggacttc	accacgg	2637

<210> 49

<211> 3783

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6427460CB1

<400> 49

gcactagtag	cccggagccc	atgggcgcg	cgagccgggc	gcggggggcg	tgaacggcgg	60
agcgggagcg	gcccggaggag	ccatggactg	cagcctcgctg	cggacgctcg	tgcacagata	120
ctgtgcagga	gaagagaatt	gggtggacag	caggaccatc	tacgtgggac	acagggagcc	180
acctccgggc	gcagaggcct	acatcccaca	gagataccca	gacaacagga	tcgtctcgtc	240
caagtacaca	ttttggaact	ttatacccaa	gaatttattt	gaacaattca	gaagagtagc	300
caacttttat	ttcctttatc	tatttctggt	gcagttgatt	attgatacac	ccacaagtc	360
agtgcacaag	ggactttccac	tcttctttgt	cattactgtg	acggctatca	aacagggtta	420
tgaagactgg	cttcgcacata	aagcagacaa	tgccatgaac	cagtgtcctg	ttcatttcat	480
tcagcacggc	aagctcgctt	ggaaacaaa	tcgaaagctg	cgagttgggg	acattgtcat	540
ggttaaggag	gacgagacct	ttccctgcga	cttgatcttc	ctttccagca	accggggaga	600
tgggacgtgc	cacgtcacca	ccgccagctt	ggatggagaa	tcagccata	aaacgcatta	660
cgcggtccag	gacaccaaag	gcttccacac	agaggaggat	atcggcggac	ttcacgccac	720
catcgagtgt	gagcagcccc	agcccagcct	ctacaagttc	gtgggtcgca	tcaacgttta	780
cagtgcactg	aatgaccccc	tggtgaggcc	cttaggatcg	gaaaacctgc	tgcttagagg	840
agctacactg	aagaacactg	agaaaatctt	tggtgtggct	atttacacgg	gaatggaaac	900
caagatggca	ttaaattatc	aatcaaaatc	tcagaagcga	tctgccgtgg	aaaaatcgat	960
gaatgcgttc	ctcattgtgt	atctctgcat	tctgatcagc	aaagccctga	taaacactgt	1020
gctgaaatac	gtgtggcaga	gtgagccctt	tcgggatgag	ccgtgggtata	atcagaaaac	1080
ggagtcggaa	aggcagagga	atctgttcct	caaggcatte	acggacttcc	tggtcttcat	1140
ggtcctcttt	aactacatca	tccctgtgtc	catgtacgtc	acggtcgaga	tgcagaagtt	1200
cctcggctct	tacttcatca	cctgggacga	agacatgttt	gacgaggaga	ctggcgaggg	1260
gcctctgggt	aacacgtcgg	acctcaatga	agagctggga	caggtggagt	acatcttcac	1320
agacaagacc	ggcaccctca	cggaacacaa	catggagttc	aaggagtgtc	gcacggaagg	1380
ccatgtctac	gtgccccacg	tcacttgcaa	cgggcagggtc	ctcccagagt	cgtcaggaa	1440
cgacatgatt	gactcgtccc	ccagcgtcaa	cgggaggagg	cgcgaggagc	tggttttccg	1500
ggccctctgt	ctctgccaca	ccgtccaggt	gaaagacgat	gacagcgtag	acggccccag	1560
gaaatcgccg	gacgggggga	aatcctgtgt	gtacatctca	tcctcgcccc	acgaggtggc	1620
gctggtcgaa	ggtgtccaga	gacttggctt	tacctacct	aggctgaagg	acaattacat	1680
ggagatatta	aacaggggaga	accacatcga	aaggtttgaa	ttgctggaaa	ttttgagttt	1740
tgactcagtc	agaaggagaa	tgagtgtaat	tgtaaaatct	gctacaggag	aaatttatct	1800
gttttgcaaa	ggagcagatt	cttcgatatt	cccccgagt	atagaaggca	aagttgacca	1860
gatecgagcc	agagtggagc	gtaacgcagt	ggagggggtc	cgaactttgt	gtgttgctta	1920
taaaaggctg	atccaagaag	aatatgaagg	catttgttaag	ctgctgcagg	ctgccaaagt	1980
ggcccttcaa	gatecgagaga	aaaagtttag	agaagcctat	gagcaaata	agaaagatct	2040
tactctgctt	ggtgctacag	ctgttgagga	ccggctgcag	gagaaagctg	cagacaccat	2100
cgaggccctg	cagaaggccg	ggatcaaagt	ctgggttctc	acgggagaca	agatggagac	2160
ggcgcgggcc	acgtgctacg	cctgcaagct	cttcgcgagg	aacacgcagc	tgctggagct	2220
gaccaccaag	aggatcgagg	agcagagcct	gcacgacgtc	ctgttcgagc	tgagcaagac	2280
ggtcctgcgc	cacagcggga	gctgaccagg	agacaacctc	tcgggacttt	cagcagatat	2340
gcaggactac	ggtttaatta	tcgacggagc	tgactgtct	ctgataatga	agcctcgaga	2400
agacgggagt	tcgggcaact	acaggggagt	cttcttgga	atctgccgga	gctgcagcgc	2460
ggtgctctgc	tgccgcagtg	cgcccttgca	gaaggctcag	attgttaaat	taatcaaatt	2520
ttcaaaagag	cacccaatca	cgtagcaat	tggcgatgg	gcaaatgatg	tcagcatgat	2580
tctggaagcg	cacgtgggca	taggtgtcat	cggcaaggaa	ggcgcggcag	ctgccaggaa	2640
cagcgactat	gcaatcccaa	agtttaagca	tttgaagaag	atgctgcttg	ttcaccggca	2700
tttttattac	attaggatct	ctgagctcgt	gcagtacttc	ttctataaga	acgtctgctt	2760
catcttccct	cagtttttat	accagttctt	ctgtgggttt	tcacaacaga	ctttgtacga	2820
caccgcgtat	ctgaccctct	acaacatcag	cttcacctcc	ctccccatcc	tcctgtacag	2880
cctcatggag	cagcatgttg	gcattgacgt	gctcaagaga	gacccgaccc	tgtacaggga	2940
cgctgcacaag	aatgtccctg	tgcgctggcg	cgtgttcac	tactggagcg	tcctgggact	3000
gtttgacgca	ctggtgttct	tctttgggtg	ttatttcgtg	tttgaaaata	caactgtgac	3060
aagcaacggg	cagatatattg	gaaactggac	gtttggaacg	ctggtattca	ccgtgatgg	3120

gttcacagtt	acactaaagc	ttgcattgga	cacacactac	tggacttggga	tcaaccattt	3180
tgatcatctgg	gggtcgctgc	tggtctacgt	tgtcttttca	cttctctggg	gaggagtgat	3240
ctggcggttc	ctcaactacc	agaggatgta	ctacgtgttc	atccagatgc	tgtccagcgg	3300
gcccgccttg	ctggccatcg	tgtgtctggt	gaccatcagc	ctccttcccg	acgtcctcaa	3360
gaaagtccctg	tgccggcagc	tgtgggccaac	agcaacagag	agagtccagc	agaatgggtg	3420
cgcacagcct	cgggaccgcg	actcagaatt	cacctctctt	gcctctctgc	agagcccagg	3480
ctaccagagc	acctgtccct	cggccgcctg	gtacagctcc	cactctcagc	aggtgacact	3540
cgccggcctgg	aaggagaagg	tgtccacgga	gccccaccc	atcctcggcg	gttcccatca	3600
ccactgcagt	tccatcccaa	gtcacagctg	ccctaggtcc	cgtgtgggaa	tgctcgtgtg	3660
atggatgggc	ctaagcctgt	ggagactgtg	cacgtgcctc	ttcctggccc	ccagcaggca	3720
aggagggggg	tcacaggcct	tgccctcgaa	catggcaccc	tgcccgccctg	gaccagcac	3780
tgt						3783

<210> 50

<211> 2105

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474127CB1

<400> 50

ccagcgccca	gggaagcggc	tcaaccacct	gaatccggaa	aacgccaaaca	agtagtttct	60
cgctcggagaa	ggcggtctca	cctgggcgcc	aagactcagt	cccgtgccc	agagaacctc	120
gtccactcgg	aaaccaaagc	agaaccactt	ttctctcggt	ctcgttaagt	catgtctgag	180
tcacagagat	gggcaagatc	gagaacaacg	agaggggtgat	cctcaatgtc	gggggcaccc	240
ggcacgaaac	ctaccgcagc	accctcaaga	ccctgcctgg	aacacgcctg	gcccttcttg	300
cctcctccga	gccccaggc	gactgcttga	ccacggcggg	cgacaagctg	cagccgtcgc	360
cgcctccact	gtcgccgcgc	ccgagagcgc	ccccgtgtc	ccccggggcca	ggcggctgct	420
tcgaggcgcg	cgcgggcaac	tgcagttccc	gcggcggcag	ggccagcgac	catcccgtg	480
gcggccgcga	gttcttcttc	gaccggcacc	cgggcgtctt	cgcctatgtg	ctcaattact	540
accgcaccgg	caagctgcac	tgccccgcag	acgtgtgcgg	gcccgtcttc	gaggaggagc	600
tggcctttctg	gggcatcgac	gagaccgacg	tggagccctg	ctgctggatg	acctaccggc	660
agcacccgca	gcgcgaggag	gcgctggaca	tcttcgagac	ccccgacctc	attggcggcg	720
accccgcgca	cgacgaggac	ctggcgggcca	agaggctggg	catcgaggag	cgcggcgggc	780
tcggggggccc	cgacggcaaa	tctggccgct	ggaggaggct	gcagccccgc	atgtggggccc	840
tcttcgaaga	cccctactcg	tccagagccg	ccaggtttat	tgtttttgct	tctttattct	900
tcctcctggg	ttcaattaca	actttttgcc	tggaaacaca	tgaagctttc	aatattgtta	960
aaaacaagac	agaaccagtc	atcaatggca	caagtgttgt	tctacaatat	gaaattgaaa	1020
cagatcctgc	cttgacgtat	gtagaaggag	tgtgtgtggt	gtggtttact	tttgaatttt	1080
tagtccgtat	tgttttttca	cccaataaac	ttgaattcat	caaaaatctc	ttgaatatca	1140
ttgactttgt	ggccatccta	cctttctact	tagagggtgg	actcagtggg	ctgtcatcca	1200
aagctgctaa	agatgtgctt	ggcttctcta	gggtggtaag	gtttgtgagg	atcctgagaa	1260
ttttcaactg	caccgcgat	tttgtaggtc	tgagggtgct	tggacatact	cttcgagcta	1320
gtactaatga	atttttgctg	ctgataattt	tcttggtctt	aggagttttg	atatttgcta	1380
ccatgatcta	ctatgccgag	agagtgggag	ctcaacctaa	cgacccttca	gctagtgagc	1440
acacacagtt	caaaaacatt	cccattgggt	tctggtgggc	tgtagtgacc	atgactaccc	1500
tgggttatgg	ggatatgtac	ccccaaacat	ggtcaggcat	gctggtggga	gcctgtgtgt	1560
ctctggctgg	agtgtgaca	atagccatgc	cagtgcctgt	catttgtcaat	aattttggaa	1620
tgtactactc	cttggaatg	gcaaagcaga	aacttccaag	gaaaagaaag	aagcacatcc	1680
ctcctgtctc	tcaggcaagc	tcacctactt	tttgcaagac	agaattaaat	atggcctgca	1740
atagtacaca	gagtacaca	tgtctgggca	aagacaatcg	acttctggaa	cataacagat	1800
cagtgttatc	aggtgacgac	agtacaggaa	gtgagccgcc	actatcacc	ccagaaaggc	1860
tccccatcag	acgctctagt	accagagaca	aaaacagaag	aggggaaaca	tgtttcttac	1920
tgacgacagg	tgattacacg	tgtgcttctg	atggagggat	caggaaagg	tatgaaaaat	1980
cccgaagctt	aaacaacata	gcgggcttgg	caggcaatgc	tctgaggctc	tctccagtaa	2040
catcaccta	caactctcct	tgctctctga	ggcgctctcg	atctcccatc	ccatctatct	2100
tgtaa						2105

<210> 51

<211> 2069

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7476949CB1

<400> 51

atgagcaagg	acctggcagc	aatggggcct	ggagcttcag	gggacggggt	caggactgag	60
acagctccac	acatagcact	ggactccaga	gttgggtctgc	acgcctacga	catcagcgtg	120
gtgggtcatct	actttgtctt	cgctattgct	gtggggatct	ggctcgtccat	ccgtgcaagt	180
cgagggacca	ttggcggtta	tttccctggcc	gggaggtcca	tgagctgggtg	gccaattgga	240
gcattctctga	tgtccagcaa	tgtgggcagt	ggcttgttca	tggcctgggc	tgggacaggg	300
gctgccggag	gccttgcctg	agggtggcttc	gagtggaacg	caacctggct	gctcctggcc	360
cttggctggg	tcttcgtccc	tgtgtacatc	gcagcagggtg	tggtcacaat	gccgcagtat	420
ctgaagaagc	gattttggggg	ccagaggatc	caggtgtaca	tgtctgtcct	gtctctcatc	480
ctctacatct	tcaccaagat	ctcgactgac	atcttctctg	gagccctctt	catccagatg	540
gcattgggct	ggaacctgta	cctctccaca	gggatcctgc	tgggtggtgac	tgccgtctac	600
accattgacg	gtggcctcat	ggccgtgatc	tacacagatg	ctctgcagac	ggtgatcatg	660
gtagggggag	ccctggctct	catgtttctg	ggctttcagg	acgtgggctg	gtaccagggc	720
ctggagcagc	ggtacaggca	ggccatccac	aatgtcacag	tccccaacac	cacctgtcac	780
ctcccacggc	ccgatgtctt	ccacattctt	cgggaccctg	tgagcgggga	catcccttgg	840
ccaggtctca	ttttcgggct	cacagtgtctg	gccacctggg	gttgggtgcac	agaccagggtc	900
atttgtgcagc	ggtctctctc	ggccaagagt	ctgtctcatg	ccaagggagg	ctccgtgctg	960
gggggctacc	tgaagatcct	ccccatgttc	ttcatcgtca	tgcctggcat	gatcagccgg	1020
gccctgttcc	cagacgaggt	gggctgcgtg	gacctgatg	tctgccaaag	aatctgtggg	1080
gcccagtggtg	gatgttccaa	cattgcctac	cctaagttgg	tcatggccct	catgcctgtt	1140
ggtctgcggg	ggctgatgat	tgccgtgatc	atggccgctc	tcatgagctc	actcacctcc	1200
atcttcaaca	gcagcagcac	cctgttcacc	attgatgtgt	ggcagcgctt	ccgcaggaag	1260
tcaacagagc	aggagctgat	ggtgggtggg	agagtgtttg	tgggtgttct	ggttgtcatc	1320
agcatcctct	ggatccccat	catccaaagc	tccaacagtg	ggcagctctt	cgactacatc	1380
caggctgtca	ccagttacct	ggccccaccc	atcaccgctc	tcttctctgt	ggccatcttc	1440
tgcaagaggg	tcacagagcc	cggagctttc	tggggcctcg	tgtttggcct	gggagtgggg	1500
cttctgcgta	tgatcctgga	gttctcatac	ccagcgccag	cctgtgggga	ggtggaccgg	1560
aggccagcag	tgctgaagga	cttccactac	ctgtactttg	caatcctcct	ctgcgggctc	1620
actgccatcg	tcattgtcat	tgtcagcctc	tgtacaactc	ccatccctga	ggaacagctc	1680
acacgcctca	catggtggac	tcggaactgc	ccccctctctg	agctggagaa	ggaggcccac	1740
gagagcacac	cggagatatac	cgagaggcca	gccggggagt	gccctgcagg	aggtggagcg	1800
gcagagaact	cgagcctggg	ccaggagcag	cctgaagccc	caagcagggtc	ctggggaaaag	1860
ttgctctgga	gctggttctg	tgggctctct	ggaacacogg	agcaggccct	gagcccagac	1920
gagaaggctg	cgctagaaca	gaagctgaca	agcattgagg	aggagccact	ctggagacat	1980
gtctgcaaca	tcaatgctgt	ccttttgcgtg	gccatcaaca	tcttctctctg	gggctatattt	2040
gcgtgattca	aacctggcctt	cactgtaga				2069

<210> 52

<211> 4245

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477249CB1

<400> 52

gcggcggcag	getcagctgc	gccggggcggg	ggcgggcgctg	gggcccgcgc	tgtaggactc	60
ggggccgacg	ccgcgggatg	gggacgcggc	ggggggagtg	aggcagtggc	ggcgggcgcg	120
gtaagcggaa	cttcggcccc	aggggctcgc	ccgctcccgc	ctctgtcttg	tgggcctcca	180
cctgcagccc	cgcgcccccc	gcgccccgcg	ggacccgagc	ggcgacgacg	ggggaatgtg	240
gcgctggatc	gggcagcagc	tgggttttga	cccaccacat	cagagtgaac	caagaacctc	300
ctacgtagcc	aacaggtttc	ctcagaatgg	cctttacaca	cctcagaaat	ttatagataa	360
caggatcatt	tcattctaagt	acactgtgtg	gaattttgtt	ccaaaaaatt	tatttgaaca	420
gttcagaaga	gtggcaaaact	tttattttct	tattatattt	ttggttcagc	ttatgattga	480
tacacctacc	agtccagttta	ccagtggact	tccattatct	tttgtgataa	cagtaactgc	540
cataaagcag	ggatatgaag	attggttacg	gcataactca	gataatgaag	taaatggagc	600
tcctgtttat	gttggtcgaa	gtgggtggcct	tgtaaaaaact	agatcaaaaa	acattcgggt	660
gggtgatatt	gttcgaatag	ccaaagatga	aatttttctc	gcagacttgg	tgcctctgtc	720
ctcagatcga	ctggatgggt	cctgtcacgt	tacaactgct	agtttggacg	gagaaactaa	780
cctgaagaca	catgtggcag	ttccagaaac	agcattatta	caaacagttg	ccaatttggg	840
cactctagta	gctgtaatag	aatgccagca	accagaagca	gacttatata	gattcatggg	900
acgaatgatc	ataaccaaac	aaatggaaga	aattgttaaga	cctctggggc	cggagagtct	960
cctgcttcgt	ggagccagat	taaaaaacac	aaaagaaatt	tttgtgtgtg	cgggtatacac	1020
tggaatggaa	actaagatgg	cattaaatta	caagagcaaa	tcacagaaac	gatctgcagt	1080

```

agaaaagtca atgaatacat ttttgataat ttatctagta attcttataat ctgaagctgt 1140
catcagcact atcttgaagt atacatggca agctgaagaa aaccttggta 1200
taacccaaaa acagaacatc aaagaaatag cagtaaggta gagtacgtgt ttacagataa 1260
aactggtaca ctgacagaaa atgagatgca gtttcgggaa tgttcaatta atggcatgaa 1320
ataccaagaa attaatggta gacttgtacc cgaaggacca acaccagact cttcagaagg 1380
aaacttatct tatcttagta gtttatccca tcttaacaac ttatcccatc ttacaaccag 1440
ttcctctttc agaaccagtc ctgaaaatga aactgaacta attaaagaac atgatctctt 1500
ctttaaagca gtcagtctct gtcacactgt acagattagc aatgttcaaa ctgactgcac 1560
tggtgatggt ccctggcaat ccaacctggc accatcgagc ttggagtact atgcatcttc 1620
accagatgaa aaggctctag tagaagctgc tgcaaggatt ggtatttgtt ttattggcaa 1680
ttctgaagaa actatggagg ttaaaactct tggaaaactg gaacggtaca aactgcttca 1740
tattctggaa ttgtattcag atcgtaggag attgttcagg cacttcagg 1800
tgagaagtta ttatttgcta aaggagctga gtcatcaatt ctccctaaat gtataggttg 1860
agaaatagaa aaaaccagaa ttcattgtaga tgaatttgct ttgaaagggc taagaactct 1920
gtgtatagca tatagaaaat ttacatcaaa agagtatgag gaaatagata aacgcatatt 1980
tgaagccagg actgccttgc agcagcggga agagaaattg gcagctgttt tcagattcat 2040
agagaaagac ctgatattac ttggagccac agcagtagaa gacagactac aagataaagt 2100
tcgagaaact attgaagcat tgagaatggc tggatatcaa gtatgggtac ttactgggga 2160
taaacatgaa acagctgtta gtgtgagttt atcatgtggc cattttcata gaaccatgaa 2220
catccttgaa cttataaacc agaaatcaga cagcgagtgt gctgaacaat tgaggcagct 2280
tgccagaaga attacagagg atcatgtgat tcagcatggg ctggtagtgg atgggaccag 2340
cctatctctt gcactcaggg agcatgaaaa actatttatg gaagtttgca gaaattgttc 2400
agctgtatta tgctgtcgta tggctccact gcagaaagca aaagtaataa gactaataaa 2460
aatatcacct gagaaaccta taacattggc tgttgggtgat ggtgctaatt acgtaagcat 2520
gatacaagaa gcccatgttg gcataggaat catgggtaaa gaaggaagac aggctgcaag 2580
aaacagtgc atgcaatag ccagatttaa gttcctctcc aaattgcttt ttgttcattg 2640
tcatttttat tatattagaa tagctaccct tgtacagtat tttttttata agaattgttg 2700
ctttatcaca cccagtttt tatatcagtt ctactgtttg ttttctcagc aaacattgta 2760
tgacagcgtg tacctgactt tatacaatat ttgttttact tccctaccta ttctgatata 2820
tagtcttttg gaacagcatg tagaccctca tgtgttacaa aataagccca ccctttatcg 2880
agacattagt aaaaaccgcc tcttaagtat taaacatttt ctttattgga ccatcctggg 2940
cttcagtcac gccctttattt tcttttttgg atcctatttt ctaataggga aagatacatc 3000
tctgcttgga aatggccaga tgtttggaaa ctggacattt ggcacttttg tcttcacagt 3060
catggttatt acagtccag ctattatatt tctggaaact cttttttgga cttggatcaa 3120
ccatctcgtt ccctggggat ctattatatt ttattttgta ttttcttgt tttatggagg 3180
gattctctgg ccatttttgg gctcccagaa tatgtatttt gtgtttatc agctcctgtc 3240
aagtggttct gcttggtttg ccataatcct catggttggt acatgtctat ttcttgatat 3300
cataaagaag gtctttgacc gacacctcca cctacaagt actgaaaagg cacagcttac 3360
tgaacaaaat gcaggtatca agtgcttgga ctccatgtgc tgtttcccg aaggagaagc 3420
agcgtgtgca tctgttgga gaatgctgga acgagttata ggaagatgta gtccaacca 3480
catcagcagg tgtgaaatct ctctaagtag ccttctgtgc agatgagtat cctatctgga 3540
acaggatgaa cctgccgctc tagataccta ataaatcagc agctggtttt accaactgaa 3600
gcaggaagtc tgctatttat tagcactctt tgggtgtaga tttcactttg tggctttggg 3660
gtaagggctt tttcactcac aaaggaagag aaagcacctt tgaagagact tcattctaag 3720
aacaataaat tttgtttcat aatctttcta aaatgtgctc agtaggagtg tcttctatg 3780
actcttttat ggtttgtata actttctttt ttaaattata catatactat ttctttttta 3840
tttttttaaa atttttttgc tttttgtctt taaaaataa tctcaacata acagtgaagt 3900
caaaggcttt cttttcttta ctctgtatgt atattttcca gttggttatt tgaggctttg 3960
aggatattat aaacacaaaa ggctgtattt ctgctccctt acctcttctt atgtctgtaa 4020
tgaagttttg aaatgagtca tgatttttaa gtttcttttg cttggtattt attgctaata 4080
taaaagtgtg tgagttagaa caggcttttt aaattatgga gtaaaagaat ctttagcattt 4140
ttgtcccttc ctaaactctg ttcttgaatg agatttatca ccatgcctgc tgttgtgcac 4200
cataacgaaa aaaaacacct tttggtaaac accattttaa attca 4245

```

<210> 53
 <211> 2124
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7477720CB1

```

<400> 53
atggctctgc agatgttcgt gacttacagt ccttggaaatt gtttgctact gctagtggct 60
cttgagtgtt ctgaagcatc ttctgatttg aatgaatctg caaattccac tgctcagtat 120
gcactctaag cttgggttgc tgctgccagc tcagagccag aggaagggat atctgttttt 180

```


gaactggatt	atgactatgt	gcaaattcct	tatgagggtca	ctctctggat	acttctagca	240
tcccttgcaa	aaataggctt	ccacctctac	cacaggctgc	caggcctcat	gccagaaagc	300
tgccctctca	tcctggtggg	ggcgctggtg	ggcgggcatca	tcttcggcac	cgaccacaaa	360
tcgcctccgg	tcattggactc	cagcatctac	ttcctgtatc	tcctgccacc	catcgttctg	420
gagggcggt	acttcatgcc	cacccggccc	ttctttgaga	acatcggtct	catcctgtgg	480
tgggcagtat	tgggggccc	gatcaacgcc	ttggggcattg	gcctctccct	ctacctcatc	540
tgccagggtga	aggcctttgg	cctggggcag	gtcaacctgc	tgcaaacct	gctgttcggc	600
agcctgatct	ccgcctggga	cccagtgggc	gtgctagccg	tgtttgagga	agcgcgctg	660
aacgagcagc	tctacatgat	gatctttggg	gaggccctgc	tcaatgatgg	cattactgtg	720
gtcttataca	atatgttaat	tgccctttaca	aagatgcata	aatttgaaga	catagaaact	780
gtcgacattt	tggctggatg	tgccccgattc	atcgttgtgg	ggcttggagg	ggtattgttt	840
ggcatcgttt	ttggatttat	ttctgcattt	atcacacgtt	tcactcagaa	tatctctgca	900
attgagccac	tcactgtctt	catgttcagc	tatttgtctt	acttagctgc	tgaaaccttc	960
tatctctccg	gcactcctggc	aatcacagcc	tgccgcagtaa	caatgaaaaa	gtacgtggaa	1020
gaaaacgtgt	cccagacatc	atacacgacc	atcaagtact	tcatgaagat	gctgagcagc	1080
gtcagcgaga	ccttgatctt	catcttcatg	ggtgtgtcca	ctgtgggcaa	gaatcacgag	1140
tggaaactggg	ccttcaccctg	gccttctgcc	gccttctgcc	aaatctggag	agccatcagc	1200
gtatttgcct	tctttctatat	cagtaaccag	tttcggactt	tccccctctc	catcaaggac	1260
cagtgcataca	ttttctacag	tgggtgttcca	ggagctggaa	gttttttact	tgcatttttg	1320
cttccctctgt	ctctttttcc	taggaagaaa	atgtttgtca	ctgctactct	agtagttata	1380
tacttttactg	tattttattca	gggaatcaca	gttggccctc	tggtcaggta	cctggatgtt	1440
aaaaaaacca	ataaaaaaga	atccatcaat	gaagagcttc	atattcgtct	gatggatcac	1500
ttaaaggctg	gaatcgaaga	tgtgtgtggg	cactggagtc	actaccaagt	gagagacaag	1560
tttaagaagt	ttgatcatag	atacttacgg	aaaatcctca	tcagaaagaa	cctaccctaa	1620
tcaagcattg	tttctttgta	caagaagctg	gaaatgaagc	aagccatcga	gatggtggag	1680
actgggatac	tgagctctac	agctttctcc	ataccctcat	aggcccagag	gatacaagga	1740
atcaaaaagac	tttccccctga	agatgtggag	tccataaggg	acattctgac	atccaacatg	1800
taccaagtct	ggcaaaggac	cctgtcctac	aacaaataca	acctcaaacc	ccaaacaagt	1860
gagaagcagg	ctaaagagat	tctgatccgc	cgccagaaca	ccttaagggg	gagcatgagg	1920
aaaggtcaca	gcctgccctg	gggaaagccg	gctggcacca	agaatatccg	ctacctctcc	1980
taccctctacg	ggaatcctca	gtctgcagga	agagacacaa	gggctgctgg	gttctcaggt	2040
aagctgcccc	ctgggtgctt	ctgctgcttt	tctgtagagt	caggtggtaa	atatctgggg	2100
gtgtggggcca	agaggcaaca	ttaa				2124

<210> 54

<211> 2195

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477852CB1

<400> 54

atgggggggtt	ttctacctaa	ggcagaaggg	cccgaggacc	aactccagaa	acttctgccc	60
tcctttcttg	tcagagaaca	agactgggac	cagcacctgg	acaagcttca	tatgctgcag	120
cagaagagga	ttctagagtc	tccactgctt	cgagcatcca	aggaaaatga	cctgtctgtt	180
cttaggcaac	ttctactgga	ctgcacctgt	gacgttcgac	aaagaggagc	cctggggggag	240
acggcgctgc	acatagcagc	cctctatgac	aacttgaggg	cggccttggg	gctgatggag	300
gtgccccag	agctggtctt	tgagcccacc	acatgtgagg	cttttgaggg	tcagactgca	360
ctgcacatcg	ctgttgtgaa	ccagaatgtg	aacctgggtg	gtgccctgct	caccgcagg	420
gccagtgtct	ctgccagagc	cacaggcact	gccttccgcc	gtagtccccg	caacctcatc	480
tactttgggtg	agcacccttt	gtcctttgct	gcctgtgtga	acagcgagga	gatcgtgcgg	540
ctgctcattg	agcatggagc	tgacatcagg	gccaggact	ccctgggtaa	cacagtatta	600
cacatcctca	ctctccagcc	caacaaaacc	tttgcctgcc	agatgtacaa	cctgtgctg	660
tcctacgatg	gacatgggga	ccacctgcag	cccctggacc	ttgtgcccac	tcaccagggt	720
ctcaccctct	tcaagctggc	tggagtggag	ggtaacactg	tgatgttcca	gcacctgatg	780
cagaagcgga	ggcacatcca	gtggacgtat	ggacccctga	cctccattct	ctacgacctc	840
acagagatcg	actcctgggg	agaggagctg	tccttctctg	agcttgtggg	ctcctctgat	900
aaacgagagg	ctcgcaaat	tctggaacag	acccagtgga	aggagctggg	gagcttcaag	960
tggaaacaagt	atggccggcc	gtacttctgc	atcctggctg	ccttgtacct	gctctacatg	1020
atctgtctta	ccacgtgctg	cgtctaccgc	ccccttaagt	ttcgtgggtg	caaccgcact	1080
cattctcgag	acatcaccat	cctccagcaa	aaactactac	aggaggccta	tgagacacgt	1140
gaagatatca	tcaggctggg	gggggagctg	gtgagcatcg	ttggggctgt	gatcactctg	1200
ctccatagaga	ttccagcat	cttcagggtt	gggtccctct	gctatttttg	aaagacgatt	1260
cttggggggc	cattccatgt	catcatgac	acctatgcct	ccctgggtgt	ggtgaccatg	1320
gtgatgcggc	tcaccaacac	caatggggag	gtgggtgccc	tgctccttgc	cctgggtgctg	1380

ggctgggtgca	gtgtcatgta	tttcaactcga	ggattccaga	tgtctgggtcc	cttcaccatc	1440
atgatccaga	agatgatttt	tggagaccta	atgcgtttct	gctggctgat	ggctgtggtc	1500
atcttgggat	ttgcctccgc	gttctatata	attttccaga	cagaggaccc	aaccagtctg	1560
gggcaattct	atgactaccc	catggcactg	ttcaccacct	ttgagctttt	tctcactgtt	1620
attgatgcac	ctgccaacta	cgacgtggac	ttgcccttca	tgttcagcat	tgtcaacttc	1680
gccttcgcca	tcattgccac	actgctcatg	ctcaacttgt	tcacgcgcat	gatgggcgac	1740
accactgga	gggtggccca	ggagagggat	gagctctgga	gggcccagg	cgtggccacc	1800
acagtgatgc	tggagcgga	gctgcctcgc	tgcctgtggc	ctcgctccgg	gatctgtggg	1860
tgcgaattcg	ggctggggga	ccgctgggtc	ctgcgggttg	agaaccacaa	tgatcagaat	1920
cctctgcgag	tgcttcgcta	tgtggaagtg	ttcaagaact	cagacaagga	ggatgaccag	1980
gagcatccat	ctgagaaaca	gccctctggg	gctgagagtg	ggactctagc	cagagcctct	2040
ttggctcttc	caacttcctc	cctgtcccg	accgcgtccc	agagcagcag	tcaccgaggc	2100
tgggagatcc	ttcgtcaaaa	caccttgggg	cacttgaatc	ttggactgaa	ccttagtgag	2160
ggggatggag	aggaggtcta	ccatttttga	ttaac			2195

<210> 55
 <211> 2055
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1471717CB1

<400> 55						
cggctctggg	ccctcagcct	ggctcatgca	caactgtctg	aagtgtctctg	gactatgggtg	60
atgaacagcg	gccttcagac	gcgaggctgg	ggaggaatcg	tgggggtttt	tattattttt	120
gccgtatttg	ctgtcctgac	agtagccatc	cttctgatca	tggagggcct	ctctgtcttc	180
ctgcacgccc	tgcgactgca	ctgggaagct	gtttggggaa	cttgagctat	ttagaagatg	240
gcaaccaagc	caacagagcc	tgtcacgata	ctcagccttc	ggaaattgag	cctggggacc	300
gcagagccac	aggttaaaga	gccaaagacg	ttcacctggg	aagatgcagt	ggagactatc	360
ggcttcgggc	gtttccacat	tgcctctctt	ctgatcatgg	gcagtactgg	ggtggttgag	420
gccatggaga	tcattgtgat	agctgtttgt	tctcctgtca	tccgctgtga	atggcaactg	480
gagaattggc	agggtggcatt	agtaaccacg	atggtgtttt	ttggctacat	ggttttcagt	540
atcctctttg	gectcctggc	tgacagatat	ggccgctgga	agattctgct	catctcgttc	600
ctgtggggag	cctattttct	cttgcctgac	tgcgtttgct	cttcgtacat	ctgggttgte	660
ttcctgcgga	cgatggtggg	ctgtggtgtg	tccggccact	cgcaagggtt	aatcataaag	720
actgaatttt	tgccacgaa	ataccgagge	tatatgttac	ccttgtctca	ggtgttctgg	780
cctgcgggct	ccctgctcat	cattggcttg	gcctctgtga	tcacccccc	catcggtggg	840
cgtgggtcct	ttcgcgtcgc	ctccatcccg	ggcatcatcc	tcacgtggc	cttcaagttt	900
attcctgaat	ctgcgcggtt	caatgtctcc	actgggaaca	ctcgggctgc	cctggccact	960
ctggagcgcg	ttgccaagat	gaaccgctcg	gtcatgcggg	aggggaagct	ggtggagccc	1020
gtcctggaaa	aaagaggaag	at ttgcagac	ctattggatg	ctaaatat	acggaccaca	1080
ttacagatct	gggtcatatg	gcttggaatc	tcttttgcct	actatggggt	tatcctggcc	1140
agtgtcgagc	tgttgagcgc	ggacttggtc	tgtggttcaa	agtcagactc	tgcgggtggtg	1200
gtgactgggg	gggactcagg	ggagagccag	agccctgct	actgccacat	gtttgcaccc	1260
tctgactatc	ggaccatgat	catcagcacc	atcggtgaaa	ttgctttgaa	tccttttaaat	1320
atactgggca	tcaattttct	gggaagacgg	ctgagccttt	ctattacccat	gggatgcacg	1380
gctttattct	gccttctcct	caacatttgc	acttcaagtg	cgggctgat	tggcttctct	1440
ttcatgctga	gggctctggg	agctgcaaac	ttcaacacgc	tctacattta	cacagctgag	1500
gtctacccca	ccacgatgcg	cgttttgggg	atgggaacca	gcggtctcct	gtgtcgcatt	1560
ggtgcaatgg	tggcgccatt	tatatcccag	gttcttatga	gtgcatcaat	actggggggc	1620
ctgtgtctct	tctcatctgt	ctgtgtttga	tgcgccattt	ctgcattcac	tctcccccac	1680
gaaaccaaag	gacgggccc	ccagcaaatt	aaatgaagac	ctgcaaagct	atgtctacca	1740
gatgagaaaa	atgaattcta	tcttcagaac	tgcggtgcat	tttttttaaaa	ccttggtttta	1800
cttctgtatg	ctactcggta	attagtaaa	tgattttttt	ttaaaaggca	tatatgggaa	1860
tggggtaggt	aactgtatat	tgatctcttc	ccttgaggaa	aatatataaa	gtactttttat	1920
aaaatataat	ttaagctttc	aaaggggtgt	gagagggaga	tgggtggggg	gaagatggct	1980
tttcttcgtt	gaaatcaagt	ctgtaaacct	ttatatgaat	aaataactaa	tttttaaactt	2040
acaaaaaaaa	aaaaa					2055

<210> 56
 <211> 4727
 <212> DNA
 <213> Homo sapiens

<220>

<221> misc_feature
 <223> Incyte ID No: 3874406CB1

<400> 56

```

aagagctgct ggagtaggca cccattttaa gaaaaaatga agaagcagca ataaagaagt 60
tghtaatcgtt acctagacaa acagagaact ggttttgaca gtgttttctag agtgcttttt 120
attatttttcc tgacagttgt gttccaccat gttacttttc tcttcagcg aataggctaa 180
atgaatatga aacagaaaag cgtgtatcag caaaccaaag cacttctgtg caagaatttt 240
cttaagaaat ggaggatgaa aagagagagc ttatttggaaat ggggcctctc aatacttcta 300
ggactgtgta ttgctctgtt ttccagttcc atgagaaatg tccagtttcc tggaaatggc 360
cctcagaatc tgggaagggg agataaatat aatagctctt ctttaattggg tgtgtataca 420
ccaatatcta atttaaccca gcagataatg aataaaacag cacttgctcc tcttttgaaa 480
ggaacaagtg tcattggggc accaaataaa acacacatgg acgaaatact tctggaaaat 540
ttaccatatg ctatgggaat catctttaat gaaactttct cttataagtt aatatttttc 600
cagggatata acagtccact ttggaaagaa gatttctcag ctcatgtctg ggatggata 660
ggtgagtttt catgtacatt gaccaaatac tggaaatagag gatttgtggc tttaacaaca 720
gctatttaata ctgccattat agaagtagct ttggtgttcc tgatgagtgt gctgttaaag 780
aaagctgtcc tcaccaattt gggtgtgttt ctccttacc ctttttgggg atgtctggga 840
ttcactgtat tttatgaaca acttccttca tctctggagt ggattttgaa tatttgtagc 900
ccttttgcct ttactactgg aatgattcag attatcaaac tggattataa cttgaatggg 960
gtaatttttt ctgacctttc aggagactca tatacaatga tagcaacttt ttctatgttg 1020
cttttggatg gtctcatcta cttgctattg gcattatact ttgacaaaat tttaccctat 1080
ggagatgagc gccattattc tcttttattt ttcttgaatt catcatcttg tttccaacac 1140
caaaggacta atgctaaggt tattgagaaa gaaatcgatg ctgagcatcc ctctgatgat 1200
tattttgaac cagtgcctcc tgaattccaa ggaaaagaag ccatcagaat cagaaatggt 1260
aagaaggaa ataaaggaaa atctggaaaa aggtgaagcat tgaaaggcct gctctttgac 1320
ataatgaag gtcaaatcac ggcaatcctg ggtcacagtg gagctggcaa atcttcactg 1380
ctaaatatc ttaatggatt gtctgttcca acagaaggat cagttaccat ctataataaa 1440
aatctctctg aaatgcaaga cttggaggaa atcagaaaga taactggcgt ctgtcctcaa 1500
ttcaatgttc aatttgacat actcacctg aaggaaaacc tcagcctgtt tgctaaaata 1560
aaagggattc atctaaagga agtggaaaca gaggtacaac gaatatattt ggaattggac 1620
atgcaaaaaca ttcaagataa ccttgctaaa catttaagt aaggacagaa aagaaagctg 1680
acttttggga ttaccatttt aggagatcct caaattttgc tcttagatga accaactact 1740
ggattggatc ccttttccag agatcaagtg tggagcctcc tgagagagcg tagagcagat 1800
catgtgatcc ttttcagtac ccagtcctat gatgaggtcg acatcctggc tgatagaaaa 1860
gtgatcatgt ccaatgggag actgaagtgt caggttctct ctatcttttt gaaaagaagg 1920
tggggctctg gatatacct aagtttacat aggaatgaaa tatgtaacct agaacaaata 1980
acatccttca ttactcatca catccccgat gctaaattaa aaacagaaaa caaagaaaag 2040
cttgtatata ctttgccact ggaaaggaca aatacatttc cagatctttt cagtgatctg 2100
gataagtgtt ctgaccagg agtgacaggt ccatgataat ccatgtcaac tctaaatgat 2160
gtctttatga aactggagg caactgaact atcgacaag atttcgaaca agtggagatg 2220
ataagagact cagaaagcct caatgaaatg gagctggctc actcttctct ctctgaaatg 2280
cagacagctg tgagtgcacat gggcctctgg agaatgcaag tctttgccat ggcacggctc 2340
cgtttcttaa agttaaaacg tcaaaactaa gtgttattga ccctattatt ggtatttggg 2400
atcgcaatat tccctttgat tgttgaaaa ctatgttata ctatgttata tgaaaagatg 2460
gattgggaat ttaaaaaacga attgtatttt ctctctctct gacaacttcc ccaggaacct 2520
cgtaccagcc tgttgatcat caataacaca gaatcaaata ttgaagattt tataaaatca 2580
ctgaagcatc aaaatatact tttggaagta gatgactttg aaaacagaaa tgggtactgat 2640
ggcctctcat acaatggagc tatcatagtt tctggtaaac aaaaggatta tagattttca 2700
gttgtgtgta ataccaagag attgcactgt tttccaattc ttatgaatat tatcagcaat 2760
gggctacttc aaatgtttta tcacacacaa catattcgaa ttgagtcaag cccattttct 2820
cttagccaca taggactctg gactgggttg ccggatgggt cctttttctt atttttgggt 2880
ctatgtagca tttctcctta tatcaccatg ggcagcatca gtgattacaa gaaaatgct 2940
aagtcaccagc tatggatttc aggctctcat acttctgctt actggtgtgg gcaggcacta 3000
gtggacgtca gcttctctat ttaattctc cttttaatgt atttaatttt ctacatagaa 3060
aacatgcagt accttcttat tacaagccaa attgtgtttg ctttggttat agttactcct 3120
ggttatgcag cttctcttgt cttcttcata tatatgatat catttatttt tcgcaaaagg 3180
agaaaaaaca gtggcctttg gtcattttac ttcttttttg cctccaccat catgttttcc 3240
atcactttaa tcaatcattt tgacctaatg atattgatta ccaccatggg attggttctc 3300
tcataacctt tgcctggatt taacactttt ttggaagtga gagaccagga gcaactacaga 3360
gaatttccag aggcacaaatt tgaattgagt gccactgatt ttctagtctg cttcataccc 3420
tactttcaga ctttgctatt cgtttttgtt ctaagatgca tggaaactaaa atgtggaaag 3480
aaaagaatgc gaaaagatcc tgttttcaga atttcccccc aaagtagaga tgctaagcca 3540
aatccagaag aacccataga tgaagatgaa gatattcaaa cagaaagaat aagaacagca 3600
actgctctga ccacttcaat cttagatgag aaacctgtta taattgccag cgtgctacac 3660
aaagaatatg caggccagaa gaaaagttgc ttttcaaaga ggaagaagaa aatagcagca 3720
agaaatatct ctttctgtgt tcaagaaggt gagatttttg gattgctagg acccagtggt 3780

```

gctggaaaaa	gttcatctat	tagaatgata	tctgggatca	caaagccaac	tgctggagag	3840
gtggaactga	aaggctgcag	ttcagttttg	ggccacctgg	ggtactgccc	tcaagagaac	3900
gtgctgtggc	ccatgctgac	gttgaggga	cacctggagg	tgatgctgc	cgtcaagggg	3960
ctcaggggaag	cggacgcgag	gctcgccatc	gcaagattag	tgagtgcctt	caaactgcat	4020
gagcagctga	atgttctctg	gcagaaatta	acagcaggaa	tcacgagaaa	gttgtgtttt	4080
gtgctgagcc	tcctgggaaa	ctcacctgtc	ttgctcctgg	atgaaccatc	tacgggcata	4140
gacccacacag	ggcagcagca	aatgtggcag	gcaatccagg	cagtcgttaa	aaacacagag	4200
agaggtgtcc	tcctgaccac	ccataacctg	gctgaggcgg	aagccttgtg	tgaccgtgtg	4260
gccatcatgg	tgtctggaag	gcttagatgc	attggctcca	tccaacacct	gaaaaacaaa	4320
cttggcaagg	attacattct	agagctaaaa	gtgaaggaaa	cgtctcaagt	gactttggtc	4380
cacactgaga	ttctgaagct	tttccacacg	gctgcagggc	agcaaaggta	ttcctctttg	4440
ttaacctata	agctgcccgt	ggcagacgtt	tacctctat	cacagacctt	tcacaaatta	4500
gaagcagtga	agcataactt	taacctggaa	gaatacagcc	tttctcagtg	cacactggag	4560
aaggatttct	tagagctttc	taaagaacag	gaagtaggaa	attttgatga	agaaattgat	4620
acaacaatga	gatggaaact	cctccctcat	tcagatgaac	cttaaacctt	caaacctagt	4680
aattttcttg	cttgatctcc	tataaactta	tgttttatgt	aataatt		4727

<210> 57

<211> 3852

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4599654CB1

<400> 57

cgccggcgat	tccgagccta	cgacgcctcc	gctagagccc	gcggggctgc	gccgactcct	60
gctctggagg	ggttgcgggt	acctgatggc	cacagagggc	tctaggaggc	cgagcgtgta	120
agcgggggtg	gcgccatgga	ggcagagcag	cggccggcgg	cgggggccag	cgaaggggcg	180
acccctggac	tggaggcggt	gcctcccgtt	gctccccgcg	ctgcgaccgc	ggcctcagg	240
ccgatcccca	aatctggggc	tgagcctaag	aggaggcacc	ttgggacgct	gctccagcct	300
acggtcaaca	agttctccct	tcgggtgttc	ggcagccaca	aagcagtggg	aatcgagcag	360
gagcgggtga	agtcagcggg	ggcctggatc	atccacccct	acagcgactt	ccggttttac	420
tgggacctga	tcattgctgt	gctgatgggt	gggaacctca	tcgtcctgcc	tgtgggcatc	480
accttcttca	aggaggagaa	ctccccgctt	tggatcgtct	tcaacgtatt	gtctgatact	540
ttcttctctac	tggatctggg	gctcaacttc	cgaacgggca	tcgtgggtgga	ggagggtgct	600
gagatcctgc	tggcaccgcg	ggccatccgc	acgcgctacc	tgcgcacctg	gttccctgggt	660
gacctcatct	cttctatccc	tgtggattac	atcttccctag	tgggtggagct	ggagccacgg	720
ttggacgtcg	aggtctacaa	aacggcacgg	ggcctacgca	tcgttcgctt	caccaagatc	780
ctaagcctgc	tgaggctgct	ccgcctctcc	cgccctatcc	gctacataca	ccagtgggag	840
gagatcttcc	acatgacctt	tgacctggcc	agtgtctgtg	ttcgcatctt	caacctcatt	900
gggatgatgc	tgtctgctatg	tcactgggat	ggctgtctgc	agttccctgg	gcccatgctg	960
caggacttcc	ctcccgcactg	ctgggtctcc	atcaaccaca	tgggtgaacca	ctcgtggggc	1020
cgccagttat	cccatgccct	gttcaaggcc	atgagccaca	tgtgtgtgcat	tggctatggg	1080
cagcaggcac	ctgtaggcat	gcccgaagtc	tggctcacca	tgtctagcat	gatcgtaggt	1140
gccacatgct	acgccatggt	catcgcccat	gccacggcac	tcattccagtc	cctggactct	1200
tcccggcgtc	agtaccagga	gaagtacaag	caggtggagc	agtaacatgtc	cttcacaaag	1260
ctgccagcag	acacgcggca	gcgcattccac	gagtactatg	agcacccgta	ccagggcaag	1320
atgttcgatg	aggaaagcat	cctgggcgag	ctgagcgagc	cgcttcgcga	ggagatcatt	1380
aacttcacct	gtcggggcct	ggtggccccc	atgccgctgt	ttgcccctgc	cgaccccgag	1440
ttcgtcactg	cagttctcac	caagctgcgc	tttgaggctt	tcagccgggg	ggatctcgtg	1500
gtgcgtgagg	gctccgtggg	gaggaagatg	tacttcatcc	agcatgggct	gctcagtgtg	1560
ctggcccgcg	gcgcccggga	cacacgcctc	accgatggat	cctacttttg	ggagatctgc	1620
ctgctaacta	ggggccggcg	cacagccagt	gttcgggctg	acacctactg	cgccctttac	1680
tcactcagcg	tggaccattt	caatgctgtg	cttgaggagt	tccccatgat	gcgcccgggc	1740
tttgagactg	tggccatgga	tcggctgctc	cgcacgggca	agaagaattc	catactgcag	1800
cggaagcgct	ccgagccaag	tccaggcagc	agtgggtggc	tcattggagca	gcatttggtg	1860
caacatgaca	gagacatggc	tcgggggtgt	cggggctcgg	ccccgagcac	aggagctcag	1920
cttagtgga	agggagctact	gtgggagcca	ctggtacatg	cgcccttcca	gcagctgct	1980
gtgacctcca	atgtggccat	tgccttgact	catcagcggg	gcccctctgc	cctctcccct	2040
gactctccag	ccaccctcct	tgtctgctct	gcttggcgct	cagcaggctc	tcagcttcc	2100
ccgctgggtg	ccgtccgagc	tggcccattg	gcattccacct	cccgcctgcc	cgccccacct	2160
gcccgaacct	tgcagccag	cctatcccgg	gcagggcgct	cccaggtctc	cctgctgggt	2220
ccccctccag	gaggaggtgg	acggcggtga	ggacctcggg	gccgcccact	ctcagctcc	2280
caacctcttc	tgcctcagcg	ggcaacaggc	gatggctctc	ctgggctgta	gggatcagga	2340
agtgagcggc	tgcctccctc	agggctcctg	gccaaacctc	caaggacagc	ccagccccc	2400

aggccaccag	tgcctgagcc	agccacaccc	cggggtctcc	agctttctgc	caacatgtaa	2460
aacctttgag	tacatccagc	cttagttctt	gggggtgcagt	agtatgtacc	caagggcaga	2520
tgcctcttgg	ggaaggccat	ggggacctga	aacattgccc	catggaaatg	tcgaccctgt	2580
gcggacattc	cgcatactgc	catgaagacg	gtctctgtgt	cctcagctca	agaatcctgt	2640
agcttgtccc	atcataatcc	attcaccocgt	tcatcatgtg	tactgagcag	ctaccatgtt	2700
caaggtaata	tgccaggcgc	tgtatgtctc	cactgccaaag	tagaagtgc	tcaaaacccct	2760
ctgacaaggga	tattcccttg	gctatgggtcc	tgccagggtgc	aggcccaggc	ccatgacccc	2820
accttttacta	agcacaagta	cttgccactg	ccatcactgc	caagtaacta	gatgtctctg	2880
tttccctgcc	aatgatectg	caggttctgc	ccggctctggt	tatcttcctg	tttccctgtag	2940
catagccagg	cactgccagt	cacctgtgcc	cccattgctg	tcagcagatg	tcttgggtcc	3000
tgagtgtggg	tatccacttt	tacccgctca	ctgccacctg	tggacactct	gtgtctaccc	3060
tctgagtggg	aacatacttc	taagtccct	gcagtctctg	tccgtgtgga	gaccatcttt	3120
ttgtaaactg	cgagcttccct	cttccctgta	ccctctgccc	cagtcgtgac	cccctaaaag	3180
ttaaggggta	gttggcacct	ccttattaat	atgccagcct	agatccccc	cgggtggagg	3240
gcaaattggct	gaatccttgt	gtgatatttt	tttcttcgct	tgtttattta	ttcatttatt	3300
taattgtatt	tattcattta	ctaactttat	gtgttaccac	ttaattttgt	ttacccattc	3360
ctttatccat	ccctcccttc	cttttcagg	aaggagacag	gaggagttag	aggaggcagg	3420
gcctctccat	gccagcctct	gtggctcctg	cccaaaccac	tcagcgcaat	acttgaacct	3480
tctcccaggt	aggggcagga	ggagccacat	gagagaggga	gaaggaccgc	gtttaccttt	3540
agagttttgt	tttgtttttt	ccttctgagt	ctgctgttgg	tgcaggaata	agggaaagge	3600
ccaaggtatc	taagcctggg	gaagggcagg	ccagccagca	cctctgcctt	ctcagggaca	3660
agagttagtcc	tttaccaccc	tcactctgcc	tgccccctct	cctactctac	agcattaaag	3720
actgtggggac	caggacccta	agtctccttt	ccttctgggt	ggggagtctt	aggggttctt	3780
ggtgtgtggg	agaagtttta	taattgcttc	caaacagctg	ggttttaata	taaaatagac	3840
acactcaaaa	aa					3852

<210> 58

<211> 1917

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5047435CB1

<400> 58

atggcagaag	gtgaaagggg	agcagacgtg	ccacatggcc	tccggggcctg	gctggccgac	60
gtggcggttg	cggcgcgtgc	cgcgggagg	cagggcagga	gggacagagg	cgggggcggy	120
ccggaaagtt	tgtccggcgg	cagcggcggt	ggggactccg	gcgggggatg	cgcgcccggc	180
ccctcagcgc	ccccagcacg	ccgcagagtc	ccgctcgcca	tgggccactc	cccacctgtc	240
ctgcctttgt	gtgcctctgt	gtctttgtct	gggtggcctga	cctttgggta	tgaactggca	300
gtcatatcag	gtgcctctgt	gccactgcag	cttgactttg	ggctaagctg	cttggagcag	360
gagttccttg	tgggcagcct	gctcctgggg	gctctcctcg	cctccctggt	tgggtggcttc	420
ctcattgact	gctatggcag	gaagcaagcc	atcctcggga	gcaacttggt	gctgctggca	480
ggcagcctga	ccctgggcct	ggcctgggtcc	ctggcctggc	tggctcctggg	ccgcgctgtg	540
gttggcttgc	ccatttccct	ctcctccatg	gcttgcctga	tctacgtgtc	agagctgggtg	600
gggccacggc	agcggggagt	gctgggtgtc	ctctatgagg	caggcatcac	cgtgggcac	660
ctgctctcct	atgcctcaaa	ctatgcaact	gctggtaacc	cctggggatg	gaggcacatg	720
ttcggctggg	ccactgcacc	tgtgtcctct	caatccctca	gcctcctctt	cctccctgct	780
ggtacagatg	agactgcaac	acacaaggac	ctcatccac	tccagggagg	tgaggccccc	840
aagctgggac	cggggaggcc	acgggtactc	tttctggacc	tcttcagggc	acgcgataac	900
atgcgaggcc	ggaccacagt	gggcctgggg	ctgggtgctc	tccagcaact	aacagggcag	960
cccaacgtgc	tgtgctatgc	ctccaccatc	ttcagctccg	ttgggtttcca	tgggggatcc	1020
tcagccgtgc	tggcctctgt	ggggcttggc	gcagtgaagg	tggcagctac	cctgaccgcc	1080
atggggctgg	tggaccgtgc	aggccgcagg	gctctgttgc	tagctggctg	tgccctcatg	1140
gccctgtccg	tcagtggcat	aggcctcgtc	agctttgccc	tgcccatgga	ctcaggccca	1200
agctgtcttg	ctgtgcccaa	tgccaccggg	cagacaggcc	tccctggaga	ctctggcctg	1260
ctgcaggact	cctctctacc	tcccattcca	aggaccaatg	aggaccaaag	ggagccaatc	1320
ttgtccactg	ctaagaaaac	caagcccatc	cccagatctg	gagacccttc	agccctcctc	1380
cggctggccc	tgaactctgc	cctccctggg	ccccctctgc	ccgctcgggg	gcatgcactg	1440
ctgcgttgga	ccgcactgct	gtgcctgatg	gtctttgtca	gtgccttctc	ccttgggttt	1500
gggccagtga	cctggcttgt	cctcagcgag	atctaccctg	tggagatacg	aggaagagcc	1560
ttcgccttct	gcaacagctt	caactgggag	gccaacctct	tcatcagcct	ctccttccct	1620
gatctcattg	gcaccatcgg	cttgtccttg	accttccctg	tctacggact	gaccgtgtgc	1680
ctcggccctg	gcttcatcta	tttatttgtt	cctgaaacaa	aaggccagtc	gttggcagag	1740
atagaccagc	agttccagaa	gagacgggtc	accctgagct	ttggccacag	gcagaactcc	1800
actggcatcc	cgtacagccg	catcgagatc	tctgcggcct	cctgagggaat	ccgtctgcct	1860

ggaaattctg gaactgtggc tttggcagac catctccage atcctgcttc ctaggcc 1917

<210> 59
 <211> 6791
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7475603CB1

<400> 59
 cgcgctccct gcctgctgct gggcggaggg aaggcggcaa gagctgcgga gcccctggaa 60
 gagcttccag gaaccctgcg ctgtgggata aaggaatgag gttcagaaag gggcaggagt 120
 tgcccgcagc cgcaccgcac gtcttcagcc cgaccgttgt cctgacctct ctgtcccgtc 180
 ccttgcaccg tctcaccatg gccttctgga cacagctgat gctgctgctc tggaagaatt 240
 tcatgtatcg ccggagacag ccggtccagc tcctggctga attgctgtgg cctctcttcc 300
 tcttcttcat cctgggtggct gttcgccact ccaccgcgc cctggagcac catgaatgcc 360
 acttcccaaa caagccactg ccacgcggcg gcaccgtgcc ctggctccag ggtctcatct 420
 gtaatgtgaa caacacctgc tttccgcagc tgacaccggg cgaggagccc gggcgccctga 480
 gcaacttcaa cgcctccctg gtctcccgcg tgcctagcga tgcccgcact gtgctgggag 540
 gggccagtg cccacaggac ctggctggcc tagggaagct gatcgccacg ctgagggctg 600
 cacgcagcac ggcccgacct caaccaacca agcagtctcc actggaacca cccatgctgg 660
 atgtcgcgga gctgctgacg tcaactgctgc gcacggaaat cctgggggttgc gactggggcc 720
 aagcccagga gcccttgcaac agcttgttgg aggcgctga ggacctggcc caggagctcc 780
 ttgcgctgcg cagcctgggtg gagcttgcga cactgctgca gagaccocga gggaccagcg 840
 gccccttggg gttgctgtca gaggccctct gcagtgtcag gggacctagc agcacagtgg 900
 gcccctccct caactggtac gaggctagtgc acctgatgga gctgggtggg caggagccag 960
 aatccgcctt gccagacagc agcctgagcc ccgctgctc ggagctgatt ggagccctgg 1020
 acagccaccc gctgtccgcg ctgctctgga gacgctgaa gcctctgatc ctcggaagc 1080
 tactctttgc accagataca ccttttacc ggaagctcat ggcccagggtg aaccggacct 1140
 tcgaggagct caccctgctg agggatgtcc gggaggtgtg ggagatgctg ggaccccgga 1200
 tcttcacctt catgaacgac agttccaatg tggccatgct gcagcggtc ctgcagatgc 1260
 aggatgaagg aagaaggcag ccagacctg gaggccggga ccacatggag gccctgcgat 1320
 cctttctgga ccctgggagc ggtggctaca gctggcagga cgcacacgct gatgtggggc 1380
 acctggtggg cagctgggc cgagtgcagg agtgcctgtc cttggacaag ctggaggcgg 1440
 caccctcaga ggcagccctg gtgtcgcggg ccttgcaact gctcgcgga catcgattct 1500
 gggcgcggct cgtcttcttg ggacctgagg actcttcaga cccacagag caccacccc 1560
 cagacctggg ccccgccac gtgcgcacaa aaatccgcat ggacattgac gtggtcacga 1620
 ggaccaataa gatcagggac aggttttggg acctggccc agccgcggag cccctgaccg 1680
 acctgocgta cgtgtggggc ggcttctgtt acctgcaaga cctgggtggag cgtgcagccg 1740
 tccgctgct cagcggcgcc aacccccggg ccggcctcta cctgcagcag atgccctatc 1800
 cgtgctatgt ggacgacgtg tctctgctg tctgagccg gtcgctgccc ctcttctctga 1860
 cgctggcctg gatctactcc gtgacactga cagtgaaggc cgtgggtgcg gagaaggaga 1920
 cgcgctgctg ggacacctg cgcgccatgg gctcagccg cgcggtgctc tggctaggct 1980
 ggttctctag ctgcctcggt ccttctctgc tcagcgcgc gctgctggtt ctgggtgctc 2040
 agctggggga catcctcccc tacagccacc cggcgctggt cttctgttct ttggcagcct 2100
 tcgcggtggc cagcgtgacc caggacctcc tgcctagcgc cttcttctcc cgcgccaaac 2160
 tggctgcggc ctggcgccgc cctggcctact tctccctcta cctgccctac gtgctgtgtg 2220
 tggcttggcg ggacggctg ccccggggtg cccgctgggc cgcgagcctg ctgtcgccc 2280
 tggccttcgg ctctggctgc gagagcctgg ctctgctgga ggagcagggc gagggcgcgc 2340
 agtggcaca cgtgggcacc cggcctacgg cagacgtctt cagcctggcc caggtctctg 2400
 gccttctgct gctggacgcg gcgctctacg gcctcgccac ctggtacctg gaagctgtgt 2460
 gccagggcca gtacgggac cctgaacctt ggaattttcc ttttcggagg agctactggt 2520
 gcggacctcg gccccecaag agtccagccc cttgcccac cccgctggac ccaaaggctc 2580
 tggtagaaga ggcaccgccc ggcctgagtc ctggcgtctc cgttcgcagc ctggagaagc 2640
 gcttctctgg aagcccgag ccagccctgc gggggctcag cctggacttc taccagggcc 2700
 acatcacccg ctctctgggc cacaacgggg ccttcacctc gggccacgac gtccgctcca 2760
 gtggcctctt cccaccagt ggtggctctg cactggcggt tctgtctca ctggttgaca 2820
 gcattggccc catcggccc gtctgggtct atgggcggct gaagggtctg agtgccgctg 2880
 tgctgaccgt ggacgagcac cgtctgctgc aggatgtggg gctgggtctc aagcagagt 3000
 tagtggggcc ccacctctct ggtgggatgc aacggaagct gtcgctggcc attgcctttg 3060
 tgggcggctc ccaagtgtgt atcctggac agcctacggc tggcgtggat cctgcttccc 3120
 gccgcggtat ttggcagctg ctgctcaaat accgagaagg tcgcacgctg atcctctcca 3180
 cccaccacct ggatgaggca gagctgctgg gagaccgtgt ggcgctggtg gcaggtggcc 3240
 gcttgtgctg ctgtggatcc ccactcttcc tgcgcgtca cctgggctcc ggctactacc 3300

tgacgctggt	gaaggcccg	ctgcccctga	ccaccaatga	gaaggctgac	actgacatgg	3360
agggcagtg	ggacaccagg	caggaaaaga	agaatggcag	ccagggcagc	agagtcggca	3420
ctcctcagct	gctggccctg	gtacagcact	gggtgcccg	ggcacggctg	gtggaggagc	3480
tgccacacga	gctggtgctg	gtgctgccct	acacgggtgc	ccatgacggc	agcttcgcca	3540
cactcttccg	agagctagac	acgcggctgg	cggagctgag	gctcactggc	tacgggatct	3600
ccgacaccag	cctcgaggag	atcttctctga	aggtggtgga	ggagtgtgct	gcggacacag	3660
atatggagga	tggcagctgc	gggcagcacc	tatgcacagg	cattgctggc	ctagacgtaa	3720
ccctacggct	caagatgccg	ccacaggaga	cagcgtgga	gaacggggaa	ccagctgggt	3780
cagccccaga	gactgaccag	ggctctgggc	cagacgccgt	gggcccggta	cagggtctgg	3840
cactgaccgg	ccagcagctc	caggccctgc	ttctcaagcg	ctttctgctt	gcccgcgca	3900
ggcgccggcg	cctgttcgce	cagatcgtgc	tgcctgccct	ctttgtgggc	ctggccctcg	3960
tgttcagcct	catcgtgctt	cctttcgggc	actaccggcg	tctgcggctc	agtccccacca	4020
tgtacgggtg	tcaggtgtcc	ttcttcagtg	aggacgcccc	aggggacctt	ggacgtgccc	4080
ggctgctcga	ggcgtgctg	caggaggcag	gactggagga	gccccagtg	cagcatagct	4140
cccacagggt	ctcggcacca	gaagtctctg	ctgaagtggc	caaggtcttg	gccagtggca	4200
actggacccc	agagtctcca	tccccagcct	gccagtgtag	ccggcccggt	gcccggcgcc	4260
tgtgcggcga	ctgcccggct	gcagctgggt	gtccccctcc	gccccaggca	gtgaccggct	4320
ctgggggaagt	ggttcagaac	cagacaggcc	ggaacctgtc	tgacttctctg	gtcaagacct	4380
acccgcgcct	ggtgcgccag	ggcctgaaga	ctaagaagtg	ggtgaatgag	gtcagatacg	4440
gaggcttctc	gctggggggc	cgagaccag	gcctgccctc	gggccaagag	ttgggcccgt	4500
cagtggagga	ggtgtggggc	ctgctgagtc	ccctgccttg	cggggccctc	gaccgtgtcc	4560
tgaaaaacct	cacagcctgg	gctcacagcc	tggatgctca	ggacagtctc	aagatctggg	4620
tcaacaacaa	aggctggcac	tccatggtgg	cctttgtcaa	ccgagccagc	aacgcaatcc	4680
tccgtgctca	cctgccccca	ggcccggccc	gccacgcccc	cagcatcacc	acactcaacc	4740
accccttgaa	cctcaccaag	gagcagctgt	ctgaggctgc	actgatggcc	tcctcggtgg	4800
acgtcctcgt	ctccatctgt	gtggtctttg	ccatgtcctt	tgtcccggcc	agcttcaact	4860
ttgtcctcgt	tgaggagcga	gtcacccgag	ccaagcacct	gcagctcatg	gggggcctgt	4920
ccccaccctt	ctactggctt	ggcaactttc	tctgggacat	gtgtaactac	ttggtgccag	4980
catgcacgtg	ggtgctcate	tttctggcct	tccagcagag	ggcatatgtg	gcccctgcca	5040
acctgcctgc	tctcctgctg	ttgctactac	tgtatggctg	gtcgatcaca	ccgctcatgt	5100
acccagcctc	cttcttcttc	tccgtgccca	gcacagccta	tgtggtgctc	acctgcataa	5160
acctctttat	tggcatcaat	ggaagcatgg	ccacctttgt	gcttgagctc	ttctctgate	5220
agaagctgca	ggaggtgagc	cggatcttga	aacaggctct	ccttatcttc	ccccacttct	5280
gcttggggccg	ggggctcatt	gacatggtgc	ggaaccaggc	catggctgat	gccttttgagc	5340
gcttggggaga	caggcagttc	cagtcacccc	tgcgctggga	ggtggtcggc	aagaacctct	5400
tggccatggg	gatacagggg	ccccctcttc	ttctcttcac	actactgctg	cagcaccgaa	5460
gccaaactcct	gccacagccc	aggggtgaggt	ctctgccact	cctgggagag	gaggacgagg	5520
atgtagcccg	tgaacgggag	cgggtggtcc	aaggagccac	ccagggggat	gtgttggtgc	5580
tgaggaaactt	gaccaaggta	taccgtgggc	agaggatgcc	agctgttgac	cgcttgtgcc	5640
tggggattcc	ccctggtgag	tgtttttggc	tgctgggtgt	gaatggagca	gggaagacgt	5700
ccacgttttcg	catggtgacg	ggggacacat	tggccagcag	ggcgaggct	gtgctggcag	5760
gccacagcgt	ggcccgggaa	cccagtgtctg	cgcacctcag	catgggatac	tgcctcaat	5820
ccgatgccat	ctttgagctg	ctgacgggcc	gcgagcacct	ggagctgctt	gcgcgcctgc	5880
gcggtgtccc	ggaggcccag	ggtgccccaga	ccgctggctc	gggcctggcg	cgtctgggag	5940
tctcatggta	cgcagaccgg	cctgcaggca	cctacagcgg	agggaaacaaa	cgcaagctgg	6000
cgcagggccct	ggcgtggtt	ggggaccag	ccgtggtgtt	tctggacgag	ccgaccacag	6060
gcatggaccc	cagcgcgcgg	cgcttccctt	ggaacagcct	tttggcctg	gtgcgggagg	6120
gccgttcagt	gatgtccacc	tcccatagca	tggaggagtg	tgaagcgctc	tgctcgcgcc	6180
tagccatcat	ggtgaatggg	cggttccgct	gcctgggcag	cccgaacat	ctcaagggca	6240
gattcgcggc	gggtcacaca	ctgacctgc	gggtgcccg	cgcaaggtcc	cagccggcag	6300
cggccttcgt	ggcgccgag	ttccctgggg	cgagctgcg	cgaggcacat	ggaggccgcc	6360
tgcgcttcca	gctgccgcgg	ggagggcgct	gcgcctggc	gcgcgtcttt	ggagagctgg	6420
cgggtgcacgg	cgcagagcac	ggcgtggagg	acttttccgt	gagccagacg	atgctggagg	6480
aggtattctt	gtacttctcc	aaggaccagg	ggaaggacga	ggacaccgaa	gagcagaagg	6540
aggcaggagt	gggagtggac	cccgcgccag	gcctgcagca	ccccaaacgc	gtcagccagt	6600
tctcagatga	ccctagcact	gccgagactg	tgtcttgagc	ctccctcccc	tgcggggccg	6660
cggggaggcc	ctgggaatgg	caagggcaag	gtagagtgcc	taggagccct	ggactcaggc	6720
tggcagaggg	gctggtgccc	tggagaaaaa	aaagagaagg	ctggagagaa	gccgtggtgg	6780
tgaaaaaaaaa	a					6791

<210> 60
 <211> 5214
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID No: 7477845CB1

<400> 60

atgctcaaaa	ggaagcagag	tccaggggtg	gaagcccagc	cagtcactga	ctttggtect	60
gatgagtctc	tgctcggataa	tgctgacatc	ctctggatta	acaaaccatg	ggttcactct	120
ttgctgcgca	tctgtgccat	catcagcgct	atctctgttt	gtatgaatac	gccaatgacc	180
ttcgagcact	atcctccact	tcagtatgtg	accttcactt	tggatacatt	attgatgttt	240
ctctacacgg	cagagatgat	agcaaaaatg	cacatccggg	gcattgtcaa	gggggatagt	300
tcctatgtga	aagatcgctg	gtgtgttttt	gatggattta	tggctctttt	cctttggggt	360
tctttgggtg	tacaggtgtt	tgaaattgct	gatatagttg	atcagatgtc	accttggggc	420
agtttgcgga	ttccagggcc	actgattatg	atccgagcat	tcoggattta	tttccgattt	480
gaactgccaa	ggaccagaat	tacaaatatt	ttaaagcgat	cgggagaaca	aatatggagt	540
gtttccattt	ttctactttt	ctttctactt	ctttatggaa	ttttaggagt	tcagatgttt	600
ggaacatttt	cttatcactg	tgttgtaaat	gacacaaagc	caggggaatg	aacctggaat	660
agtttagcta	ttccagacac	acactgctca	ccagagctag	aagaaggcta	ccagtgcaca	720
ccttgattta	aatgcatgga	ccttgaagat	ctgggactta	gcaggcaaga	gctgggtcac	780
agtggcttta	atgagatagg	aactagtata	ttcaccgctc	atgaggccgc	ctcacaggaa	840
ggctgggtgt	tcctcatgta	cagagcaatt	gacagctttc	cccgttggcg	ttctacttcc	900
tatttcatca	ctctcatttt	cttctctgcc	tggttgtgta	agaacgtgtt	tattgctgtt	960
atcattgaaa	catttgcaga	aatcagagta	cagtttcaac	aaatgtgggg	atcgagaagc	1020
agcagacctc	caacagccac	caccagcatg	tttcatgaag	atgctgctgg	aggttggcag	1080
ctggtagctg	tggatgtcaa	caagccccag	ggacgcgccc	cagcctgcct	ccagaaaatg	1140
atgcggtcat	ccgttttcca	catgttccatc	ctgagcatgg	tgaccgtgga	cgtgatcgtg	1200
gcggctagca	actactacaa	aggagaaaac	ttcaggaggc	agtacgacga	gttctacctg	1260
gcggagggtg	ctttttacagt	actttttgat	ttggaagcac	ttctgaagat	atggtgtttg	1320
ggattttactg	gatataattag	ctcatctctc	cacaaattcg	aactactact	cgtaattgga	1380
actactcttc	atgtataccc	agatctttat	cattcacaat	tcacgtactt	tcagggtactc	1440
cgagtagttc	ggctgattaa	gatttccact	gcattagaag	actttgtgta	caagatattt	1500
ggtcctggaa	aaaagcttgg	gagttttggt	gtattttactg	ccagcctctt	gattgttatg	1560
tcagcaatta	gtttgcagata	gttctgcttt	gtcgaagaac	tggacagatt	tactacgttt	1620
ccgagggcat	ttatgtccat	gttccagatc	ctcaccagg	aaggatgggt	ggacgtaattg	1680
gaccaaactc	taaatgctgt	gggacatatg	tgggcaccgc	tgggttgccat	ctatttcatt	1740
ctctatcatc	tttttgccac	tctgatcctc	ctgagtttgt	ttgttgctgt	tattttggac	1800
aacttagaac	ttgatgaaga	cctaaagaag	cttaaacat	taaagcaaag	tgaagcaaat	1860
gcggacacca	aagaaaagct	ccctttacgc	ctgcgaatct	ttgaaaaatt	tccaaacaga	1920
cctcaaattg	tgaaaatctc	aaagcttcc	tcagatttta	cagttccctaa	aatcagggag	1980
agttttatga	agcagtttat	tgaccgccag	caacaggaca	catgttgcc	tctgagaagc	2040
ctcccgacca	cctcttccctc	ctcctgcgac	cactccaaac	gctcagcaat	tgaggacaac	2100
aaatacatcg	accaaaaact	tgcgaagtct	gttttcagca	tcagggcaag	gaaccttctg	2160
aaaaggagga	ccgcagtcac	taaaatctta	aggccttgca	cccgcagcgc	ctgttctgcg	2220
ggatcatttg	aggggcagcc	cgcaaaggag	aggtaaatcc	tcagcgtgca	gcatacatc	2280
cgccaagagc	gcaggtcact	aagacatgga	tcaaacagcc	agaggatcag	caggggaaaa	2340
tctcttggaa	ctttgactca	agatcattgc	aatacagtga	tatatagaaa	tgctcaaaga	2400
gaagtccagt	aaataaagat	gattcaggaa	aaaaaggagc	tagcagagat	gcttcaagga	2460
aagtgcaaaa	aggaactcag	agagagccac	ccatacttcg	ataagccact	gttctattgtc	2520
gggcgagaac	acaggttcag	aaacttttgc	cgggtggtgg	tcagagcacg	cttcaacgcg	2580
tctaaaacag	accctgtcac	aggagctgtg	aaaaatacaa	agtaccatct	tctttatgat	2640
ttgctgggat	tggtcactta	cctggactgg	gtcatgatca	tcgtaacctc	tgactcttgc	2700
atttccatga	tgtttgagtc	cccgtttcga	agagtcatgc	atgcacctac	tttgagattt	2760
gctgagtatg	tgtttgtgat	attcatgagc	attgagctta	atctgaagat	tatggcagat	2820
ggcttatttt	tcactccaac	tgctgtcatc	agggaacttc	gtggagtaat	ggacataatt	2880
atatactctt	tgagcttgat	atttctttgt	tggatgcctc	aaaatgtacc	tgctgaatcg	2940
ggagctcagc	ttctaattgg	ccttcgggtg	ctgagacctc	tgcgcataat	caaactgggtg	3000
cccagatgga	ggaaagtgtg	tcgagaactt	ttcagcggct	tcaaggaaat	tttttttggtc	3060
tccattcttt	tgctgacatt	aatgctcggt	tttgcaagct	ttggagtcca	gctttttgct	3120
ggaaaactgg	ccaagtgcac	tgatcccaac	attattagaa	gggaagattg	caatggcata	3180
ttcagaatta	atgtcagtg	gtcaaagaac	ttaaatttaa	aattgaggcc	tggagagaaa	3240
aaacctggat	tttgggtgcc	ccgtgttttg	gcgaatcctc	ggaacttta	tttcgacaa	3300
gtgggaaacg	ctatgttcgc	gttgtttgaa	gttctctctc	tgaaaggctg	ggtggaagtg	3360
agagatgtta	ttattcatcg	tgtggggccg	atccattgaa	tctatatatt	tgttttttga	3420
ttcttgggtt	gcattgattg	actgaccctt	tttgtttggg	tagttatttc	taatttcaat	3480
gaaaacaagg	ggacggcttt	gctgaccgtc	gatcagagaa	gatgggaaga	cctgaagagc	3540
cgactgaaga	tcgcacagcc	tcttcatctt	ccgcctcgcc	cggataatga	tggtttttaga	3600
gctaaaatgt	atgacataac	ccagcatcca	ttttttaaga	ggacaatcgc	attactcgtc	3660
ctggcccag	ccgtgttgct	ctctgtcaag	tgggacgtcg	aggaccgggt	gacctacct	3720
ttggcaacaa	tgctcagttg	tttccacctc	atctttgttc	tggaggtaac	catgaagatc	3780
atagcaatgt	cgctgctgg	cttctggcaa	agcagaagaa	accgatacga	tctcctggtg	3840


```

acgtcgcttg gcggttgatg ggtggtgctt cactttgccc tcttgaatgc atatacttac 3900
atgatgggag cttgtgtgat tgtatttagg tttttctcca tctgtggaaa acatgtaacg 3960
ctaaagatgc tctctctgac agtggtcgtc agcatgtaca agagcttctt tatcatagta 4020
ggcatgtttc tcttctgtgct gtgttacgct tttgtctggag ttgttttatt tgggtactgtg 4080
aaatatgggg agaataattaa caggcatgca aatttttctt cggctggaaa agctattacc 4140
gtactgttcc gaattgtcac aggtgaagac ttgaacaaga ctgtatggta 4200
cagcctccgt tttgtactcc agatgaattt acatactggg caacagactg tggaaattat 4260
gctggggcac ttatgtattt ctgttcattt tatgtcatca ttgcctacat catgctaaat 4320
ctgctttagg ccataaattgt ggagaatttc tcttggattt attccactga ggaggaccag 4380
cttttaagtt acaatgatct tcgccacttt caaataatat ggaacatggt ggatgataaa 4440
agagaggtat tccccacgtt ccgctcaag ttctgtctgc ggctactgag tgggaggctg 4500
gaggtggacc tggacaagga caagctcctg tttaagcaca tgtgtacga aatggagagg 4560
ctccacaatg gcgggcagct cacttccat gatgtcctga gcatgcttcc ataccgggtcc 4620
gtggacatcc ggaagagctt gcagctggag gaactcctgg cgagggagca gctggagtac 4680
accatagagg aggaggtggc caagcagacc atccgcagt gtgctcaagaa gtgcctgaag 4740
cgcatcagag ctaaacagca gcagtctgac agtatcatcc acagcctgag agagagtcag 4800
cagcaagagc tgagccggtt tctgaaccgc ccagcatcg agaccacca gccagtgag 4860
gacacgaatg ccaacagtca ggacaacagc atgcaacctg agacaagcag ccagcagcag 4920
ctcctgagcc ccacgtctgc ggtcagagga ggaagtccgc aagatgcagc cgacgcaggg 4980
aaacccagca ggaaatttgg gcagtggcgt ctgccctcag ccccaaaacc aataagccat 5040
tcagtgtcct cagtcacact acggtttgga ggaaggacaa ccatgaaatc tgtcgtgtgc 5100
aaaatgaacc ccatgactga cgcygttcc tgcggttctg aagttaagaa gtggtggacc 5160
cggcagctga ctgtggagag cgacgaaagt ggggatgacc ttctggatat ttag 5214

```

<210> 61

<211> 1818

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 168827CB1

<400> 61

```

ggaaattgct tccgtgacct tgctgcagat gggagagagg gccattaag aagagagtgg 60
ggtcaggatc aacacacaca cttagtgtga ttttaaggaaa ggaaatattt tctctttgaa 120
cttatctgga tacagtcatt ttgtctctc ttggggatca cttgtccagc ctcaatggcc 180
tttcaggacc tcttagatca agttggaggc ctggggagat tccagatcct tcagatgggt 240
ttccttataa tgttcaacgt catagtatac catcaaactc agctggagaa cttcgcagca 300
ttcatacttg atcatcgtg ctgggttcat atactggaca atgacactat cctgacaat 360
gaccttggga cctcagcca ggatgccctc ctgagaatct ccatcccatt cgactcaaat 420
ctgaggccag agaagtgtc tcgctttgtc catcccagc ggaagctcat tcatctgaat 480
gggaccttcc ccaacacagc tgagccagat acagagccct gtgtggatgg ctgggtatat 540
gaccaagctc ccttccctc caccattgtg actaagtggg atctggatg cgaatctcaa 600
ccagtgaatt cagtacata atttctattc atggctggaa tgatgggtgg aggcaacct 660
tatggccatt tgtcagacag gtttgggaga aagtctgtgc tcagatggtc ttacctccag 720
ctcgccattg taggcacctg tgcggccttt gctcccacca tctctgtata ctgtccctg 780
cgcttcttgg ctggggctgc tacatttagc atcattgtaa atactgtttt gttaatgtga 840
gagtggataa ctaccaatt ctgtgccatg gcattgacat tgacactttg tctgtctagt 900
attggacata taacctggg aagcctggct ttgtcattc gagaccagt cactctccag 960
ttggtgatgt ctgcacctg ctttgtcttc tttctgttct caaggtggct ggcagagtct 1020
gtcgggtggc tcattatcaa caacaaacca gaagagggct taaaggaaact tacaaaagct 1080
gcacacagga atggaatgaa gaatgctgaa gacatcctaa ccatggaggt tttgaaatcc 1140
accatgaagc aagaactgga ggcagcacag aaaaagcatt ctctttgtga attgctccgc 1200
ataccaaca tatgtaaaag aatctgtttc ctgtcctttg tgagatttgc aagtaccatc 1260
cctttttggg gccttacttt gcacctccag catctgggaa acaatgtttt cctgttgcag 1320
actctctttg gtgcagtcac cctcctggcc aattgtgttg caccttgggc actgaatcac 1380
atgagccgtc gactaagcca gatgtctctc atgttcttac tggcaacctg ccttctggcc 1440
atcatatttg tgctcaaga aatgcagacc ctgctgtggg ttttggcaac cctgggtgtg 1500
ggagctgctt ctcttggcat tacctgttct actgcccaag aaaatgaact aattccttcc 1560
ataatcaggg gaagagctac tggaatcact ggaaactttg ctaatatgtg gggagccctg 1620
gttccctca tgatgatcct aagcatatat tctcgacccc tgccctggat catctatgga 1680
gtctttgcca tctctcttgg ccttgttgtc ctctctcttc ctgaaaccag gaaccagcct 1740
cttcttgaca gcatcagga tgtggaaaat gaggagtaaa atagcctagc tgccctcag 1800
aggagctctg tgctatat 1818

```

<210> 62

<211> 2245
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7472734CB1

<400> 62
 cccttggaac agtaggatgt tggatgatgca aaagtcaatg tttaaactca acattccact 60
 ttcctttaac taagaatagt ttttattaac ttttagtaaa actcagtcct agtccaaaaa 120
 aagccctgct ctctgatctt tgtacaagaa catcataaag caattcactt tggattttct 180
 aatatcccat ttctgagaag aatggcagac tattgaacag gtgtatttta ggtcacgtgg 240
 ggactgcatc cacctgaaaa tccaccgttg acttatcagg aaactcagag atcaggatct 300
 ttcacagagt agtcttttaa gaagattcag ttgtcaacag ctagcagtcct ctttgccaaa 360
 taattatata tgtgacttct gaaactattt ggctgcctaa agttaaagga cttggggaaa 420
 gtccctccac tgcctctctg cagtagtgct acaccactca gtgcagggcc caccaagaag 480
 aaagcagtggt caggatccac atggcactat ggtaactttg tgaaagggga cattttctcc 540
 ctctgaactt ctcttcataa agtcattgtg ctctctcttg gggatcacct gttcagtcct 600
 aatgggcttt gatgtgctcc tggatcaagt ggggtggcatg gggagattcc agatttgtct 660
 gatagctttc ttttgcatca ccaacatcct actgttccct aatatttgtt tggagaactt 720
 cactgcattc acccctagtc atcgctgctg ggtcccccct ctggacaatg acactgtgtc 780
 tgacaatgat accgggaccc tcagcaagga tgacctcctg agaatctcca tcccactgga 840
 ctcaaacctg aggccacaga agtgctcagcg ctttatccat ccccgagggc agctccttca 900
 cctgaacggg accttcccca acacaaatga gccagacacg gagccctgtg tggatggctg 960
 ggtgtacgac agaagctctt tctctccac catcgtgact gagtgggacc tggatgtgta 1020
 atctcagtc ctaaaatcaa tggttcaatc cctatttatg gctgggtcac ttcctggagg 1080
 tctaataat ggccatcttt cagacagggt tgggagaaag ttcgtgctca gatggcttta 1140
 cctccagctc gccattgtag gcacctgtgc ggcctttgct cccaccatcc tctgtacttg 1200
 ctccctgcgc ttcttggtcg gggctgctac atttagcctc attgtaaata ctgttttgtt 1260
 aattgtagag tggataactc accaattctg tgccatggca ttgacattga cactttgtgc 1320
 tgctagtatt ggacatataa ccctgggaag cctggccttt gtcattcgag accagtgcct 1380
 cctccagttg gtgatgtctg caccatgctt tgtctctctt ctgttctcaa ggtggctggc 1440
 agagtctgct cgggtggctc ttatcaacaa caaaccagaa gagggcttaa aggaacttag 1500
 aaaagctgca cacaggaatg gaatgaagaa tgctgaagac atcctaacca tggaggtttt 1560
 gaaatccacc atgaagcaag aactggaggc agcacagaaa aagcattctc tttgtgaatt 1620
 gctccgcata cccaacatat gtaaaagaat ctgtttctct tcttttgtga gatttgcaag 1680
 taccatccct ttttggggcc ttactttgca cctccagcat ctgggaaaca atgttttctt 1740
 gttgcagact ctctttggtg cagtcaccct cctggccaat tgtgttgca cttgggcact 1800
 gaatcacatg agcgtcgac taagccagat gcttctcatg ttcctactgg caacctgcct 1860
 tctggccatc atatttgtgc ctcaagaaat cgagaccctg cgtgtgggtt tggcaaccct 1920
 ggggtgtggga gctgcttctc ttggcattac ctgttctact gccaagaaa atgaactaat 1980
 tctttccata atcaggggaa gagctactgg aatcactgga aactttgtct atattggggg 2040
 agccctggct tccctcatga tgatcttaag catatattct cgacccctgc cctggatcat 2100
 ctatggagtt tttgcaatcc ttctggcctc tctgtctctc ctccttctct aaaccaggaa 2160
 ccagcctctt cttgacagca tccaggatgt ggaaaatgag ggagtaaata gcctagctgc 2220
 ccctcagagg agctctgtgc tatag 2245

<210> 63
 <211> 3196
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7473473CB1

<400> 63
 gggggggcgc ggggagcgct actaccatga actgcctggt cctcctcccc agagctgctc 60
 atccgggtcg ggctggagac acagtcaggg gaccccgctc ccgcgcgcgc gccccctctt 120
 ctttcggctc aatcttctct tccacctttt cctcctcttc ctccaccttc tttgctgca 180
 tccccccctc ccccgccgcg gatcctggcc gctgctctcc agaccagga tgccgggggg 240
 caagagaggg ctggtggcac cgcagaacac atttttggag aacatcgctc ggcgtctcag 300
 tgaatcaagt ttcttactgg gaaatgcccc gattgtggat tggcctgtag tttatagtaa 360
 tgacgggttt ttgaaactct ctggatatca tctgagctgac gtcattgcaga aaagcagcac 420
 ttgcagtttt atgtatgggg aattgactga caagaagacc attgagaaa tccaggcaaac 480
 ttttgacaac tacgaatcaa actgctttga agttcttctg tacaagaaaa acagaacccc 540

tgttttggttt	tatatgcaaa	ttgcaccaat	aagaaatgaa	catgaaaagg	tggctcttgtt	600
cctgtgtact	ttcaaggata	ttacgttgtt	caaacagcca	atagaggatg	attcaacaaa	660
agggttgacg	aaatttgccc	gattgacacg	ggcttttgaca	aatagccgaa	gtgttttgca	720
gcagctcacg	ccaatgaata	aaacagaggt	ggctcataaa	cattcaagac	tagctgaagt	780
tcttcagctg	ggatcagata	tccttccctca	gtataaacia	gaagcgccaa	agacgccacc	840
acacattatt	ttacattatt	gtgcttttaa	aactacttgg	gattgggtga	ttttaattct	900
tacctctac	accgccatta	tggttcctta	taatgtttcc	ttcaaaacia	agcagaacia	960
catagcctgg	ctgttactgg	atagtgtggt	ggacgttatt	tttctggttg	acatcgtttt	1020
aaattttcac	acgactttcg	tggggcccg	tggagaggtc	atttctgacc	ctaagctcat	1080
aaggatgaac	tatctgaaaa	ccttggtttgt	gatcgatctg	ctgtcttgtt	taccttatga	1140
catcatcaat	gcctttgaaa	atgtggatga	gggaatcagc	agtctcttca	gttcttttaa	1200
agtgggtgct	ctctttacgac	tgggcctgtg	ggctaggaaa	ctggaccatt	acctagaata	1260
tggagcagca	gtcctcgtgc	tcctgggtgtg	tgtgtttgga	ctgggtggccc	actggctggc	1320
ctgcataatg	tatagcatcg	gagactacga	ggtcattgat	gaagtcacta	acaccatcca	1380
aatagacagt	tggctctacc	agctggcttt	gagcattggg	actccatata	gctacaatac	1440
cagtgtctgg	atatgggaag	gaggaccagg	caaggattca	ttgtacgtgt	cctctctcta	1500
cctttaccatg	acaagcctta	caaccatagg	atttggaaac	atagctccta	ccacatagtg	1560
ggagaagatg	ttttcgggtg	ctatgatgat	ggttggcgct	cttcttttatg	caactatttt	1620
tggaaatgtt	acaacaattt	tccagcaa	gtatgccaac	accaaccgat	accatgagat	1680
gctgaataat	gtacgggact	tcctaaaact	ctatcaggct	ccaaaaggcc	ttagtgagcg	1740
agtcattggat	tatatgtct	caacatggct	catgtcaaaa	ggcattgata	cagaaaaggt	1800
cctctccatc	tgtcccaagg	acatgagagc	tgatatctgt	gttcattctaa	accggaaggt	1860
ttttaatgaa	catcctgctt	ttcgattggc	cagcgatggg	tgtctgcgcg	ccttgccggt	1920
agagttccaa	accattcact	gtgctcccg	ggacctcatt	taccatgctg	gagaaagtgt	1980
ggatgccctc	tgctttgttg	tgtcaggatc	ccttggaaag	atccaggatg	atgaggtggt	2040
ggctatttta	gggaagggtg	atgtatttgg	agacatcttc	tgggaaggaaa	ccacccttgc	2100
ccatgcattg	gcgaacgtcc	gggcactgac	gtactgtgac	ctacacatca	tcaagcgga	2160
agccttgctc	aaagtccctg	actttttatac	agcttttgca	aactccttct	caaggaatct	2220
cactcttact	tgcaatctga	ggaaacggat	catctttcgt	aagatcagtg	atgtgaagaa	2280
agaggaggag	gagcgctcc	ggcagaagaa	tgaggtgacc	ctcagcatc	ccgtggacca	2340
cccagtcaga	aagctcttcc	agaagttcaa	gcagcagaag	gagctgcgga	atcagggtct	2400
aacacagggt	gaccctgaga	ggaaccaact	ccaggtagag	agccgctcct	tacagaatgg	2460
agcctccatc	accggaacca	gcgtgggtgac	tgtgtcacag	attactccca	ttcagacgtc	2520
tctggcctat	gtgaaaacca	gtgaatccct	taagcagaac	aaccgtgatg	ccatggaact	2580
caagcccaac	ggcgggtgctg	accaaaaatg	tctcaaagtc	aacagcccaa	taagaatgaa	2640
gaatggaaat	ggaaaagggt	ggctgcgact	caagaataat	atgggagccc	atgaggagaa	2700
aaaggaagac	tgggaataatg	tcactaaagc	tgagtcaatg	gggctattgt	ctgaggaccc	2760
caagagcagt	gattcagaga	acagtgtgac	caaaaaccca	ctaagaaaaa	cagattcttg	2820
tgacagtggg	attacaaaaa	gtgaccttcg	tttggataag	gctggggagg	cccgaagtcc	2880
gctagagcac	agtcctatcc	aggctgatgc	caagcacccc	ttttatccca	tccccgagca	2940
ggcctttacag	accacactgc	aggaagtcaa	acacgaactc	aaagaggaca	tccagctgct	3000
cagctgcaga	atgactgccc	tagaaaagca	ggtggcagaa	attttaaaaa	tactgtcgga	3060
aaaaagcgta	cccaggcct	catctcccaa	atcccaaagt	ccactccaag	taccccccca	3120
gataccatgt	caggatattt	ttagtgtctc	aaggcctgaa	tcacctgaat	ctgacaaaaga	3180
tgaatccac	ttttaa					3196

<210> 64

<211> 1602

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477725CB1

<400> 64

atggcctttg	aggagctctt	gagtcaggtt	ggaggccttg	ggagatttca	gatgcttcat	60
ctgggttttta	ttcttccctc	tctcatgtta	ttaatccctc	atatactgct	agagaacttt	120
gctgcagcca	ttcctgggtca	tctgtgtgtg	gtccacatgc	tggacaataa	tactggatct	180
ggaatgaaa	ctggaatcct	cagtgaagat	gcctcttga	gaatctctat	cccactagac	240
tcaaatctga	ggccagagaa	gtgtcgtcgc	tttgtccatc	cccagtggca	gcttcttcac	300
ctgaatggga	ctatccacag	cacaagtgcg	gcagacacag	aacctgtgt	ggatggctgg	360
gtatatgatc	aaagctactt	cccttcgacc	attgtgacta	agtgggacct	ggtatgtgat	420
tatcagtcac	tgaatcagtg	ggttcaatc	ctacttctga	ctggaatgct	ggtgggaggg	480
atcatagggtg	gccatgtctc	agacaggttt	ggcggaagat	ttattctcag	atgggtgttg	540
ctccagcttg	ccattactga	cacctgcgct	gccttcgctc	ccaccttccc	tgtttactgt	600
gtactacgct	tcttggcagg	tttttcttcc	atgatcatta	tatcaaataa	ttctttgccc	660

attactgagt	ggataaggcc	caactctaaa	gccctggtag	taatattgtc	atctggtgcc	720
cttagtattg	gacagataat	cctgggaggc	ttggcttatg	tcttcgaga	ctggcaaacc	780
ctgcacgtgg	tggcgtctgt	acctttcttt	gtcttctttc	ttctttcaag	gtggctggtg	840
gaatctgctc	ggtggttgat	aatcaccaat	aaactagatg	agggcttaaa	ggcacttaga	900
aaagttgcac	gcacaaatgg	aataaagaat	gctgaagaaa	ccctgaacat	agaggttgta	960
agatccacca	tgcaggagga	gctggatgca	gcacagacca	aaactactgt	gtgtgacttg	1020
ttccgcaacc	ccagtatgcg	taaaaggatc	tgtatcctgg	tatttttgag	atgtgcaaac	1080
acaatacctt	tttatggtac	catggtcaat	cttcagcatg	tggggagcaa	cattttctctg	1140
ttgcaggtag	tttatggagc	tgtcgtctct	atagttcgat	gtcttgctct	tttgacacta	1200
aatcatatgg	gccgtcgaat	aagccagata	ttgttcatgt	tcctggtggg	cctttccatt	1260
ttggccaaca	cgtttgtgcc	caaagaaatg	cagaccctgc	gtgtggcctt	ggcatgtctg	1320
ggaatcggtt	gttctgctgc	tactttttcc	agtgttgctg	ttcacttcac	tgaactcatc	1380
cccactgttc	tcagggaag	agcttcagga	atagatttaa	cggctagtag	gattggagca	1440
gcactggctc	ccctcttgat	gaccttaacg	gtatttttta	ccactttgcc	atggatcatt	1500
tatggaatct	ccccatcat	tggtggcctt	attgtcttcc	tcctaccaga	aaccaagaat	1560
ctgcctttgc	ctgacaccat	caaggatgtg	gaaaatcagt	ga		1602

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/004520 A3

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/47, 16/18, C12Q 1/68, G01N 33/68

(21) International Application Number: PCT/US01/21448

(22) International Filing Date: 5 July 2001 (05.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/216,547	7 July 2000 (07.07.2000)	US
60/218,232	14 July 2000 (14.07.2000)	US
60/220,112	21 July 2000 (21.07.2000)	US
60/221,839	28 July 2000 (28.07.2000)	US

(71) Applicant (*for all designated States except US*): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **RAUMANN, Brigitte, E.** [US/US]; 5801 South Dorchester Avenue #3B, Chicago, IL 60637 (US). **THORNTON, Michael** [US/US]; 9 Medway Road, Woodside, CA 94062 (US). **DING, Li** [CN/US]; 3353 Alma Street #146, Palo Alto, CA 94306 (US). **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **HARLAND, Lee** [GB/GB]; 18 Chaucer Court, New Dover Road, Canterbury, Kent CT1 3AU (GB). **BURFORD, Neil** [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **GREENE, Barrie, D.** [US/US]; 1332 10th Avenue #104, San Francisco, CA 94122 (US). **SANJANWALA, Madhu, S.** [US/US]; 210 Sylvia Court, Los Altos, CA 94024 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **YAO, Monique, G.** [US/US]; 111 Frederick Court, Mountain View, CA 94043 (US). **YANG, Junming** [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). **PATTERSON, Chandra** [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). **GANDHI, Ameena, R.** [US/US]; 837 Roble Avenue #1, Menlo Park, CA 94025 (US). **HAFALIA, April, J., A.** [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). **TRIBOULEY, Catherine, M.** [FR/US]; 1121 Tennessee Street #5, San Francisco, CA 94107 (US). **WALIA,**

Narinder, K. [US/US]; 890 Davis Street #205, San Leandro, CA 94577 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). **WALSH, Roderick, T.** [IE/GB]; 8 Boundary Court, St. Lawrence Road, Canterbury, Kent CT1 3EZ (GB). **RAMKUMAR, Jayalaxmi** [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). **LU, Yan** [CN/US]; 3885 Corrina Way, Palo Alto, CA 94303 (US). **LU, Dyung, Aina, M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **AZIMZAI, Valda** [US/US]; 5518 Boulder Canyon Drive, Castro Valley, CA 94552 (US). **LAL, Preeti** [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). **ELLIOTT, Vicki, S.** [US/US]; 3770 Polton Place Way, San Jose, CA 95121 (US). **NGUYEN, Danniell, B.** [US/US]; 1403 Ridgewood Drive, San Jose, CA 95118 (US). **XU, Yuming** [US/US]; 1739 Walnut Drive, Mountain View, CA 94040 (US). **SEILHAMER, Jeffrey, J.** [US/US]; 12555 La Cresta, Los Altos Hills, CA 94022 (US). **BOROWSKY, Mark, L.** [US/US]; 122 Orchard Avenue, Redwood City, CA 94061 (US). **KHAN, Farrah, A.** [IN/US]; 3617 Central Road #102, Glenview, IL 60025 (US). **KEARNEY, Liam** [US/US]; 50 Woodside Avenue, San Francisco, CA 94127 (US). **THANGAVELU, Kavitha** [IN/US]; 1950 Montecito Avenue #23, Mountain View, CA 94043 (US). **DAS, Debopriya** [IN/US]; Apartment 13, 1179 Bonita Avenue, Mountain View, CA 94040 (US). **POLICKY, Jennifer, L.** [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US).

(74) Agents: **HAMLET-COX, Diana** et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

[Continued on next page]

(54) Title: TRANSPORTERS AND ION CHANNELS

(57) Abstract: The invention provides human transporters and ion channels (TRICH) and polynucleotides which identify and encode TRICH. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of TRICH.

WO 02/004520 A3



before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(88) Date of publication of the international search report:

20 March 2003

INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 01/21448

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C07K16/18 C12Q1/68 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, WPI Data, EP0-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL [Online] 7 January 2000 (2000-01-07) "Human DNA sequence from clone RP1-137F1 on chromosome 6p21.1-21.2 Contains two genes for novel members of the potassium channel subfamily K (KCNK). Contains ESTs, STSs, GSSs and a CpG island." Database accession no. AL136087 XP002212498 see nts 51861-67180 and complement thereof;/product="dJ137F1.1 (novel member of the potassium channel subfamily K)";/translation="MYRPRARA.....;/db_xref="SPTREMBL:Q9H592"</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1,3,4, 11,12

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

6 September 2002

Date of mailing of the international search report

09. 01. 2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

MADDOX, A

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL [Online] 23 December 1998 (1998-12-23) "CIT-HSP-2384B13.TRB CIT-HSP Homo sapiens genomic clone 2384B13, genomic survey sequence." Database accession no. AQ310967 XP002212499 the whole document	11,12
P,X	WO 00 52164 A (CURTIS RORY A J ;MILLENNIUM PHARM INC (US)) 8 September 2000 (2000-09-08) SEQ ID NOS:5-8 - figure 3	1,3, 6-16,18, 19,22, 25,26
P,X	WO 01 32872 A (SILOS SANTIAGO INMACULADA ;CURTIS RORY A J (US); MILLENNIUM PHARM) 10 May 2001 (2001-05-10) SEQ ID NOS:4-6 figure 3	1,3, 6-16,18, 19,22, 25,26
P,X	DATABASE EMBL [Online] 2 May 2001 (2001-05-02) "Homo sapiens 2P domain potassium channel Talk-2 (KCNK17) mRNA, complete cds." Database accession no. AF358910 XP002212647 the whole document	1-4,11, 12,45
P,X	GIRARD C ET AL: "GENOMIC AND FUNCTIONAL CHARACTERISTICS OF NOVEL HUMAN PANCREATIC 2P DOMAIN K+ CHANNELS" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 1, no. 282, 23 March 2001 (2001-03-23), pages 249-256, XP002908280 ISSN: 0006-291X figure 1A	1-4,9, 11,12,45
X	WO 00 05367 A (KATO SEISHI ;KIMURA TOMOKO (JP); PROTEGENE INC (JP); SAGAMI CHEM R) 3 February 2000 (2000-02-03) SEQ ID NO:68 and 88 page 79, line 32 -page 81, line 26	1-19,22, 25-45,77
X	WO 00 27871 A (CENTRE NAT RECH SCIENT) 18 May 2000 (2000-05-18) figure 14A	1-19,22, 25-45,77

	-/--	

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 26253 A (SMITHKLINE BEECHAM PLC) 11 May 2000 (2000-05-11) SEQ ID NOS:1-4 ---	1-19,22, 25-45,77
E	WO 01 55367 A (HUMAN GENOME SCIENCES INC ;ROSEN CRAIG A (US); BARASH STEVEN C (US) 2 August 2001 (2001-08-02) see SEQ ID NOS:3748 and 881 ---	3,11,12
L	DATABASE GENESEQ [Online] 8 January 2002 (2002-01-08) "Human musculoskeletal system related polynucleotide SEQ ID NO 3748." Database accession no. AAL37383 XP002212648 Cited to indicate SEQ ID NO:3748 of W00155367 the whole document ---	3,11,12
L	DATABASE GENESEQ [Online] 8 January 2002 (2002-01-08) "Human musculoskeletal system related polynucleotide SEQ ID NO 881." Database accession no. AAL35539 XP002212649 Cited to indicate SEQ ID NO:881 of W00155367 the whole document ---	3,11,12
A	WO 00 26245 A (INCYTE PHARMA INC ;AZIMZAI YALDA (US); CORLEY NEIL C (US); YUE HEN) 11 May 2000 (2000-05-11) the whole document -----	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/21448

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
45 and 77 both completely, 1-44 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 01/21448

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

· Invention 1: 45 and 77 both completely,
1-44 all partially

Polypeptide and correpsonding nucleotide as defined by SEQ ID NOS:1 and 33,methods,hosts,compositions and antibodies based on said sequences

Inventions 2-32: Claims 1-108 in so far as is applicable for the sequences as defined below

Polypeptide and corresponding polynucleotide defined by SEQ ID NOS:2-32 and 34-64, each individual polypeptide sequence of 2 through 32 representing an individual invention in combination with the correpsonding polynucleotide sequence of 34 through 64, where invention 2 is represented by SEQ ID NOS:2 and 34 and each subsequent sequential pair representing another invention through to invention 32 represented by SEQ ID NOS:32 and 64;methods,hosts,compositions and antibodies based on said sequences.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 01/21448

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 18 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claim(s) 32 and 34 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Present claims 20, 21, 23, and 24 relate to a product/compound/method defined by reference to a desirable characteristic or property, namely agonists and antagonists of the polypeptide of claim 1.

The claims cover all products/compounds/methods having this characteristic or property, whereas the application does not provide support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound/method by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has not been carried out for claims 20, 21, 23, and 24.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21448

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0052164	A	08-09-2000	AU 3509100 A	21-09-2000
			WO 0052164 A2	08-09-2000
			US 2002028483 A1	07-03-2002
			US 2002045209 A1	18-04-2002

WO 0132872	A	10-05-2001	AU 1356701 A	14-05-2001
			EP 1228205 A2	07-08-2002
			WO 0132872 A2	10-05-2001

WO 0005367	A	03-02-2000	AU 4799499 A	14-02-2000
			CA 2336225 A1	03-02-2000
			EP 1100898 A2	23-05-2001
			WO 0005367 A2	03-02-2000
			JP 2002522018 T	23-07-2002

WO 0027871	A	18-05-2000	EP 1129187 A2	05-09-2001
			WO 0027871 A2	18-05-2000

WO 0026253	A	11-05-2000	EP 1127073 A1	29-08-2001
			WO 0026253 A1	11-05-2000
			US 6426197 B1	30-07-2002

WO 0155367	A	02-08-2001	AU 2950801 A	07-08-2001
			AU 3095801 A	07-08-2001
			AU 3645901 A	07-08-2001
			AU 3646001 A	07-08-2001
			AU 3646101 A	07-08-2001
			AU 3646201 A	07-08-2001
			AU 3646301 A	07-08-2001
			AU 3646401 A	07-08-2001
			AU 3646501 A	07-08-2001
			AU 3646601 A	07-08-2001
			AU 3794301 A	07-08-2001
			AU 3794401 A	07-08-2001
			AU 3794701 A	07-08-2001
			AU 3794901 A	07-08-2001
			AU 3795001 A	07-08-2001
			AU 3795101 A	07-08-2001
			AU 3795201 A	07-08-2001
			AU 3795301 A	07-08-2001
			AU 3795401 A	07-08-2001
			AU 3795501 A	07-08-2001
			AU 3795701 A	07-08-2001
			AU 3795801 A	07-08-2001
			AU 3972601 A	07-08-2001
			AU 3972701 A	07-08-2001
			AU 3972801 A	07-08-2001
			AU 4140201 A	07-08-2001
			AU 4140301 A	07-08-2001
			AU 4140401 A	07-08-2001
			AU 4140501 A	07-08-2001
			AU 4140601 A	07-08-2001
			AU 4140701 A	07-08-2001
			AU 4140801 A	07-08-2001
			AU 4140901 A	07-08-2001
			AU 4141001 A	07-08-2001
			AU 4141101 A	20-08-2001
			AU 4141201 A	07-08-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21448

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0155367	A	AU 4141301 A	07-08-2001
		AU 4141401 A	07-08-2001
		AU 4141501 A	07-08-2001
		AU 4141601 A	07-08-2001
		AU 4141701 A	07-08-2001
		AU 4141801 A	07-08-2001
		AU 4141901 A	07-08-2001
		AU 4313401 A	07-08-2001
		AU 4313501 A	07-08-2001
		AU 4313601 A	07-08-2001
		AU 4313701 A	14-08-2001
		AU 4526201 A	07-08-2001
		AU 4719001 A	07-08-2001
		AU 4719101 A	07-08-2001
WO 0026245	A	11-05-2000	
		AU 1606800 A	22-05-2000
		EP 1144443 A2	17-10-2001
		WO 0026245 A2	11-05-2000